

AD_____

Award Number: W81XWH-07-1-0470

TITLE:

Inhibition of Ras Signaling in Neurofibromatosis.

PRINCIPAL INVESTIGATOR:

Nicolas Nassar, PhD

CONTRACTING ORGANIZATION:

The Research Foundation of SUNY
Stony Brook, NY 11794-3362

REPORT DATE:

July 2010

TYPE OF REPORT:

Final Report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

x Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-07-2010		2. REPORT TYPE Final		3. DATES COVERED (From - To) 15 JUN 2007 - 14 JUN 2010	
4. TITLE AND SUBTITLE Inhibition of Ras Signaling in Neurofibromitosis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-07-1-0470	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Nicolas Nassar, PhD				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The Research Foundation of SUNY Stony Book, NY 11794-3362				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) US Army Medical Research and Material Command Fort Detrick, Maryland 21702- 5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The purpose of the proposed research is to screen for small molecules that interfere with the signaling of Ras in Neurofibromitosis. During this research period, an 'open conformation' of Ras that is non-signaling has been generated. The current hypothesis is that this new conformation is adopted by Ras in the GTP-bound form before it adopts its active effector-binding conformation and therefore is a target for inhibition by small molecules that would attenuate the signaling of hyper-activated Ras in Neurofibromitosis and in other human cancers. Small molecules are being screened for their ability to inhibit or decrease the in vitro binding of Ras to downstream effectors including Raf.					
15. SUBJECT TERMS Ras, neurofibromatosis, cycling, activation, GTP-binding					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	1
Reportable Outcomes.....	2
Conclusion.....	2
References.....	2
Appendices.....	

Introduction and Significance

There is now compelling evidence that aberrant Ras and Ras-like GTPases signaling contributes to many human diseases especially cancer and X-linked neurodegenerative diseases. In these disorders, Ras is constitutively active or GTP-bound either because of a mutation that slows its ability to hydrolyze GTP or because an upstream regulator is constitutively activated (e.g. a membrane-bound receptor such as EGFR). Despite the accumulated knowledge and decades of research, few available drugs directly target Ras or Ras-like GTPases. A need for such a drug(s) is urgent for example to fight Neurofibromatosis where K-Ras is predominantly activated. We argue that the lack of anti-Ras drugs lays in the lack of understanding of the path for Ras cycling between the GTP- and GDP-bound forms. We hypothesize that describing the path for GTP hydrolysis and the path for nucleotide exchange will help design new molecules that stabilize Ras in a non-signaling conformation.

Key Research Accomplishments:

1- To investigate the cycling of Ras between the active and inactive forms (Aim 1). We have continued to study the cycling of Ras between the GTP- and GDP-bound forms by manipulating the flexibility of the switch regions. Recent published work by other investigators suggests that our data especially the ones related to the 'open conformation' of Ras are not an experimental artifact but are transient conformations of Ras along its path between the active and inactive forms. For example, recent molecular dynamics results by the McCammon group (see Lukman *et al.*, 2010; Grant *et al.*, 2009; Gorfe *et al.*, 2008) confirm our approach. Most importantly, the structure of GTP-bound RasT35S in the 'inactive state 1' recently reported by Shima *et al.* (2010) is remarkably similar to the open conformation we found. In the 'open conformation', the switch 1 and 2 domains adopt a conformation that are reminiscent of nucleotide free Ras despite the presence of GDP or GTP. This opening of the switch regions results in the appearance of new pockets on Ras that can be used to dock small molecule inhibitors. Overall, these findings validate our hypothesis and support our approach of screening the open conformation of Ras to find new Ras inhibitors that could eventually be developed into anti-cancer drugs. We are currently running a high throughput screen (HTS) searching a library for small molecules that bind to the open conformation of Ras. Since last report in 2009, we have published our work on the dominant negative form of Ras, RasS17N, (Nassar *et al.*, 2010). We have submitted our molecular dynamics results obtained in collaboration with Dr. Jin Wang (Chemistry, Stony Brook) regarding the reaction of GTP-hydrolysis catalyzed by Ras as investigated using a new-targeted molecular dynamics algorithm for publication (Lu *et al.*, 2011 in review)

2- In addition, we are collaborating with Dr. Francesco Peri (Italy), who has developed a series of Ras inhibitors, in order to improve on the binding and specificity of his compounds.

3- Other studies. In collaboration with Dr. Nick Carpino (Microbiology, Stony Brook University), we are studying the structure-function relationship of the Sts proteins, which act downstream of the T cell receptor (TCR) complex. We previously showed that Sts-1 is the prototype of a new family of protein tyrosine phosphatases (PTPs) (Mikhailik *et al.*, 2007). We have published the crystal structure of the phosphatase domain of Sts-1, Sts-1_{PGM}, in complex with a sulfate moiety (Jankonic *et al.*, 2010). The postdoctoral fellow in charge of the Ras project contributed to demonstrating that Sts-2 regulates the level of tyrosine phosphorylation on targets within T cells, among them the critical T cell tyrosine kinase Zap-70 (Boris *et al.*, 2011).

In parallel, we demonstrated that Sts-1 and Sts-2 are instrumental in down-modulating proteins that are dually modified by both protein tyrosine phosphorylation and ubiquitination (Carpino *et al.*, 2009). Specifically, both naïve and activated T cells derived from genetically engineered mice that lack the Sts proteins display strikingly elevated levels of tyrosine phosphorylated, ubiquitinated proteins following TCR stimulation. The accumulation of the dually modified proteins is transient, and in activated T cells but not naïve T cells is significantly enhanced by co-receptor engagement. These observations hint at a novel regulatory mechanism downstream of the T cell receptor

Publications (total of 4 for this progress period).

1. Nick Carpino, Yunting Chen, Nicolas Nassar and Hye-Won Oh "The Sts proteins target tyrosine phosphorylated, ubiquitinated proteins within TCR signaling pathways" *Mol. Immunol* 46, 3224-3231 (2009).
2. Jakoncic J, Sondgeroth B, Carpino N, Nassar N. "The 1.35 Å resolution structure of the phosphatase domain of the suppressor of T-cell receptor signaling protein in complex with sulfate" *Acta Crystallogr Sect F Struct Biol Cryst Commun*. 2010 Jun 1;66(Pt 6):643-7.
3. San Luis, Boris, Ben Sondgeroth, Nicolas Nassar, and Nick Carpino "STS-2 IS A PHOSPHATASE THAT NEGATIVELY REGULATES ZETA- ASSOCIATED PROTEIN (ZAP)-70 AND T CELL RECEPTOR (TCR) SIGNALING PATHWAYS" in revision *J. Biol. Chem*.
4. Qiang Lu, Nicolas Nassar and Jin Wang "An investigation of Ras catalyzed GTP hydrolysis using classical and quantum targeted molecular dynamics" submitted to *JACS*.