

# Near Surface Phenomena in the Inhomogeneous Coastal Boundary Layer

G. de Leeuw  
TNO Physics and Electronics Laboratory  
P.O. Box 96864  
2509 JG The Hague  
The Netherlands  
phone: +31 70 3740462  
FAX: +31 70 3740654  
email: [deleeuw@fel.tno.nl](mailto:deleeuw@fel.tno.nl)

Grant Number N00014-96-1-0581

## LONG-TERM GOALS

The long-term goals of the research are to understand and assess the effects of atmospheric phenomena on the detection of targets at low altitudes over sea in coastal environments with long range infrared search and track (LR-IRST) systems. Atmospheric propagation effects considered are transmission losses due to aerosols and water vapour, blurring and scintillation due to turbulent fluctuations of air temperature and changes in refractivity due to vertical temperature gradients.

## OBJECTIVES

The objectives of the research performed in the framework of the Grant are:

- to describe the aerosol source function in coastal areas at short fetches;
- to quantify the effect of the surf zone on the production of sea spray aerosol;
- to determine the turbulence and refractivity in the inhomogeneous coastal boundary layer and their effects on imaging of low altitude point targets;
- to improve the description of the aerosol size distribution as function of height and meteorological parameters.

This work is supported by the US Office of Naval Research (ONR) Marine Meteorology program.

## APPROACH

### *Aerosol source function.*

Aerosol particle size distributions were measured during EOPACE (Electro-Optical Propagation Assessment in the Coastal Environment) experiments on piers, boats and ships. A suitable selection of these data will be used to formulate expressions for the aerosol source function, and relate the results to meteorological and wave parameters. In addition, data from a variety of other locations will be used in this effort to cover situations and conditions that are representative for 'typical' scenarios and thus allow for extrapolation of the results to other areas. When bubble size distributions are also available, an attempt will be made to discriminate between bubble-mediated production of aerosols and spume droplet production [De Leeuw et al., 1996].

# Report Documentation Page

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>Near Surface Phenomena in the Inhomogeneous Coastal Boundary Layer</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>TNO Physics and Electronics Laboratory, PO Box 96864, 2509 JG The Hague, The Netherlands,</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

### *Surf-zone production of aerosols.*

During the EOPACE Surf experiments in San Diego (Scripps Pier, La Jolla) and Monterey (Moss Landing Marine Institute), the production of aerosols in the surf zone was quantified from particle size distribution measurements at different levels upwind and downwind from the surf zone. Prior to these measurements, the optical particle counters used in these surf experiments were calibrated and intercompared, to avoid compatibility problems that are frequently observed between un-like instrumentation.

### *Turbulence and refractivity.*

Optical turbulence and refractivity are due to different phenomena, i.e. turbulent fluctuations of the air temperature and variation of the mean air temperature with height. During EOPACE IOP's in Monterey and San Diego, over-water transmission links were established between, respectively, Moss Landing and Monterey (path length 22 km) and between Imperial Beach Pier and the Point Loma submarine base (path length 14 km). Visible and IR sources were mounted on the piers. The turbulent fluctuations of the transmitted signals and the positions of the images were measured with camera systems in both the IR and in the visible. Simultaneously, meteorological and aerosol parameters were measured to support the analysis and understand the observed phenomena, and to test propagation models such as IRBLEM [Forand et al., 1998]. A problem is the atmospheric inhomogeneity in off-shore flow, affecting temperature and humidity profiles, and the presence of waves. Data used include those from buoys along the propagation path, and those recorded using a small boat equipped with meteorological instrumentation and IR sources. The latter allows for the determination of the range dependence of the turbulence and refractivity effects.

## **WORK COMPLETED**

During FY98, TNO-FEL participated in EOPACE IOP8, which took place in Duck (NC) in February 1998. IOP8 was a 'small-scale' experiment involving only 2 of the 'regular' EOPACE participants. The experiments in Duck were regarded as a test for a full-scale EOPACE experiment in 1999, to validate surf experiments on the Californian coast (La Jolla, Monterey, San Diego Bay). The attraction of the Duck site is the 600-m long pier and the supporting measurements of environmental parameters such as waves, bottom topography and meteorological data. During the Duck trial, TNO-FEL only measured profiles of aerosol particle size distributions over the surf with Rotorods, to complement the SPAWAR measurements with optical particle counters. Other important information is the evolution of the aerosol plumes measured with lidar by NRL, and bubble information measured with acoustical techniques by NRL. The TNO-FEL optical bubble measuring system could be used during only a very short time because the instrument was damaged in the violent surf.

The analysis of the earlier EOPACE experiments on surf aerosol production, long range transmission and scintillation, and air mass characterisation has been continued. Results are summarised below.

## **RESULTS**

### *Surf-produced sea spray aerosol.*

During the Surf experiments in La Jolla (1996, 1997) and Monterey (1996), particle size distributions were measured by TNO-FEL at the base of the pier and by University of Sunderland (CMAS) at the end of the pier. From comparison of the results in on- and off-shore winds, empirical relations were

developed for the production of sea spray aerosol over the surf zone [De Leeuw et al., 1997; Neele et al., 1997; 1998a; 1998b; 1998c]. Primarily, only the wind speed was identified as a parameter. Together with Delft Hydrolics, a model was developed that describes the aerosol production over the surf in terms of the wave spectrum measured in deep water and the bottom topography [Neele et al., 1998b; 1998c]. This model is based on a parameterisation of the aerosol mass flux over the surf derived from lidar measurements at a Polish site [Petelski and Chomka, 1996; Chomka and Petelski, 1997]. The surf-production model reasonably predicts the observed aerosol fluxes in many cases, and even the evolution of the fluxes over a period of several hours. However, in other cases the model predictions may be more than one order of magnitude off from the observations. This may be due to the use of inappropriate wave spectra, e.g., for the experiments in Monterey Bay, but also other factors may play a role. It is noted that the parameterisation used in the model needs improvement.

The effect of the surf-produced sea spray aerosol on the extinction was estimated using aerosol transport models, in the framework of a PhD study [Vignati, 1999]. A simple line source model, taking into account diffusion and deposition, showed the horizontal extent of the surf-produced aerosol [De Leeuw et al., 1997a; 1997b; Neele et al., 1998a]. In a following effort, the Coastal Aerosol Transport model (CAT) is now under development, to describe processes in a column advected out over the sea [Vignati, 1999; Vignati et al., 1998]. The aerosol is transported in the vertical by diffusion and gravitational effects, and deposition takes place at the surface. Surface production of sea spray aerosol is accounted for, and the model is initialised with the vertical aerosol distribution at the land-sea transition, i.e. the surf produced aerosol and a continental aerosol distribution. Initial exercises with the model show the relative contributions of the surf-produced and surface-produced sea spray aerosol, and the distances over which the surf aerosol dominates the aerosol concentrations [Vignati et al., 1998]. Preliminary results from a small-scale surf-characterisation experiment in Duck (NC) show that the model may reasonably predict observed plume transport, at least qualitatively.

#### *Transmission and scintillation.*

Transmission and scintillation were measured by TNO-FEL over a 22 km path over Monterey Bay between Monterey and Moss Landing in 1996, and over a 15 km path across San Diego between the Point Loma Subbase and Imperial Beach Pier in 1996 (April and November) and 1997 [De Jong et al., 1996; 1998a; 1998b; 1998c; Zeisse et al., 1997; De Jong and de Leeuw, 1997; Forand et al., 1997; 1998]. These measurements span several seasons with a wide variety of atmospheric conditions. Anomalous transmission effects were observed during all experiments. The base transmission was reasonably well forecasted with MODTRAN, but large excursions were observed with very high transmissions due to refraction and atmospheric lensing which cannot be explained by either MODTRAN or IRBLEM [De Jong et al., 1998c; Forand et al., 1998]. The anomalous transmissions occur over periods varying from minutes to hours. Scintillation was observed at time scales from seconds to minutes, which apparently cannot be explained by current theory [De Jong et al., 1998c].

#### *Air mass characterisation.*

TNO-FEL participated in the air mass characterisation experiments in 1996 and 1997 with aerosol retrievals using AVHRR data, in a strong cooperation with the Naval Postgraduate School (NPS) in Monterey, and lidar measurements of aerosol backscatter profiles. The aerosol optical depths clearly show the variations off the Californian coast, and the lidar profiles show the vertical layering. During the 1996 air mass characterisation experiment, the aerosol optical depth was mainly due to aerosols contained in layers above the marine boundary layer, which appeared very shallow. An elevated layer

was advected off shore in the Santa Ana condition encountered during the experiments. However, elevated aerosol layers are not unusual as demonstrated by the TARFOX data [Veefkind et al., 1998]. The air mass characterisation data are analysed in a strong cooperation with NPS and CMAS, taking into account the meteorological situation [Veefkind et al., 1997; Wash et al., 1997; 1998; Jordan et al., 1998].

## **IMPACT**

The results can be used to assess the effects of the atmosphere on the performance of thermal imagers over sea, and in particular the performance of LR-IRST systems.

## **RELATED PROJECTS**

The EOPACE results of TNO-FEL are exchanged with other EOPACE participants. Cooperations are underway to commonly analyse the EOPACE data. The efforts described above are in conjunction with other projects addressing electro-optical propagation over sea in coastal environments, in part basic research, in part applied research, including work on long-range transmission, IRST and backgrounds. Data are used from other areas, e.g. the North Sea, the North Atlantic, the Mediterranean and the Baltic. These projects are supported by the Netherlands Ministry of Defence, the EU or other funding agencies.

## **REFERENCES**

- Chomka, M. and T. Petelski (1997). Modeling the sea aerosol emission in the coastal zone. *Oceanologia* 39, 211-225.
- De Leeuw, A.M.J. van Eijk, A.I. Flossmann, W. Wobrock, P.G. Mestayer, B. Tranchant, E. Ljungstrom, R. Karlsson, S.E. Larsen, M. Roemer and P.J.H. Builtjes (1996). Transformation and vertical transport of N-compounds in the marine atmosphere. In: Proceedings of EUROTRAC Symposium '96, Eds. P.M. Borrell, P. Borrell, T. Cvitas, K. Kelly and W. Seiler, Computational Mechanics Publ., Southampton, pp. 195-198.
- Petelski, T. and M. Chomka (1996). Marine aerosol fluxes in the coastal zone - BAEX experimental data. *Oceanologia* 38, 469-434.
- Veefkind, J.P., G. de Leeuw, P.A. Durkee, P.B. Russell and P.V. Hobbs (1998). Aerosol optical depth retrieval using ATSR-2 and AVHRR data during TARFOX. Accepted for publication in *J. Geophys. Res.*
- Vignati, E. (1999). Modelling interactions between aerosols and gaseous compounds in the polluted marine atmosphere. PhD thesis. In preparation.

## **PUBLICATIONS**

- De Leeuw, G., F.P. Neele, A.M.J. van Eijk, E. Vignati, M.K. Hill and M.H. Smith (1997a). Aerosol production in the surf zone and effects on IR extinction. In: Propagation and Imaging through the atmosphere (L.R. Bissonnette and C. Dainty, Editors), Proc. SPIE Vol. 3125, 14-27.
- De Leeuw, G., F.P. Neele, E. Vignati, M.K. Hill and M.H. Smith (1997b). Surf effect on sea spray aerosol. Proc. Nordic Aerosol Symposium NOSA 97, Chalmers University of technology, Göteborg, Sweden, 23-24 October 1997.
- De Jong, A.N. et al. (1996). EOPACE transmission measurements Spring '96, preliminary results. TNO report FEL-96-A090.

- De Jong, A.N., and G. de Leeuw (1997). Low elevation transmission measurements at EOPACE, part III: Scintillation effects. In: Propagation and Imaging through the atmosphere (L.R. Bissonnette and C. Dainty, Editors), Proc. SPIE, Vol. 3125, 135-147.
- De Jong, A.N. et al. (1998a). Transmission measurements during EOPACE, November 1996 and August/September 1997, preliminary results. TNO report FEL-97-A269 (January 1998).
- De Jong, A.N., G. de Leeuw, P.J. Fritz, M.M. Moerman (1998b). Long-range transmission at low elevations over the ocean. In: RTO meeting proceedings 1: E-O Propagation, Signature and System Performance Under Adverse Meteorological Conditions Considering Out-Of-Area Operations, paper 15 (12 pp).
- De Jong, A.N., G. de Leeuw and P.J. Fritz (1998c). Scintillation measurements during EOPACE 1996 and 1997 campaigns. In: Propagation and Imaging Through the Atmosphere II (SD10) SPIE, San Diego, July 1998
- Forand, J.L., D. Dion, M. Duffy, S.G. Gathman, K.M. Littfinn, A.N. de Jong, G. de Leeuw and K.L. Davidson (1998). Extensive analysis of low-level IR transmission measurements taken over a 15 km path during EOPACE with IRBLEM. In: Propagation and Imaging Through the Atmosphere II (SD10) SPIE, San Diego, July 1998
- Jordan, M.S., C.H. Wash, J.P. Veefkind and G. de Leeuw (1998). Satellite and ship-based lidar measurements of optical depth during EOPACE. IGARRS'98.
- Neele, F.P., G. de Leeuw, A.M.J. van Eijk, E. Vignati, M. K. Hill and M. H. Smith (1998a). Aerosol production in the surf zone and effects on IR extinction. In: RTO meeting proceedings 1: E-O Propagation, Signature and System Performance Under Adverse Meteorological Conditions Considering Out-Of-Area Operations, paper 24 (11 pp.).
- Neele, F.P., G. de Leeuw, M. Jansen and M. Stive (1998b). Quantitative assessment of surf-produced sea spray aerosol. In: Propagation and Imaging Through the Atmosphere II (SD10) SPIE, San Diego, July 1998
- Neele, F.P., G. de Leeuw, M. Jansen, M. Stive and M.H. Smith (1998c). Aerosol production in the surf zone. *J. Aerosol Sci.* 29, Suppl.1, pp. S181-S182.
- Veefkind, P., G. de Leeuw, K.L. Davidson, M. Jordan, C. Wash, P.H. Durkee, M.K. Hill and M.H. Smith (1997). EOPACE air mass characterisation experiment. In: Propagation and Imaging through the atmosphere (L.R. Bissonnette and C. Dainty, Editors), Proc. SPIE Vol. 3125, 66-76.
- Vignati, E., G. de Leeuw and R. Berkowicz (1998). Aerosol transport in the coastal environment and effects on extinction. In: Propagation and Imaging Through the Atmosphere II (SD10) SPIE, San Diego, July 1998
- Wash, C.H., M. Jordan, P.A. Durkee, P. Veefkind and G. de Leeuw (1997). EOPACE satellite and in situ optical depth studies. BAC'97 Proceedings, San Diego, Dec. 2-4, 1997.
- Wash, C.H., M.S. Jordan, P.A. Durkee, J.P. Veefkind and G. de Leeuw (1998). Satellite and Ship-based LIDAR measurements of Optical Depth during EOPACE. In: RTO meeting proceedings 1: E-O Propagation, Signature and System Performance Under Adverse Meteorological Conditions Considering Out-Of-Area Operations, paper 23 (5 pp).
- Zeisse, C., S. Gathman, D. Jensen, K. Littfin, B. Moison, K. Davidson, P. Frederickson, A. de Jong, P. Fritz, G. de Leeuw, J.L. Forand and D. Dion (1997). Low elevation transmission measurements at EOPACE, Part I: Molecular and aerosol effects. In: Propagation and Imaging through the atmosphere (L.R. Bissonnette and C. Dainty, Editors), Proc. SPIE, Vol. 3125, 109-122.

## **IN-HOUSE/OUT-OF-HOUSE RATIOS**

All work in the framework of the Grant was performed by TNO personnel.