

Geoacoustic Parameters of Marine Sediments: Theory and Experiment

M. J. Buckingham
Marine Physical Laboratory, Scripps Institution of Oceanography
University of California, San Diego
9500 Gilman Drive, La Jolla, California 92093-0238, USA
phone: (858) 534-7977 fax: (858) 534-7461 email: mjb@mpl.ucsd.edu

Grant Number: N00014-07-1-0109

<http://extreme.ucsd.edu/>

LONG-TERM GOALS

There are four long term goals, as follows, in descending order of priority. 1) Characterize fully, from a physics-based model, the inter-relationships that exist between the geo-acoustic properties (*i.e.*, compressional-wave speed and attenuation, shear-wave speed and attenuation, frequency, density, porosity, grain size, grain shape and overburden pressure) of saturated, unconsolidated granular media such as marine sediments. 2) Develop a deep-diving acoustic monitoring system capable of operating to the greatest depths in the ocean. 3) Relate the bubble structure to the acoustic properties of ship wakes. 4) Detect extremely high-energy neutrinos from their acoustic signatures in the ocean.

OBJECTIVES

The scientific objectives of the sediment research may be conveniently divided into five categories. (1) Incorporate the effects of pore fluid viscosity into my analytical theory of wave propagation in saturated granular materials. (2) Continue developing my recently introduced Doppler geo-spectroscopy measurement technique for estimating the geo-acoustic properties of marine sediments using a high-Doppler airborne sound source. (3) Develop analytical and numerical models of wave propagation in a 3-layer waveguide (atmosphere-ocean-sediment) from a moving airborne sound source. (4) Develop inversions, based on the 3-D numerical forward model, for extracting sediment parameters and sub-bottom structure from Doppler geo-spectroscopy data. (5) Identify the relationship between the geometrical properties of individual grains (*e.g.*, size and shape) and the physical properties (*e.g.*, porosity) of the bulk granular material.

The objective of the deep ambient noise project is to develop and deploy a broadband (0.05 - 60 kHz), multi-sensor system capable of monitoring sound to the deepest depths (11,000 km) in the ocean. The system will return depth profiles of the noise level and noise coherence in the vertical.

The objective of the ship-wake research is to relate the strong spatial variability of the void fraction to the acoustic properties of the bubbly wake.

The neutrino detection project is a high-risk, high return venture with Prof. Giorgio Gratta and his research group at Stanford. The objective is to detect extremely high-energy neutrinos, of uncertain extra-terrestrial origin, from their acoustic signatures in the ocean.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2007		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Geoacoustic Parameters Of Marine Sediments: Theory And Experiment				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, San Diego, Marine Physical Laboratory, Scripps Institution of Oceanography, La Jolla, CA, 92093				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT There are four long term goals, as follows, in descending order of priority. 1) Characterize fully, from a physics-based model, the inter-relationships that exist between the geo-acoustic properties (i.e., compressional-wave speed and attenuation, shear-wave speed and attenuation, frequency, density, porosity, grain size, grain shape and overburden pressure) of saturated, unconsolidated granular media such as marine sediments. 2) Develop a deep-diving acoustic monitoring system capable of operating to the greatest depths in the ocean. 3) Relate the bubble structure to the acoustic properties of ship wakes. 4) Detect extremely high-energy neutrinos from their acoustic signatures in the ocean.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

APPROACH

Sediments

1) *Pore-fluid viscosity and the grain-shearing theory of sound waves and shear waves sediments*

In the original version of my grain-shearing (GS) theory of waves in saturated porous materials, effects due to the viscosity of the pore fluid were neglected. Such effects manifest themselves at lower frequencies, below 10 kHz, which is well below the frequency band of many of the published measurements of wave properties of sediments. In an effort to fill the low-frequency gap, one component of the ONR-supported SAX99 experiment, conducted in the northern Gulf of Mexico, off Fort Walton Beach, was aimed at extending the frequency range of sound speed and attenuation data to the band below 10 kHz. The new, lower-frequency measurements returned a sound speed (attenuation) that is somewhat lower (higher) than predicted by the GS theory. To accommodate these observations, the GS theory was extended by incorporating the viscosity of the microscopically thin layer of pore fluid separating contiguous grains. The generalized form has been designated the VGS theory, where the first initial serves as a reminder that viscous effects are now included in the model. At higher frequencies, above 10 kHz, the VGS dispersion curves approach those of the GS theory asymptotically; and at lower frequencies, below 10 kHz, the VGS theory shows a very reasonable match to the SAX99 dispersion data. The VGS theory fits the available dispersion and attenuation data over the approximate frequency range 1 - 400 kHz.

2) *Doppler geo-spectroscopy*

This recently introduced technique for measuring the low-frequency (80 Hz to 1 kHz) sound speed in marine sediments relies on a high-speed airborne sound source (a light aircraft) for ensonifying the ocean and sediment. Most of the received sound is in the form of engine and propeller harmonics, whose frequencies extend from about 80 Hz up to 1 kHz. As the source approaches toward (recedes from) the sensor station, the harmonics are Doppler up-shifted (down-shifted). Normal modes are excited in the water column and are detected on our autonomous line array of 11 non-uniformly spaced hydrophones (known as the “FlyBy array”). A high-resolution spectrum of a single harmonic exhibits sharp peaks, representing the Doppler up- and down-shifted modal field. The magnitude of the modal shifts depends on the speed of the source and the properties of the sea bed. From a precision measurement of the shifted modal frequencies, an inversion returns the sound speed in the sediment; and the remaining geo-acoustic parameters are determined from their correlations with the sound speed, which are known from the VGS theory. Two of my graduate students, Eric Giddens (who now has his Ph.D.) and Melania Guerra, have been directly involved in our Doppler geo-spectroscopy research, along with my engineer, Fernando Simonet. One of my earlier graduate students, Dr. Thomas Hahn, who now holds a faculty position at the University of Miami, assisted us during our participation in SAX04. One of my current graduate students, David Barclay, participated in the Doppler geo-spectroscopy component of the MAKAI experiment, conducted off the western tip of Kauai.

3) *Theoretical models of sound from a moving source in a 3-layer waveguide*

Two analyses of sound in a 3-layer waveguide (atmosphere, ocean and sediment) from a high-Doppler, horizontally moving, unaccelerated airborne source have been developed based on multiple integral transforms in conjunction with appropriate boundary conditions. The first is a 2-D model (horizontal line source perpendicular to the source track), which yields an exact analytical solution for the field in all three layers, including a new 3-layer dispersion relation that takes account of all the Doppler shifts introduced by the moving source. The second is a 3-D model (point source), whose solution is a double wavenumber integral that is evaluated numerically. Both field solutions exhibit interesting fore-aft asymmetries, which appear to be consistent with the geo-spectroscopy experimental observations.

4) *Geoacoustic inversions for Doppler geo-spectroscopy*

Based on a comparison of the predictions from the 3-D, double-wavenumber integral model and the data obtained from our Doppler geo-spectroscopy experiments, a “best match” is obtained, from which the sediment parameters are estimated. Essentially, a cost function is formed and minimized. This work was started by my (then) graduate student, Eric Giddens and is being continued by David Barclay.

5) *Grain shape and sediment porosity*

Of all the physical properties, it could be argued that the porosity of a sediment is the most important in determining its wave properties. The porosity is related to the mean grain size, although not uniquely, suggesting that another parameter is involved, perhaps grain shape or roughness. By examining the shapes of individual grains under a microscope, with careful analysis of the resultant computer-generated images, the relationships between mean grain diameter, rms roughness and bulk-sediment porosity are being investigated. This work is being performed by my graduate student, David Barclay.

Deep ambient noise

This is a new project in which two Vitroflex glass hemispheres, of approximately 0.4 m internal diameter, form a complete sphere containing data acquisition and storage electronics. Throughputs connect external hydrophones and pressure/temperature sensors to the interior of the sphere. As the system descends into the ocean under the influence of gravity at a terminal speed of about 1 m/s, the broadband (0.05 - 60 kHz) ambient noise is detected on two hydrophones, separated vertically by about 0.5 m. At the lowest point in its descent, which may be as deep as 11,000 m, a drop weight is released via a burn wire, and the system returns to the surface under its own buoyancy. Recovery will be achieved with the aid of an Argos antenna, an RF locator beacon and a high-intensity strobe light.

Ship wakes

Aerial photography using a 35mm Canon 5D camera returns high-resolution (12.8 Megapixel), high-dynamic-range (16-bit) digital images of the bubbly wake produced by various types of maritime vessel. In such images, the detailed geometrical shape of the bubbly wake can be observed, as can the pronounced patchy structure of the bubble concentration along the length of the wake. By extracting the spatial statistics of the void fraction distribution from the images, a numerical model of the “acoustic” wake is being constructed, which will be incorporated into 3-D numerical acoustic propagation/scattering models.

Neutrino detection

Particle physicists have theoretical reasons for believing that extremely high energy (non-solar) neutrinos should exist. Since such particles possess mass, they interact with matter, including the ocean, where they deposit energy, causing thermal expansion, which results in a brief burst of sound. Working in conjunction with Prof. Giorgio Gratta and his research group at Stanford, we are using the US Navy’s AUTECH array of bottom-mounted hydrophones off Andros Island, Bahamas in an attempt to detect neutrino sound signatures. AUTECH is located in the Tongue of the Ocean, which is relatively quiet, since it is protected from shipping noise. However, over the bandwidth of the detection system (approximately 50 kHz), there are plenty of sources of ambient noise, including marine mammals. Much of the effort to date has been directed at characterizing this background noise, against which the neutrino signals must be detected.

WORK COMPLETED

Sediments

1) Pore-fluid viscosity and the grain-shearing theory of sound waves and shear waves sediments

The VGS theory has been completed and its predictions compared with the dispersion and attenuation data from SAX99, as shown in Fig. 1. The new theory has been published in the September 2007 issue of *Journal of the Acoustical Society of America*.

2) Doppler geo-spectroscopy

During September 2005, as part of the ONR-sponsored MAKAI experiment, conducted off the west coast of Kauai, we performed Doppler geo-spectroscopy experiments using our Fly-By line array of 11 non-uniformly spaced hydrophones. The array was deployed either horizontally on the seabed, aligned parallel with the depth contours, or vertically in the water column. A light aircraft (Maule MXT7-180 STOL), rented from a local tour-flight company at Lihue airport, was used as the sound source. Multiple low-level flights were made over the sensor station, yielding a variety of interesting data sets. A further set of flying experiments was conducted at a deeper site, close to the Kilo Moana, the research ship participating in MAKAI. At this location the Fly-By array was deployed vertically. Along with the Doppler shifted acoustic signature of the aircraft, the Fly-By array recorded ambient noise at both sites, although most of the time the weather was calm, as a result of which the noise intensity was almost undetectably low.

3) Theoretical models of sound from a moving source in a 3-layer waveguide

Both models are complete and have been published in the October 2006 issue of the *Journal of the Acoustical Society of America*.

4) Geoacoustic inversions for Doppler geo-spectroscopy

Refinements to the geo-acoustic inversion procedure have continued, including the development of a high-resolution, adaptive filtering technique for enhancing the modal peaks in the Doppler-shifted spectrum.

5) Grain shape and sediment porosity

This work is progressing, based on photomicrographs of various types of grains collected from different locations worldwide.

Deep ambient noise

The data acquisition and storage electronics have been designed and fabricated. The circuits have been assembled into the Vitrovex sphere, along with pressure and temperature sensors. Standard ITC hydrophones (with depth rating of only 1000 m) have been installed external to the sphere on a specially designed support frame. The recovery antennas are not yet available, nor are the deep-diving hydrophones. Salt-water tank tests on the stability of the system and its descent rate have been satisfactorily performed. The system works exactly as required.

Ship wakes

Aerial photographs of ship wakes have been taken routinely during flights for other purposes. Our collection of wake photographs is continually growing, including some showing the detailed bubble structure in a wake.

Neutrino detection

Many noise data sets from AUTECH have been analyzed and various features identified in the spectra.

RESULTS

Sediments

1) Pore-fluid viscosity and the grain-shearing theory of sound waves and shear waves sediments

Fig. 1 shows plots of sound speed ratio and sound attenuation as functions of frequency. The symbols in the plots represent data acquired during the SAX99 experiment in the Gulf of Mexico, the blue lines are the predictions of the GS theory and the red lines represent the new VGS theory. Below 10 kHz, the red lines follow the trends of the data very reasonably, in particular matching the relatively high dispersion displayed by Schock's chirp sonar data (red diamonds). Above 10 kHz, the two theories are essentially the same and both match the data satisfactorily.




Fig. 1. Sound speed ratio versus frequency (left) and sound attenuation versus frequency (right)

2) Doppler geo-spectroscopy

Preliminary analysis of the MAKAI data indicates that the low-frequency (≈ 100 Hz) sediment sound speed is in the region of 1650 m/s, consistent with expectations for a water-saturated medium sand.

3) Theoretical models of sound from a moving source in a 3-layer waveguide

Complete solutions have been derived for the 2-D and 3-D fields in a Pekeris waveguide in the presence of source motion. As part of both the 2-D and 3-D analyses, new dispersion relations have been derived, allowing the Doppler frequency shifts of the modes to be determined from straightforward algebraic expressions. [JASA, **120**, 1825-1841 (2006)]

4) Geoacoustic inversions for Doppler geo-spectroscopy

Significantly improved spectral resolution has been achieved using adaptive filtering.

5) *Grain shape and sediment porosity*

Our analysis of the photomicrographs of individual sediment grains yields properties such as cross-sectional area, perimeter and grain shape. From the latter, a Fourier analysis yields a spectrum from which the grain roughness may be deduced. Originally, the whole analysis procedure was quite slow, with each grain taking several hours. Over the past year, the process has been significantly refined, through the development of enhanced software, and now some 500 grains can be analyzed in a morning. In addition, the original measurements of porosity required large sediment samples (bucket size). A technique has now been devised whereby porosity can be measured accurately from a test-tube size sample. This is an important advance, since many of our samples of sand and sediment are quite small.

Deep ambient noise

As the deep ambient noise measurement system has not yet been deployed in the ocean, there are no acoustic data to report.

Ship wakes

An example of the detail that may be seen in high-resolution, aerial photographs of ship wakes is shown in Fig. 2. This particular image was taken from a Cessna 172, with window removed, using a Canon 5D camera with an image-stabilized 28-135mm zoom lens and a circularly polarizing filter to enhance the contrast between the bubbly wake and the bubble-free sea surface. The original 12.8 Megapixel, 16-bit image is of much better quality than the low-resolution version shown here. The spatial distribution of the void fraction in the wake, as seen in Fig. 2, could be the basis of realistic numerical models of acoustic propagation through ship wakes.



Fig. 2. Detail of the wake from the Morgan Foss, a powerful new Dolphin Class harbor tug. Strong vorticity is evident in the wake and the bubble distribution shows rapid spatial variations, with well-defined regions of high and low concentration.

Neutrino detection

We have not seen any neutrinos yet!

IMPACT/APPLICATIONS

My theoretical and experimental work on the wave and physical properties of marine sediments, and dispersive media in general, is broadly based and has gained a following in the ocean acoustics research community. At ASA meetings and international conferences on acoustics, my theories are cited regularly, suggesting that the work is influencing other scientists in their approach to the complicated issues associated with wave propagation in granular materials.

TRANSITIONS

Several research groups in the USA and elsewhere are using the results of my theoretical work in their own programs, including investigators at the Applied Physics Laboratory, University of Washington, the University of Hawaii, NRL Washington D.C., NRL Stennis and in UK government research

laboratories. This includes my work on ambient noise, waves in sediments, acoustic propagation in shallow ocean channels, sound in multi-layer waveguides, and underwater sound fields from high-Doppler, airborne sources.

RELATED PROJECTS

U.S.A.

1. Dr. Michael Richardson, N.R.L., Stennis, and I are continuing to collaborate on the collection and interpretation of sediment data. I am also closely linked to Dr. Eric Thorsos and the group at APL, University of Washington in connection with sediments and other issues.
2. I have been working with Drs. Michael Porter and Martin Siderius, HLS Research, in helping to plan the MAKAI experiment (selection of sites, etc). I am also helping HLS with research on sediments and ambient noise. Recently, I gave a seminar at HLS on my grain-shearing theory of wave propagation in granular materials.
3. Prof. Giorgio Gratta, Stanford, and I are continuing research on the underwater acoustic detection of extremely high energy neutrinos. Acoustic data for this project are being provided by the U.S. Navy's AUTECH range off Andros Island, Bahamas.

Canada

1. Prof. Ross Chapman, University of Victoria, B.C., and I are collaborating on a shallow water experiment aimed at determining low-frequency (80 Hz to 1 kHz) sound speed and attenuation in marine sediments. In particular, we shall try to use the head wave for extracting the required information. For this frequency band, Ross has a low-intensity air gun source and we use an airborne source of opportunity, two completely different ways of exciting the head wave, but which should yield compatible answers.

United Kingdom

1. Prof. Tim Leighton, Institute of Sound and Vibration Research, University of Southampton, and I are discussing several joint research projects on underwater acoustics. These will involve the interchange of graduate students, post-docs and perhaps more senior staff between ISVR and SIO. In May 2006, under a UK initiative known as "SETsquared Collaborative US UK Research Programme", I met Prof. Phil Nelson, Vice Chancellor, University of Southampton, with a view to starting a joint research program between SIO and ISVR. It is not yet clear how this will develop, but under the SETsquared initiative, the UK has funds to supportive such collaborative efforts. (I hold a visiting Professor appointment at ISVR).
2. Dr. Nicholas Pace, University of Bath and I are discussing the possibility of using an airborne source for low-frequency measurements of sediment properties in the Mediterranean.
3. Nathan Price and Gareth Somerset, SEA Ltd. are developing a system for inverting ambient noise measured on a vertical line array to obtain sediment parameters. Their system uses the vertical coherence of ambient noise, as I proposed some years ago, combined with my recent theory of waves in sediments, to yield the majority of sediment properties.

4. Dr Alastair Cowley, DERA, Winfrith is continuing to collaborate with me on phased array techniques applied to acoustic daylight imaging. Several years ago, his team of engineers conducted tests in San Diego Bay using our ADONIS array head of 128 hydrophones with their high-speed beamformer. This phased array system, without the spherical reflector that we used in our original acoustic daylight experiments, yielded recognizable images of targets at ranges of approximately 10 m solely from the acoustic illumination provided by the ambient noise in the ocean.

PUBLICATIONS

Journals & Chapters in Books

1. M. J. Buckingham, "On pore-fluid viscosity and the wave properties of saturated granular materials including marine sediments", *J. Acoust. Soc. Am.*, **122**, 1486-1501 (2007) [published, refereed]
2. M. J. Buckingham and E. M. Giddens, "Theory of sound propagation from a moving source in a three-layer Pekeris waveguide", *J. Acoust. Soc. Am.*, **120**, 1825-1841 (2006) [published, refereed]
3. M. J. Buckingham and E. M. Giddens, "On the acoustic field in a Pekeris waveguide with attenuation in the bottom half-space", *J. Acoust. Soc. Am.*, **119**, 123-142 (2006) [published, refereed]
4. M. J. Buckingham, "Causality, Stokes' wave equation and acoustic pulse propagation in a viscous fluid", *Phys. Rev. E.*, **72**, 026610(9) (2005) [published, refereed].
5. M. J. Buckingham, "Compressional and shear wave properties of marine sediments: comparisons between theory and data", *J. Acoust. Soc. Am.*, **117**, 137-152 (2005) [published, refereed].
6. M. J. Buckingham, "Acoustic remote sensing of the sea bed using propeller noise from a light aircraft," in *Sounds in the Sea: Introduction to Acoustical Oceanography*, edited by H. Medwin, (Cambridge University Press, Cambridge, 2005) pp. 581-597 [published, refereed].
7. P. D. Thorne and M. J. Buckingham, "Measurements of the form function and total scattering cross section for suspensions of sands," *J. Acoust. Soc. Am.*, **116**, 2976-2890 (2004) [published, refereed].
8. M. J. Buckingham, "A three-parameter dispersion relationship for Biot's fast compressional wave in a marine sediment," *J. Acoust. Soc. Am.*, **116**, 769-776 (2004) [published, refereed].
9. M. J. Buckingham, "On the sound field from a moving source in a viscous medium," *J. Acoust. Soc. Am.*, **114**, 3112-3118 (2003) [published, refereed].
10. T. R. Hahn, T. K. Berger, and M. J. Buckingham, "Acoustic resonances in the bubble plume formed by a plunging water jet," *Proc. Roy. Soc. Lond. A*, **459**, 1751-1782 (2003) [published, refereed].
11. M. J. Buckingham, E. M. Giddens, F. Simonet and T. R. Hahn, "Propeller noise from a light aircraft for low-frequency measurements of the speed of sound in a marine sediment," *J. Comp. Acoust.*, **10** (4), 445-464 (2002) [published, refereed].

12. M. J. Buckingham, E. M. Giddens, J. B. Pompa, F. Simonet and T. R. Hahn, "Sound from a light aircraft for underwater acoustics experiments?," *Acta Acust. united with Acust.*, **88** (5), 752-755 (2002) [published, refereed].
13. M. J. Buckingham and M. D. Richardson, "On tone-burst measurements of sound speed and attenuation in sandy marine sediments," *IEEE J. Ocean. Eng.*, **27** (3), 429-453 (2002) [published, refereed].
14. M. J. Buckingham and M. S. Garcés, "Airborne acoustics of explosive volcanic eruptions," *J. Comp. Acoust.*, **9** (3), 1215-1225 (2001) [keynote address, published, refereed].
15. N. G. Lehtinen, S. Adam, G. Gratta, T. K. Berger and M. J. Buckingham, "Sensitivity of an underwater acoustic array to ultra-high energy neutrinos," *Astroparticle Phys.*, **697**, 1-14 (2001) [published, refereed].
16. M. D. Richardson, K. B. Briggs, D. L. Bibee, P. A. Jumars, W. B. Sawyer, D. B. Albert, T. K. Berger, M. J. Buckingham, *et al.*, "Overview of SAX99: environmental considerations," *IEEE J. Ocean. Eng.*, **26** (1), 26-53 (2001) [published, refereed].
17. M. J. Buckingham, "Precision correlations between the geoacoustic parameters of an unconsolidated, sandy marine sediment," *J. Comp. Acoust.*, **9** (1), 101-123 (2001) [published, refereed].
18. M. J. Buckingham, "Wave propagation, stress relaxation, and grain-to-grain shearing in saturated, unconsolidated marine sediments," *J. Acoust. Soc. Am.*, **108**, 2796-2815 (2000) [published, refereed].
19. C. L. Epifanio, J. R. Potter, G. B. Deane, M. L. Readhead, and M. J. Buckingham, "Imaging in the ocean with ambient noise: the ORB experiments," *J. Acoust. Soc. Am.*, **106**, 3211-3225 (1999) [published, refereed].
20. M. J. Buckingham, "Theory of compressional and transverse wave propagation in consolidated porous media," *J. Acoust. Soc. Am.*, **106**, 575-581 (1999) [published, refereed].
21. M. J. Buckingham, "On the phase speed and attenuation of an interface wave in an unconsolidated marine sediment," *J. Acoust. Soc. Am.*, **106**, 1694-1703 (1999) [published, refereed].
24. M. J. Buckingham, "Acoustic daylight imaging in the ocean," in *Handbook on Computer Vision and Applications*, vol. 1, Sensors and Imaging, B. Jähne, H. Haubecker, and P. Geibler, Eds. San Diego: Academic Press, 1999, pp. 415-424 [published, refereed].
25. M. J. Buckingham, "Acoustic pulse propagation in dispersive media," in *New Perspectives on Problems in Classical and Quantum Physics. Part II. Acoustic Propagation and Scattering - Electromagnetic Scattering*, vol. 2, P.P. Delsanto and A.W. Sáenz, Eds. Amsterdam: Gordon and Breach, 1998, pp. 19-34 [published, refereed].
26. M. J. Buckingham, "Theory of compressional and shear waves in fluid-like marine sediments," *J. Acoust. Soc. Am.*, **103**, 288-299 (1998) [published, refereed].

27. G. B. Deane, M. J. Buckingham, and C.T. Tindle, "Vertical coherence of ambient noise in shallow water overlying a fluid seabed," *J. Acoust. Soc. Am.*, **102**, 3413-3424 (1997) [published, refereed].
28. N. M. Carbone, G. B. Deane, and M. J. Buckingham, "Estimating the compressional and shear wave speeds of a shallow-water seabed from the vertical coherence of ambient noise in the water column," *J. Acoust. Soc. Am.*, **103**, 801-813 (1997) [published, refereed].
29. M. J. Buckingham, "Sound speed and void fraction profiles in the sea surface bubble layer," *Appl. Acoust.*, **51**, 225-250 (1997) [published, refereed].
30. M. J. Buckingham and N. M. Carbone, "Source depth and the spatial coherence of ambient noise in the ocean," *J. Acoust. Soc. Am.*, **102**, 2637-2644 (1997) [published, refereed].
31. M. J. Buckingham, "Theory of acoustic attenuation, dispersion, and pulse propagation in unconsolidated granular materials including marine sediments," *J. Acoust. Soc. Am.*, **102**, 2579-2596 (1997) [published, refereed].

Proceedings of Conferences and Workshops

1. M. J. Buckingham, F. Simonet and D. R. Barclay, "Geo-acoustic inversion experiments in shallow water off Kauai using sound from a light aircraft", in *Underwater Acoustic Measurements: Technologies and Results*, edited by J. S. Papadakis and L. Bjørnø (FORTH, Heraklion 2007), pp. 125-131 [published, refereed]
2. M. J. Buckingham, E. M. Giddens and F. Simonet, "Inversion of the propeller harmonics from a light aircraft for the geoacoustic properties of marine sediments", in *Acoustic Sensing Techniques for the Shallow Water Environment: Inversion Methods and Experiments*, edited by A. Caiti, N. R. Chapman, J-P. Hermende and S. M. Jesus (Springer, Dordrecht, 2006) pp. 257-263 [published, refereed]
3. M. J. Buckingham, S. Simonet and D. Barclay, "Doppler spectroscopy and ambient noise inversions in the Makai experiment", Eight European Conference on Underwater Acoustics, Carvoeiro, Portugal, 12-15 June 2006.
4. M. J. Buckingham, "Sediment acoustics measurements using a light aircraft", SA04 Workshop, 22-23 March 2006.
5. M. J. Buckingham, "Inversions for sediment geoacoustic parameters using a high-Doppler airborne sound source", ONR SW Progress Review, MBARI, 14-16 March 2005.
6. M. J. Buckingham, "Inversions for the geoacoustic properties of marine sediments using a high-Doppler, airborne sound source", Seventh International Conference on Theoretical and Computational Acoustics, Hangzhou, Zhejiang, China, 19-23 September 2005 [KEYNOTE].
7. M. J. Buckingham and E. M. Giddens, "Low frequency sound speed measured using a light aircraft as an acoustic source", International Conference on Boundary Influences in High Frequency, Shallow Water Acoustics, University of Bath, U.K., 5-9 September 2005 [INVITED].

8. M. J. Buckingham, E. M. Giddens, F. Simonet and M. Guerra, "Ocean acoustics experiments in the Gulf of Mexico during SAX04 using a light aircraft as a sound source", International Conference on Underwater Acoustics Measurements: technologies and results, Heraklion, Crete, Greece, 28 June - 1 July 2005 [INVITED].
9. M. J. Buckingham, "From air to sea to sediment: aircraft sound for ocean-acoustics experiments", Conference Honouring Prof. John Papadakis on the Occasion of His Retirement, Heraklion, Crete, November 2004 [INVITED].
10. M. J. Buckingham and E. M. Giddens, "A light aircraft as a low-frequency sound source for acoustical oceanography", Proc. Pan Ocean Remote Sensing Conference, Concepcion, Chile, ed. J. Stuardo, December 2004.
11. M. J. Buckingham, "Wave and material properties of marine sediments: theoretical relationships for geoacoustic inversions", Proc. Conference on High Frequency Ocean Acoustics, La Jolla, CA, eds. M. Porter, M. Siderius and W. M. Kuperman, pp. 1-5, March 2004 [INVITED, published, refereed].
12. M. J. Buckingham, E. M. Giddens and F. Simonet, "Inversion of propeller harmonics from a light aircraft for the geoacoustic properties of marine sediments", Proc. 2nd Workshop on Experimental Acoustic Inversion Methods for Assessment of the Shallow Water Environment", eds. J.-P. Hermand, S. Jesus, A. Caiti and R. Chapman, Ischia, Italy, Kluwer, June 2004 [INVITED].
13. M. J. Buckingham, "Rapid environmental area assessment with ambient noise", Proc. 7th European Conference on Underwater Acoustics, ed. D. Simons, Delft, Netherlands, 1, 529-535, July 2004 [INVITED, published, refereed].
14. M. J. Buckingham and E. M. Giddens, "Aeroplane sound in the sea", Proc. 7th European Conference on Underwater Acoustics, ed. D. Simons, Delft, Netherlands, July 2004 [INVITED].
15. M. J. Buckingham, "Light aircraft, propeller noise and Döppler shifts: tools for estimating the geoacoustic properties of marine sediments", 1st Informal MiniWorkshop on Acoustic Cosmic Ray Detection, Physics Dept., Stanford University, 13-14 September 2003 [INVITED by Prof. G. Gratta].
16. M. J. Buckingham, "Light aircraft, propeller noise and Döppler shifts: tools for estimating the geoacoustic properties of marine sediments", Seminar, Dept. of Aerospace and Mechanical Engineering, University of Southern California, October 2003 [INVITED by Prof. H. K. Cheng].
17. M. J. Buckingham, E. M. Giddens, F. Simonet and T. R. Hahn, "Wave properties of sediments determined using the sound of a light aircraft", Proc. International Conference on Sonar-Sensors and Systems (ICONS-2002), 11-13 December 2002, Cochin, India, H. R. S. Sastry (ed.), 1. [KEYNOTE ADDRESS, published, refereed].
18. M. J. Buckingham, E. M. Giddens, J. B. Pompa and F. Simonet, "Preliminary experiments on light-aircraft noise as the source of sound in ocean-acoustic inversion applications", Proc. First International Conference on Inverse Problems: Modelling and Simulation", Fethiye, Turkey, 14-21 July 2002, A. Hasanov (ed.). [INVITED, published, refereed].

19. M. J. Buckingham, E. M. Giddens, J. B. Pompa, F. Simonet and T. R. Hahn, "A light aircraft as a source of sound for performing geo-acoustic inversions of the sea bed", Proc. 6th European Conference on Underwater Acoustics, Gdansk, Poland, 24-27 June 2002, A. Stepnowski (ed.), pp. 465-470. [INVITED, published, refereed].
20. M. J. Buckingham, E. M. Giddens, J. B. Pompa, F. Simonet and T. R. Hahn, "An airborne sound source and an ocean receiver for remote sensing of the sea bed", Proc. 6th Pan Ocean Remote Sensing Conference (PORSEC), Bali, Indonesia, 3-6 September 2002, B. P. Pasaribu, R. Kaswadi, I. W. Nurjaya and J. L. Gaol (eds.), pp. 203-207 [published, refereed].
21. M. J. Buckingham, "Wave propagation in dispersive marine sediments: a memorial to David G. Crighton", Proc. Fifth Int. Conf. Theoretical & Computational Acoustics, Beijing, May 21-25, E. C. Shang (ed.), 7-14 (2002) [KEYNOTE ADDRESS, published, refereed].
22. M. J. Buckingham, "Acoustical oceanography in perspective", Proc. Inst. Acoust. Conf. on Acoustical Oceanography, T. G. Leighton, G. J. Heald, H. D. Griffiths, and G. Griffiths (eds), Bath Univ. Press, **23**, 2, 1-10, (2001) [KEYNOTE ADDRESS, published, refereed].
23. M. J. Buckingham and T. K. Berger, "Low frequency sound from a bubble plume", Proc. 17th Int. Conf. on Acoustics, Rome, Italy, 2-7 September 2001, A. Alippi (ed.), **1**, 2-3 [CLOSING KEYNOTE ADDRESS, published, refereed].
24. M. J. Buckingham, "New theoretical basis for determining the geoacoustic parameters of the seabed," in *Experimental Acoustic Inversion Methods for exploration of the shallow water environment*, A. Caiti, J.-P. Hermand, S.M. Jesus, and M.B. Porter, Eds. Dordrecht: Kluwer, 2000, pp. 195-209. [INVITED, published, refereed].
25. M. J. Buckingham, "Wave propagation in saturated porous media," in *Proceedings of the First International Conference on Acoustics, Noise and Vibration*, vol. **in press**, A. Guran, Ed. Montreal, (2000). [KEYNOTE ADDRESS, refereed].
26. A. Cowley, R.A. O'Leary, F. Simonet, T.K. Berger, and M.J. Buckingham, "Acoustic Daylight imaging in underwater environments," Proc. Fifth European Conference on Underwater Acoustics, Lyons, **1**, 375-380 (2000) [published, refereed].

PATENTS

1. D. N. Riches and M. J. Buckingham, "Improvements in or relating to tilt sensors", Patent No. 3504/78, January 1978.
2. M. J. Buckingham and J. Edwards, "Suspension apparatus", Patent Application No. 8127155, September 1981.
3. M. J. Buckingham, "Improvements in or relating to sonar systems (HARP), Patent Application No. 8200720, January 1982.
4. M. J. Buckingham, "Acoustic imaging in the ocean using ambient noise", Patent Application No. 08/012894, Notice of Allowance issued in February 1994.

5. M. J. Buckingham, "Method and apparatus for measuring the speed and attenuation of sound", 28 November 2003, UCSD Ref. No. SD2004-080-1 [provisional application].

HONORS/AWARDS/PRIZES

1. M. J. Buckingham, Royal Aerospace Establishment, Clerk Maxwell Premium, IERE, U.K., for research on the detection of gravitational radiation (1972).
2. M. J. Buckingham, Royal Aerospace Establishment, A. B. Wood Medal, Institute of Acoustics, U.K. (1982)
3. M. J. Buckingham, Commendation for Distinguished Contributions to Ocean Acoustics at the Naval Research Laboratory, Washington D. C., U.S.A. (1984)
4. M. J. Buckingham, Alan Berman Publication Award from the Naval Research Laboratory, Washington D. C., U.S.A. (1988).
5. M. J. Buckingham, Scripps Institution of Oceanography, Science Writing Award for Professionals in Acoustics from the Acoustical Society of America (December 1997), for the article on "Seeing underwater with background noise", Scientific American, v. 274 (No. 2), 40-44 (1996).
6. M. J. Buckingham, Scripps Institution of Oceanography, Finalist, Discover Magazine Awards for Technological Innovation, June 1998 (Sight category) for pioneering acoustic daylight imaging.
7. M. J. Buckingham, Scripps Institution of Oceanography, Multiple entries in Marquis Who's Who and Strathmore's Who's Who.
8. M. J. Buckingham, Scripps Institution of Oceanography, Technical Program Chair of the 134th Meeting of the Acoustical Society of America, San Diego, California, 1-5 December 1997.
9. M. J. Buckingham, Scripps Institution of Oceanography, Technical Program Chair of the 148th Meeting of the Acoustical Society of America, San Diego, California, 15-19 November 2004.

My graduate students have been awarded four prizes by the Acoustical Society of America:

1. Thomas Berger, First Prize for Best Student Paper, 136th Meeting of the Acoustical Society of America, 12-16 October 1998: "Low-frequency acoustic emissions of a plunging water jet. Part 1: experiment".
2. Thomas Hahn, Third Prize for Best Student Paper, 136th Meeting of the Acoustical Society of America, 12-16 October 1998: "Low-frequency acoustic emissions of a plunging water jet. Part 2: theory".
3. Eric Giddens, First Prize for Best Student Paper, awarded by the Underwater Acoustics and Engineering Acoustics Technical Committees at the 144th Meeting of the Acoustical Society of America, Cancun, Mexico, 2-6 December 2002: "Sound from a light aircraft for underwater acoustic applications".

4. Eric Giddens, First Prize for Best Student Paper, awarded by the Acoustical Oceanography Technical Committee at the 148th Meeting of the Acoustical Society of America, San Diego, California, 15-19 November 2004: “Geoacoustic inversions in shallow water using Doppler-shifted modes from a moving source”.