

AD _____
(Leave blank)

Award Number: W81XWH-08-1-0321

TITLE: A Novel Anti-Beta2-Microglobulin Antibody Inhibition of Androgen Receptor Expression, Survival, and Progression in Prostate Cancer Cells

PRINCIPAL INVESTIGATOR: Wen-Chin Huang, Ph D

CONTRACTING ORGANIZATION: Cedars-Sinai Medical Center
Los Angeles, California 90048

REPORT DATE: May 2010

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: (Check one)

- ☒ Approved for public release; distribution unlimited
- ☐ Distribution limited to U.S. Government agencies only;
report contains proprietary information

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 31-MAY-2010		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 1 MAY 2009 - 30 APR 2010	
4. TITLE AND SUBTITLE A Novel Anti-Beta2-Microglobulin Antibody Inhibition of Androgen Receptor Expression, Survival, and Progression in Prostate Cancer Cells				5a. CONTRACT NUMBER W81XWH-08-1-0321	
				5b. GRANT NUMBER PC073356	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Wen-Chin Huang, Ph.D. Email: Wen-Chin.Huang@cshs.org				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Cedars-Sinai Medical Center Los Angeles, California 90048				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command, Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We previously demonstrated that a key lipogenic transcription factor, sterol regulatory element-binding protein-1 (SREBP-1), regulated androgen receptor (AR) transcriptional expression through AR promoter activity and cell growth in prostate cancer cells. In this annual summary report, we further investigate the molecular mechanism by which SREBP-1 promotes prostate cancer growth and progression, we manipulated permanent SREBP-1 expression in prostate cancer cells. Alteration of SREBP-1 expression leads to regulate AR expression, cell growth, migration and invasion in prostate cancer cells. SREBP-1 also showed to induce fatty acid and lipid formation in prostate cancer cells through increase of fatty acid synthase expression. Additionally, SREBP-1 induced oxidative stress and NADPH oxidase 5 (Nox5) expression in prostate cancer cells. In subcutaneous xenograft mouse models, SREBP-1 significantly increased LNCaP tumor burdens and promoted prostate tumor initiation. These findings reveal that the novel lipogenic mechanism by which SREBP-1 regulating AR activity, fatty acid and lipid biosynthesis, and oxidative stress contributes to prostate cancer growth and progression, and further provides a new potential target to prevent and treat prostate cancer malignancy.					
15. SUBJECT TERMS anti-β2-microglobulin monoclonal antibody; androgen receptor; prostate cancer; sterol regulatory element-binding protein-1.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	20	19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	8
Reportable Outcomes.....	9
Conclusion.....	9
References.....	10
Appendices.....	11


INTRODUCTION:

Prostate cancer progression from an androgen-dependent (AD) to an androgen-independent (AI) state is well recognized clinically as a fatal event. Androgen signaling mediated by the androgen receptor (AR), a ligand-activated transcription and survival factor, is known to play a key role regulating this lethal progression (1, 2). The central molecule of this project is β 2-microglobulin (β 2M). β 2M is a non-glycosylated protein composed of 119 amino acid residues, and the mature (secreted) form contains 99 amino acid residues with a molecular mass of 11,800 Da (3, 4). β 2M associates with the heavy chain of major histocompatibility complex class I (MHC I) on cell surfaces (5). This complex is essential for the presentation of protein antigens recognized by cytotoxic T lymphocytes (6) and serves as a major component of body's immune surveillance mechanism (7). We previously showed that β 2M plays an unexpected role mediating prostate cancer osteomimicry, cell growth, survival and progression (8, 9), and AR expression. In this project, we evaluate the molecular mechanism of AR gene expression at the transcriptional level regulated by β 2M during prostate cancer progression. We also focus on the β 2M-mediated signaling and AR as a therapeutic target using a novel anti- β 2M monoclonal antibody (β 2M mAb) for the treatment of lethal prostate cancer malignancy. There are two specific aims proposed in this project: **Specific Aim 1:** To determine the molecular mechanism by which the β 2M-mediated signaling regulates AR expression in prostate cancer cells. **Specific Aim 2:** To determine the anti-tumor efficacy of β 2M Ab on prostate cancer cells *in vitro* and pre-established prostate tumors in mice *in vivo*.

BODY:

1) Blockade of β 2M downregulated AR and PSA expression in human prostate cancer cells—We

previously reported that β 2M is a growth and signaling-promoting factor for human prostate cancer cells (8). Target β 2M using a sequence specific β 2M siRNA (8) or β 2M Ab (10) greatly inhibited prostate tumor growth and induced cell programmed death via a caspase-9 cascade pathway *in vitro* and *in vivo*. To test if interrupting β 2M from extracellular sources may also affect AR and PSA expression as well as cell growth of prostate cancer cells, we employed a new agent, anti- β 2M monoclonal antibody (β 2M mAb), to neutralize extracellular β 2M and interrupt its downstream signaling. As shown in Fig. 1A and 1B, β 2M mAb (0 to 10 μ g/mL) significantly decreased both mRNA and protein levels of AR and PSA in LNCaP (AD) and C4-2B (AI) cells in a dose-dependent pattern determined by semi-quantitative RT-PCR and Western blot. Considered the specificity of β 2M mAb inhibitory effect, purified β 2M protein could rescue AR and PSA inhibition by



2) Sterol regulatory element-binding protein-1 binding site within the 5'-flanking promoter region of human AR gene is responsible for AR transcriptional activity regulated by β 2M mAb—Subsequently, we sought to characterize the transcriptional mechanism of AR expression regulated by β 2M mAb in prostate cancer cells. A luciferase reporter construct that contained the 5'-flanking human AR (hAR) promoter fragment (-5400 to +580) was transiently transfected into LNCaP and C4-2B cells. Consistent with previous RT-PCR and Western blot results (Fig. 1A and 1B), β 2M mAb (0 to 10 μ g/mL) significantly decreased hAR promoter luciferase activity in a concentration-dependent pattern (Fig. 2A). Purified β 2M protein could restore the inhibition of hAR promoter reporter activity by β 2M mAb as well. Isotype control IgG did not decrease hAR promoter luciferase activity in LNCaP and C4-2B cells. To further identify the responsible *cis*-acting element in



the hAR promoter region, we conducted hAR promoter deletion study. Three deletion constructs of hAR promoter fragment (ΔA , ΔB and ΔC , Fig. 2B) were generated and confirmed the DNA sequence. After transfected into LNCaP and C4-2B cells, $\beta 2M$ mAb significantly inhibited the original hAR, ΔB (deletion of -1100 to -600) and ΔC (deletion of -1600 to -1100) promoter luciferase activities (Fig. 2B). However, $\beta 2M$ mAb did not affect the promoter luciferase activity of the ΔA construct (deletion of -600 to -40, Fig. 2B), and a decrease in the basal promoter luciferase activity was observed only the ΔA construct in LNCaP and C4-2B cells. Control IgG did not significantly change the promoter activities of all these vector constructs. These results suggested that the region A within the hAR promoter fragment may contain the potential *cis*-acting element mediated AR transcriptional activity by $\beta 2M$ mAb. Because the original hAR promoter reporter vector contains approximate 6 kb (from -5400 to +580) in length, we further used a restriction enzyme, *SacI*, to generate a shorter promoter luciferase construct, the hAR/*SacI* vector (2 kb only, deletion of -4700 to -740), and tested this reporter vector activity in LNCaP and C4-2B cells exposed with $\beta 2M$ mAb or IgG. After luciferase activity assay, the basal promoter activity of this hAR/*SacI* vector slightly decreased in LNCaP and C4-2B cells compared with the original hAR promoter (Fig. 2B and 2C), but no significant difference. It implied that the fragment, -4700 to -740, within the hAR promoter region, is not responsible for AR transcriptional activity regulated by $\beta 2M$ mAb in prostate cancer cells.

To further determine the $\beta 2M$ mAb-mediated *cis*-acting factor in the region A, based on the computer databank searched and analyzed, we predicted that three potential *cis*-acting elements in the region A may be responsible for AR transcription by $\beta 2M$ mAb, early growth response gene-1 (EGR-1) binding site (-181 to -170), sterol regulatory element-binding protein-1 (SREBP-1) binding site (-347 to -336) and activator protein-1 (AP-1) binding site (-475 to -465) (Fig. 2C). Subsequently, we generated three deletion constructs which are individually deleted these three transcription factor binding sites from the hAR/*SacI* promoter luciferase vector and tested their reporter activity in prostate cancer cells. Among these three deletion constructs, the promoter luciferase activities of $\Delta EGR-1$ and $\Delta AP-1$ binding site constructs significantly inhibited by $\beta 2M$ mAb similar with the hAR/*SacI* construct activity in LNCaP and C4-2B cells (Fig. 2C). Only a slight drop of promoter luciferase activity was observed in a $\Delta SREBP-1$ binding site construct while treated with $\beta 2M$ mAb, and decreased the basal promoter luciferase activity of this construct (Fig. 2C). Control IgG did not significantly change the promoter reporter activities of all these deletion constructs. These hAR promoter deletion data, taken together, demonstrated that SREBP-1 binding site located within the 5'-flanking hAR




Fig. 3. The role of SREBP-1 in regulating endogenous AR expression in prostate cancer cells. A, $\beta 2M$ mAb decreased the expression of both precursor SREBP-1 (125 kDa) and mature nuclear SREBP-1 (68 kDa) in a concentration-dependent pattern (0 to 10 μ g/mL) in LNCaP and C4-2B cells determined by Western blot. The inhibitory effect of endogenous SREBP-1 expression was restored by pre-incubation of $\beta 2M$ mAb with purified $\beta 2M$ protein. Isotype control IgG (10 μ g/mL) did not affect SREBP-1 expression. $\beta 2M$ mAb did not change the expression of SREBP-2 which is an isoform of SREBP-1. β -actin was used as a loading control. NE, nuclear extracts. B, A sequence-specific siRNA of SREBP-1 decreased the expression of precursor and nuclear SREBP-1 proteins in LNCaP and C4-2B cells. Due to downregulation of SREBP-1 by SREBP-1 siRNA, AR and nuclear AR (NE) expression was also inhibited in LNCaP and C4-2B cells. For the specificity of this siRNA, SREBP-2 expression did not change by SREBP-1 siRNA. Control non-specific siRNA (Con) did not affect SREBP-1, AR and SREBP-2 expression. C, Overexpressing SREBP-1 by using a SREBP-1 expression vector (EV) increased the expression of precursor and nuclear SREBP-1 as well as endogenous AR protein in LNCaP and C4-2B cells. SREBP-1 expression vector did not affect SREBP-2 expression in prostate cancer cells. Con. control empty vector expression vector.

promoter region is important for hAR promoter activity regulated by β 2M mAb.

3) The role of SREBP-1 in regulating AR expression and cell viability in prostate cancer cells—We have shown that β 2M mAb inhibited AR expression through the interaction of SREBP-1 and SREBP-1 binding site in prostate cancer cells. To examine whether β 2M mAb also affected endogenous SREBP-1 expression, we performed Western blot to evaluate precursor and nuclear SREBP-1 protein expression in LNCaP and C4-2B cells. As shown in Fig. 3A, β 2M mAb (0 to 10 μ g/mL) specifically inhibited expression of precursor (125 kDa) and mature nuclear (68 kDa) SREBP-1 proteins in a concentration-dependent manner but did not affect expression of SREBP-2, which is a SREBP-1 isoform. Purified β 2M protein rescued the inhibitory effect of endogenous SREBP-1 expression by β 2M mAb. Control IgG did not decrease SREBP-1 and SREBP-2 expression. Next, to investigate the role of SREBP-1 in regulating AR expression, we conducted the functional studies to knock-down and overexpress SREBP-1 in prostate cancer cells. A sequence-specific siRNA of SREBP-1 caused great decrease of both precursor and nuclear SREBP-1 proteins in LNCaP and C4-2B cells (Fig. 3B). Due to downregulation of SREBP-1, we also observed that expression of total AR and nuclear AR proteins was inhibited by SREBP-1 siRNA in LNCaP and C4-2B cells (Fig. 3B). SREBP-2 expression was not affected by SREBP-1 siRNA. Control non-specific siRNA did not inhibit expression of SREBP-1, SREBP-2 and AR. Conversely, we overexpressed SREBP-1 by using a SREBP-1 expression vector in prostate cancer cells. As an expectation, overexpressing SREBP-1 affected increase of precursor and nuclear SREBP-1 as well as AR protein expression, but not SREBP-2 in LNCaP and C4-2B cells (Fig. 3C).

4) SREBP-1 induces oxidative stress, Nox5 and catalase expression in prostate cancer cells. ROS and Nox (a ROS generator), have been reported to regulate cell proliferation, progression and metastasis, and radiation resistance of prostate cancer cells (11-13). Our cDNA microarray data showed that Nox5 was greatly up-regulated in SREBP-1 overexpressing LNCaP cells compared to control cells (unpublished data). To further determine whether SREBP-1 induces prostate cancer cell proliferation through activation of Nox5 and ROS, we first examined expression of Nox5 in parental LNCaP, control Neo and SREBP-1 overexpressing H1 and H2 (two highest SREBP-1 overexpressing clones) cells. Consistent with the result of cDNA microarray, Nox5 protein increased in H1 and H2 compared to Neo and





Fig. 4. SREBP-1 induces cell proliferation through alteration of ROS, Nox5 and catalase expression in prostate cancer cells. A, SREBP-1 induced Nox5 and inhibited catalase expression in prostate cancer cells (H1 and H2 clones) determined by Western blot. SREBP-1 also increased p-Akt expression, a survival signaling protein. β -actin was used as an internal loading control. B, SREBP-1 increased the levels of hydrogen peroxide (DCF, see Material and Methods) in H1 and H2 prostate cancer cells. The levels of superoxide (DHE) were not significantly changed by SREBP-1. The relative DCF (%), the right panel) and DHE (%), the left panel) were assigned as 100% in LNCaP cells. **, $P < 0.005$, significant differences from Neo cells. Data represent the mean $\pm$ SD of two independent two triplicate experiments. C, DPI, a Nox specific inhibitor and ROS scavenger, inhibited cell proliferation of Neo and H2 prostate cancer cells in a dose-dependent pattern (0 to 5 μ M) during a two-day treatment. H2 cells with high Nox5 expression (Fig. 3A) and ROS status (Fig. 3B) increased the resistance of DPI-mediated suppression of cell proliferation. The relative cell proliferation (%) was assigned as 100% in each cell clones without DPI treatment. **, $P < 0.005$, significant differences from Neo cells at the same dose of DPI. Data represent the mean $\pm$ SD of two independent four replicate experiments.

parental LNCaP cells (Fig. 4A). We also found that SREBP-1 increased p-Akt expression in prostate cancer cells (Fig. 4A), which is involved in prostate cancer cell proliferation, survival and progression (14). Next, we assayed ROS status (the levels of hydrogen peroxide and superoxide) in these SREBP-1 overexpressing prostate cancer cells. SREBP-1 induced the levels of hydrogen peroxide in prostate cancer cells (Fig. 4B, the right panel). However, the levels of superoxide were not significantly changed by SREBP-1 (Fig. 4B, the left panel). Additionally, expression of catalase, a key enzyme of hydrogen peroxide degradation, decreased in SREBP-1 overexpressing H1 and H2 cells (Fig. 4A). Down-regulation of catalase may also cause hydrogen peroxide accumulation in prostate cancer cells. To further investigate if SREBP-1 induces prostate cancer cell proliferation through activation of Nox5 and ROS, a Nox specific inhibitor, DPI, was used to treat with these prostate cancer cells. DPI has also been showed to decrease hydrogen peroxide levels in prostate cancer cells (15). As shown in Fig. 4C, cell proliferations of both Neo and H2 cells were affected by DPI in a concentration-dependent inhibition. However, overexpressing SREBP-1 H2 cells with high Nox5 expression (Fig. 4A) and ROS status (Fig. 4B) increased the resistance of DPI-mediated suppression of cell proliferation (Fig. 4C). These data indicated that SREBP-1 induced prostate cancer cell proliferation via Nox5 and ROS.

5) SREBP-1 promotes tumor initiation, burden and castration resistance of human prostate tumors in mouse subcutaneous xenograft models. Because SREBP-1 induced cell proliferation, *in vitro* migration and invasion of prostate cancer cells, we further compared tumor initiation and growth of control Neo and overexpressing SREBP-1 (H2) LNCaP cells in mice *in vivo*. Remarkably, H2 developed 100% subcutaneous human prostate tumors (8/8) but Neo only formed 50% tumors (4/8) in mice during 8 weeks observation. Furthermore, H2 prostate tumors exhibited a 14-fold increased the growth rate over that of the Neo tumors, as assessed by tumor volumes (Neo: $8.8 \pm 5.0 \text{ mm}^3$ and H2: $124.0 \pm 40.0 \text{ mm}^3$), after 8-week prostate cancer cell inoculation (Fig. 5A). Consistent with previous Western blot results, IHC data showed that H2 tumors highly expressed SREBP-1 (both cytoplasm and nucleus), Nox5 (cell membrane) and AR (nucleus) proteins compared to Neo harvested from mouse subcutaneous tumor specimens using anti-SREBP-1, Nox5 and AR antibodies (Fig. 5B).



KEY RESEARCH ACCOMPLISHMENTS:

- Blockade of $\beta 2M$ using $\beta 2M$ mAb significantly downregulated AR and PSA expression and induced an apoptotic caspase-dependent pathway in prostate cancer cells.
- We identified a *cis*-acting element, SREBP-1 binding site, within the 5'-flanking promoter region of hAR gene is responsible for AR transcriptional activity regulated by $\beta 2M$ mAb.

- We further demonstrated that a transcription factor, SREBP-1, interacting with SREBP-1 binding site within the hAR promoter region mediated by β 2M mAb in prostate cancer cells.
- SREBP-1 plays a key role in regulation of AR expression and cell viability in prostate cancer cells
- SREBP-1 induces oxidative stress, Nox5 and catalase expression in prostate cancer cells.
- SREBP-1 promotes tumor initiation, burden and castration resistance of human prostate tumors in mouse subcutaneous xenograft models.

REPORTABLE OUTCOMES:

During the second year of this DoD geant from May 1, 2009 to April 30, 2010, I have published another peer-reviewed research article in *J Biol Chem* (285: 7947-59, 2010, PMID: PMC2832945) and presented a poster presentation in 2010 AACR annual meeting. Please see the **APPENDICES** section.

CONCLUSION:

β 2M is a signaling and growth-promoting factor inducing prostate cancer cell proliferation, survival and progression. Interrupting β 2M and its related signaling pathways by a novel agent, β 2M mAb resulted in the inhibition of AR and PSA expression and the induction of apoptosis of prostate cancer cells. The molecular mechanism of AR inhibitory expression by β 2M mAb was through decreasing the interaction between a lipogenic transcription factor, SREBP-1, and its binding *cis*-acting element located in the 5'-flanking AR promoter region determined by electrophoretic mobility shift assay (EMSA) and chromatin immunoprecipitation assay (ChIP) (16). The functional study of SREBP-1 revealed that knocked-down or overexpressed SREBP-1 by utilizing a sequence-specific siRNA or an expression vector showed to decrease or increase total and nuclear AR proteins and cell viability in prostate cancer cells. Additionally, SREBP-1 induced oxidative stress and Nox5 expression in prostate cancer cells. In subcutaneous xenograft mouse models, SREBP-1 significantly increased LNCaP tumor burdens and promoted prostate tumor initiation. In summary, β 2M mAb may be a potent and attractive pleiotropic therapeutic agent to inhibit AR expression, cell proliferation, survival and fatty acid and lipid metabolism through down-regulation of a lipogenic transcription factor, SREBP-1, in prostate cancer cells. SREBP-1 provides alternative target for prostate cancer.

References:

1. Dehm SM, Tindall DJ. Regulation of androgen receptor signaling in prostate cancer. *Expert Rev Anticancer Ther* 2005;5(1):63-74.
2. Heinlein CA, Chang C. Androgen receptor in prostate cancer. *Endocr Rev* 2004;25(2):276-308.
3. Gussow D, Rein R, Ginjaar I, *et al.* The human beta 2-microglobulin gene. Primary structure and definition of the transcriptional unit. *J Immunol* 1987;139(9):3132-8.
4. Cunningham BA, Wang JL, Berggard I, Peterson PA. The complete amino acid sequence of beta 2-microglobulin. *Biochemistry* 1973;12(24):4811-22.
5. Pedersen LO, Hansen AS, Olsen AC, Gerwien J, Nissen MH, Buus S. The interaction between beta 2-microglobulin (beta 2m) and purified class-I major histocompatibility (MHC) antigen. *Scand J Immunol* 1994;39(1):64-72.
6. Townsend AR, Rothbard J, Gotch FM, Bahadur G, Wraith D, McMichael AJ. The epitopes of influenza nucleoprotein recognized by cytotoxic T lymphocytes can be defined with short synthetic peptides. *Cell* 1986;44(6):959-68.
7. Seliger B. Strategies of tumor immune evasion. *BioDrugs* 2005;19(6):347-54.
8. Huang WC, Wu D, Xie Z, *et al.* {beta}2-Microglobulin Is a Signaling and Growth-Promoting Factor for Human Prostate Cancer Bone Metastasis. *Cancer Res* 2006;66(18):9108-16.
9. Huang WC, Xie Z, Konaka H, Sodek J, Zhau HE, Chung LW. Human Osteocalcin and Bone Sialoprotein Mediating Osteomimicry of Prostate Cancer Cells: Role of cAMP-Dependent Protein Kinase A Signaling Pathway. *Cancer Res* 2005;65(6):2303-13.
10. Huang WC, Havel JJ, Zhau HE, *et al.* {beta}2-Microglobulin Signaling Blockade Inhibited Androgen Receptor Axis and Caused Apoptosis in Human Prostate Cancer Cells. *Clin Cancer Res* 2008;14(17):5341-7.
11. Lu JP, Monardo L, Bryskin I, *et al.* Androgens induce oxidative stress and radiation resistance in prostate cancer cells through NADPH oxidase. *Prostate Cancer Prostatic Dis* 2010;13(1):39-46.
12. Kumar B, Koul S, Khandrika L, Meacham RB, Koul HK. Oxidative stress is inherent in prostate cancer cells and is required for aggressive phenotype. *Cancer Res* 2008;68(6):1777-85.
13. Brar SS, Corbin Z, Kennedy TP, *et al.* NOX5 NAD(P)H oxidase regulates growth and apoptosis in DU 145 prostate cancer cells. *Am J Physiol Cell Physiol* 2003;285(2):C353-69.
14. Li L, Ittmann MM, Ayala G, *et al.* The emerging role of the PI3-K-Akt pathway in prostate cancer progression. *Prostate Cancer Prostatic Dis* 2005;8(2):108-18.
15. Lim SD, Sun C, Lambeth JD, *et al.* Increased Nox1 and hydrogen peroxide in prostate cancer. *Prostate* 2005;62(2):200-7.
16. Huang WC, Zhau HE, Chung LW. Androgen receptor survival signaling is blocked by anti-{beta}2-microglobulin monoclonal antibody via a mitogen-activated protein kinase/lipogenic pathway in human prostate cancer cells. *J Biol Chem* 2010;285(11):7947-56.

Androgen Receptor Survival Signaling Is Blocked by Anti- β 2-microglobulin Monoclonal Antibody via a MAPK/Lipogenic Pathway in Human Prostate Cancer Cells^{*[5]}

Received for publication, December 7, 2009, and in revised form, January 11, 2010. Published, JBC Papers in Press, January 13, 2010, DOI 10.1074/jbc.M109.092759

Wen-Chin Huang^{‡§1}, Haiyen E. Zhau^{‡§}, and Leland W. K. Chung^{‡§2}

From the [‡]Uro-Oncology Research Program, Department of Medicine, Samuel Oschin Comprehensive Cancer Institute, Cedars-Sinai Medical Center, Los Angeles, California 90048 and the [§]Molecular Urology and Therapeutics Program, Department of Urology and Winship Cancer Institute, Emory University School of Medicine, Atlanta, Georgia 30322

A new *cis*-acting element, sterol regulatory element-binding protein-1 (SREBP-1) binding site, within the 5'-flanking human androgen receptor (AR) promoter region and its binding transcription factor, SREBP-1, was identified to regulate AR transcription in AR-positive human prostate cancer cells. We further characterized the molecular mechanism by which a novel anti- β 2-microglobulin monoclonal antibody (β 2M mAb), shown to induce massive cell death in a number of human and mouse cancer cell lines, interrupted multiple cell signaling pathways in human prostate cancer cells. β 2M mAb decreased AR expression through inactivation of MAPK and SREBP-1. By inactivation of MAPK, β 2M mAb decreased prostate cancer cell proliferation and survival. By inhibition of SREBP-1, β 2M mAb reduced fatty acid and lipid levels, an integral component of cell membrane, cell signaling mediators, and energy metabolism. These results provide for the first time a molecular link between the β 2M intracellular signaling axis mediated by MAPK and SREBP-1 and involving lipid signaling, which collectively regulates AR expression and function. Antagonizing β 2M by β 2M mAb may be an effective therapeutic approach simultaneously targeting multiple downstream signaling pathways converging with MAPK, SREBP-1, and AR, important for controlling prostate cancer cell growth, survival, and progression.

β 2-Microglobulin (β 2M)³ is a co-receptor of a major histocompatibility complex class I antigen. β 2M has been implicated

in the regulation of the host immune mechanism and is essential for the recognition of foreign antigens by T-lymphocytes (1). Recent reports from our laboratory and others assigned additional biological functions to β 2M as a diagnostic and prognostic indicator for multiple myeloma, prostate, and breast cancers (2–5); a growth factor and a signaling molecule (6, 7); a new androgen and androgen receptor (AR) target gene (8); and an attractive new therapeutic target for both liquid (9) and solid (10, 11) tumor malignancies. Blockade of β 2M and its related signaling pathways by sequence-specific siRNA or antibody resulted in the inhibition of AR expression and activity and the induction of extensive prostate cancer cell death *in vitro* as well as prostate tumor regression in immune-compromised mice (7, 10). In addition, anti- β 2M monoclonal antibody (β 2M mAb) has been shown not to significantly affect the growth of normal cells, consistent with experimental observations where transgenic mice with a β 2M deficit had normal organ function and life expectancy (9, 10, 12). Therefore, β 2M and its signaling axis may offer an opportunity for improving the clinical targeting of prostate cancer.

AR is a key growth and survival regulatory transcription factor for androgen target organs during normal development and neoplastic progression. Recognition of the importance of the AR signaling axis, particularly in castration-resistant prostate cancer, has prompted discoveries targeting androgen biosynthetic pathways using abiraterone as an agent for a Phase III trial (13, 14). Novel strategies to target AR directly through AR gene transcription and translation (10) or interfering in the interaction between AR and its co-factors and their downstream functions in prostate cancer cells have also been successfully attempted (15–17). AR activity is regulated by a host of factors including steroid hormones, thyroid hormones, vitamin D₃ (18), insulin-like growth factor I, insulin-like growth factor I receptor, keratinocyte growth factor, epidermal growth factor (19), interleukin-6 (20), and agents elevating and activating intracellular cAMP, G protein-coupled receptors, or a PKA signaling pathway (21, 22). The details of the transcriptional/translational mechanisms regulating AR within cancer cells remain unclear. Previous studies demonstrated that the 5'-flanking region of human AR promoter activity can be reg-

^{*} This work was supported, in whole or in part, by National Institutes of Health Grant P01 CA98912 (to L. W. K. C.). This work was also supported by Department of Defense Grants W81XWH-07-1-0172 (to L. W. K. C.) and W81XWH-08-1-0321 (to W.-C. H.) and Georgia Cancer Coalition-Cancer Research Awards (to W.-C. H.).

^[5] The on-line version of this article (available at <http://www.jbc.org>) contains supplemental text and Fig. S1.

¹ To whom correspondence may be addressed: Uro-Oncology Research Program, Dept. of Medicine, Cedars-Sinai Medical Center, 8750 Beverly Blvd., Atrium 103, Los Angeles, CA 90048. Tel.: 310-423-7378; Fax: 310-423-8543; E-mail: wen-chin.huang@cshs.org.

² To whom correspondence may be addressed: Uro-Oncology Research Program, Dept. of Medicine, Cedars-Sinai Medical Center, 8750 Beverly Blvd., Atrium 103, Los Angeles, CA 90048. Tel.: 310-423-7622; Fax: 310-423-8543; E-mail: leland.chung@cshs.org.

³ The abbreviations used are: β 2M, β 2-microglobulin; mAb, monoclonal antibody; AR, androgen receptor; ChIP, chromatin immunoprecipitation assay; EMSA, electrophoretic mobility shift assay; ERK, extracellular signal-regulated kinase; FAS, fatty acid synthase; hAR, human androgen receptor; MAPK, mitogen-activated protein kinase; PKA, protein kinase A; PSA, prostate-specific antigen; siRNA, small interfering RNA; SREBP-1, sterol regula-

tory element-binding protein-1; PI3K, phosphatidylinositol 3-kinase; reverse transcription; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; EGR, early growth response gene; si β 2M, β 2M siRNA; AP-1, activator protein-1.

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway

ulated by transcription factors Sp1 (23), cAMP-responsive element-binding protein (24), FOXO3a (25), and lymphoid enhancer-binding factor-1/T cell-specific factor (LEF-1/TCF) (26), whose activities are subjected to modulation by several known cell signaling pathways such as cAMP/PKA, PI3K/Akt, MAPK, and Wnt/ β -catenin in prostate cancer cells. In this study, we identified an additional transcription factor, SREBP-1, which affected lipid metabolism and accumulation, as a new downstream transcription factor under regulation by β 2M mAb in prostate cancer cells.

SREBP-1 belongs to the SREBP family, which is a basic helix-loop-helix leucine zipper transcription factor (27, 28). Three major isoforms of SREBP have been identified, SREBP-1a, SREBP-1c, and SREBP-2 (28). SREBP-1 has been determined to regulate genes involved in fatty acid and cholesterol biosynthesis (27, 29), whereas SREBP-2 is more specific in the control of cholesterol metabolism (30). Dysregulation of SREBPs and their downstream regulated genes such as fatty acid synthase (FAS), which has been proposed to be a metabolic oncogene (31, 32), was shown to be involved in the development and progression of prostate cancer (33, 34). The expression of SREBP-1 was observed to be highly elevated in clinical human prostate cancer specimens compared with nontumor prostate tissues, and this may be relevant to androgen-refractory progression (34).

The objective of this study is to determine the pleiotropic β 2M-mediated signaling mechanism by which a novel monoclonal antibody, β 2M mAb, inhibited AR mRNA and protein expression and its transcription activity in AR-positive human prostate cancer cell lines. The results of this study suggest that β 2M regulated multiple growth and survival signaling pathways through the control of transcription factors and their modifiers such as AR, MAPK, and PI3K/Akt (7, 10, 35). In particular, we demonstrated that marked down-regulation of AR as the consequence of targeting β 2M by β 2M mAb was due to the inactivation of a lipogenic transcription factor, SREBP-1, known to be associated with androgen-refractory progression of clinical prostate cancer (34). Accompanying reduction of SREBP-1 expression in prostate cancer cells, β 2M mAb also decreased FAS expression and fatty acid and lipid levels, which are the main components of cell membrane and energy storage. Our data reveal for the first time a lipogenic pathway through MAPK and SREBP-1 that is critical for controlling AR expression, activity, and function in prostate cancer cells.

EXPERIMENTAL PROCEDURES

Prostate Cancer Cell Lines, Cell Culture, and Reagents—The LNCaP (androgen-dependent) human prostate cancer cell line and the LNCaP lineage-derived C4-2B bone metastatic subline (androgen-independent) were cultured in T-medium (Invitrogen) supplemented with 5% fetal bovine serum, 100 IU/ml of penicillin, and 100 μ g/ml of streptomycin. These prostate cancer lines were maintained in 5% CO₂ at 37 °C. β 2M mAb was purchased from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Human SREBP-1 expression vector and SREBP-1 siRNA were obtained from OriGene Technologies, Inc. (Rockville, MD) and Santa Cruz Biotechnology, respectively. The selective inhibitors for signaling pathways of MAPK/ERK, U0126; PI3K,

LY294002; and PKA, *N*-[2-(*p*-bromocinnamylamino)ethyl]-5-isoquinolinesulfonamide (H-89) were purchased from Cell Signaling Technology, Inc. (Beverly, MA).

RT-PCR—Total RNA was isolated from prostate cancer cells using a RNeasy kit (Qiagen). Total RNA was used as the template for RT according to the manufacturer's instructions (Invitrogen). The oligonucleotide primer sets used for PCR analysis of cDNA were as follows: β 2M, 5'-ACGCGTCCGAA-GCTTACAGCATTC-3' (forward) and 5'-CCAAATGCGGC-ATCTAGAAACCTCCATG-3' (reverse); AR, 5'-ATGGCTG-TCATTTCAGTACTCCTGGA-3' (forward) and 5'-AGATGG-GCTTGACTTTCCCAGAAAG-3' (reverse); PSA, 5'-ATG-TGGGTCCCGTTGTCTTCCTCACCTGTC-3' (forward) and 5'-TCAGGGGTTGGCCACGATGGTGTCTCCTT-ATC-3' (reverse); and GAPDH, 5'-ACCACAGTCCATGC-CATCA-3' (forward) and 5'-TCCACCACCCTGTTGC-TGT-3' (reverse). The thermal profiles for β 2M, AR, PSA, and GAPDH cDNA amplification were 25–30 cycles starting with denaturation for 1 min at 94 °C, followed by 1 min of annealing at 64 °C (for β 2M), 61 °C (for AR), 55 °C (for PSA), and 60 °C (for GAPDH), and 1 min of extension at 72 °C. RT-PCR products were analyzed by 1.2% agarose gel electrophoresis. Quantity one-4.1.1 Gel Doc gel documentation software (Bio-Rad) was used for quantification of mRNA expression.

Western Blot Analysis—The cell lysates were prepared from prostate cancer cells as described previously (10). The concentration of protein was determined by the Bradford method using Coomassie Plus protein reagent (Pierce). Western blot analysis was performed by a Novex system (Invitrogen). Primary antibodies against human β 2M, AR, PSA, SREBP-1, SREBP-2, FAS, MAPK/ERK (Santa Cruz Biotechnology), Akt, phospho-Akt (Ser⁴⁷³), and phospho-p44/p42 MAPK (Thr²⁰²/Tyr²⁰⁴) (Cell Signaling Technology) were used. The corresponding secondary antibodies conjugated with horseradish peroxidase were purchased from GE Healthcare. Detection of protein bands was assayed by enhanced chemiluminescence Western blotting detection reagents (GE Healthcare).

Plasmid Construction—A luciferase reporter construct that contained the 5'-flanking region (–5400 to +580) of the full-length human AR (hAR) promoter was kindly provided by Dr. Donald J. Tindall (Mayo Clinic, Rochester, MN). The deletion constructs including, Δ A (deletion of –600 to –40), Δ B (deletion of –1100 to –600), and Δ C (deletion of –1600 to –1100) within the hAR promoter luciferase vector; and Δ EGR-1 binding site (5'-TCGCCCACGCTG-3', –181 to –170), Δ SREBP-1 binding site (5'-CCTCGCCTCCAC-3', –347 to –336), and Δ AP-1 binding site (5'-GCTTGGTCATG-3', –475 to –465) within the hAR/SacI promoter (deletion of –4700 to –740) luciferase vector were generated by a QuikChange site-directed mutagenesis kit (Stratagene, La Jolla, CA). All of the plasmid construct DNA sequences were confirmed by DNA sequencing.

Transfection and Luciferase Activity Assay—LNCaP and C4-2B cells were plated at a density of 1.5×10^5 cells/well in 12-well plates 24 h before transfection. Plasmid DNAs were transfected into cells using Lipofectamine 2000 (Invitrogen) according to the manufacturer's protocol. Each transfection

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway

reaction contained 1.25 μ g of tested DNA construct and 0.25 μ g of a transfection efficiency control cytomegalovirus promoter β -galactosidase plasmid. After 6 h of incubation, DNA-liposome mixtures were replaced by fresh medium without fetal bovine serum. After overnight incubation, the transfected cells were treated with reagents or vehicles. After 24 h of additional incubation, the cells were harvested and lysed in 1 $\times$ reporter lysis buffer (Promega). For luciferase activity assay, 20 μ l of the lysate supernatant was mixed with 100 μ l of the luciferase substrate (Promega) and detected by a luminometer (Monolight 3010 luminometer; PharMingen, San Diego, CA). For β -galactosidase activity assay, 100 μ l of the supernatant was mixed with 100 μ l of 2 $\times$ β -galactosidase substrate (200 mM sodium phosphate buffer, pH 7.3, 2 mM MgCl₂, 100 mM β -mercaptoethanol and 1.33 mg/ml *o*-nitrophenyl- β -D-galactopyranoside) and incubated at 37 °C for 30 min. β -Galactosidase activity was detected by a microplate reader (model 680; Bio-Rad) at 405 nm wavelength. The data were presented as the normalized luciferase activity (the means $\pm$ S.D.) defined as the luciferase activity normalized to internal control β -galactosidase activity. All of the experiments were performed as three independent experiments with duplicate assays.

Electrophoretic Mobility Shift Assay (EMSA)—LNCaP and C4-2B cells were cultured in T-medium with 5% fetal bovine serum until 80% confluence. The cells were then switched to 1-day complete serum-free condition and then treated with β 2M mAb (5 μ g/ml), β 2M mAb preincubated with β 2M or control IgG for an additional 24 h. The nuclear extracts were prepared by a NucBusterTM protein extraction kit (Novagen, San Diego, CA). The specific double-stranded oligonucleotide of SREBP-1 binding site within hAR promoter region used as a probe was 5'-TTCTCCCTCCCTCGCCTCCACCTGTTGGTT-3'. The double-stranded oligonucleotide was end-labeled with [γ -³²P]ATP (3,000 Ci/mmol at 10 mCi/ml) using T4 polynucleotide kinase (New England BioLabs, Beverly, MA). Forty thousand cpm of the labeled probe and 5 μ g of nuclear extracts were incubated with EMSA buffer containing 20 mM HEPES (pH 8.0), 100 mM KCl, 0.2 mM EDTA, 0.5 mM dithiothreitol, 20% glycerol, 500 ng of sonicated salmon sperm DNA, and 1 μ g of poly(dI-dC) at 30 °C for 30 min. The samples were subjected to 6% nondenaturing DNA retardation gel and a Novex TBE system (Invitrogen). For the competition experiment, a ³²P-unlabeled oligonucleotide probe (100 $\times$) was preincubated with nuclear extracts for 30 min at room temperature before the addition of the ³²P-probe. After electrophoresis, the gels were dried with a Gel Dryer (model 583; Bio-Rad) and exposed to BioMax film (Kodak).

Chromatin Immunoprecipitation (ChIP) Assay—ChIP was performed by a ChIP-IT kit (Active Motif, Inc., Carlsbad, CA). Briefly, LNCaP and C4-2B cells were serum-starved for 24 h and then treated with or without β 2M mAb (5 μ g/ml) for an additional 24 h. The formaldehyde-fixed chromatin was prepared from these prostate cancer cells and were sheared into 200–1,500 bp of DNA fragments by enzymatic shearing mixture for 10 min at 37 °C. A portion of the fixed and sheared chromatin fragments was reversed and used as input DNA. The other chromatin DNA fragments were immunoprecipitated by anti-SREBP-1 antibody (Santa Cruz Biotechnology), and DNA

was extracted and purified from the immunoprecipitate. A PCR primer pair to amplify the SREBP-1 binding site is 5'-TGG-CAGCCAGGAGCAGGTATT-3' (forward) and 5'-TTTC-CTGGAGGCCAGCACTCAC-3' (reverse). A negative PCR primer pair included in the ChIP-IT kit was used as a negative control. To further quantify the PCR products of ChIP, we conducted quantitative real time PCR. Each purified DNA sample was mixed with Power SYBR[®] Green PCR Master Mix (Applied Biosystems, Foster City, CA) and a primer pair of the SREBP-1 binding site (see above), and quantitative PCR was performed using an iCycler iQ real time PCR detection system (Bio-Rad) for two independent experiments with duplicate assays. The results were normalized by input (assigned as 1.0-fold without treating β 2M mAb) for each cell line.

Immunohistochemical Staining—Primary anti-AR antibody purchased from Santa Cruz Biotechnology (1:100 dilution) was used. Tissue specimens were deparaffinized, rehydrated, and subjected to pressure-cooking antigen retrieval at 125 °C and 20 p.s.i. for 30 s, 10 min of double endogenous enzyme block, 4 °C for overnight primary antibody reaction, and 30 min of EnVision+ dual link and streptavidin-peroxidase system incubation. The signals were detected by adding substrate hydrogen peroxide using diaminobenzidine as the chromogen and counterstained with hematoxylin. The staining reagents were obtained from Dako Corporation (Carpinteria, CA).

Statistical Analysis—The statistical analyses were performed as described previously (35). Student's *t* test and two-tailed distribution were applied in the analysis of statistical significance.

RESULTS

Blockade of β 2M Down-regulated AR and PSA Expression in Human Prostate Cancer Cells—In support of our previous report (7), we observed that β 2M knockdown can be achieved efficiently by genetic manipulation using β 2M siRNA (si β 2M) in a human prostate cancer cell line, C4-2B. We observed that mRNA levels of β 2M, AR, and PSA were dramatically decreased in β 2M knockdown C4-2B cells compared with parental (P) and control scramble siRNA (Scramble) C4-2B cells (Fig. 1A). In addition to decreased mRNA levels of β 2M, AR, and PSA, endogenous proteins of secreted (from conditioned media) and soluble (from cell lysates) β 2M, AR, and PSA were also greatly reduced in si β 2M C4-2B cells compared with P and Scramble C4-2B cells (Fig. 1B). Furthermore, these results were supported by an *in vivo* subcutaneous C4-2B xenograft mouse model (7) where immunohistochemical staining confirmed that AR expression was greatly decreased in si β 2M-treated C4-2B tumors compared with P and Scramble siRNA-treated C4-2B tumors (Fig. 1C). Mouse serum PSA levels were also markedly decreased in si β 2M-treated C4-2B tumors (1.03 $\pm$ 0.52 ng/ml, *n* = 5, after 28 days of treatment) compared with Scramble siRNA-treated C4-2B tumors (19.70 $\pm$ 9.04 ng/ml, *n* = 5). These *in vitro* and *in vivo* data suggested that blockade of intracellular β 2M by β 2M siRNA greatly inhibited the expression of AR and PSA mRNA and protein in prostate cancer cells.

To test whether interrupting β 2M from extracellular sources may also affect AR and PSA expression as well as cell growth of prostate cancer cells, we employed a new β 2M mAb to neutral-

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway




FIGURE 1. β 2M siRNA inhibited expression of AR and PSA mRNA and protein in prostate cancer cells *in vitro* and *in vivo*. A, si β 2M dramatically decreased expression of β 2M, AR, and PSA mRNA in C4-2B prostate cancer cells analyzed by RT-PCR. Expression of GAPDH was used as a loading control. P, parental nontransfected C4-2B cells; Scramble, control scramble siRNA transfected C4-2B cells. B, si β 2M also markedly inhibited expression of secreted β 2M (from conditioned media, CM), soluble β 2M (from cell lysates), AR, and PSA protein in si β 2M C4-2B cells compared with P and Scramble C4-2B cells assayed by Western blot. β -Actin was used as an internal loading control. C, immunohistochemical staining analysis of AR in P (parental, untreated), si β 2M (β 2M siRNA-treated), and Scramble (control scramble siRNA-treated) C4-2B tumor specimens from subcutaneous mouse xenografts (7). This result revealed that si β 2M greatly inhibited AR expression in C4-2B subcutaneous tumors *in vivo*. Scale bars, 100 μ m.




FIGURE 2. β 2M mAb decreased expression of AR and PSA in prostate cancer cells. A, β 2M mAb decreased the steady-state mRNA levels of AR and PSA in a dose-dependent manner (0–10 μ g/ml) in LNCaP and C4-2B AR-positive prostate cancer cells determined by semi-quantitative RT-PCR. The inhibitory effect was restored by preincubation of β 2M mAb with purified β 2M protein. Isotype control IgG (10 μ g/ml) did not significantly affect AR and PSA mRNA expression. The relative mRNA levels of AR and PSA, normalized by GAPDH mRNA, were measured by Gel Doc gel documentation software (Bio-Rad). The relative mRNA levels (%) were assigned as 100% in the absence of β 2M mAb treatment. *, $p < 0.05$; **, $p < 0.005$, significant differences from the β 2M mAb-untreated group. The data represent the means $\pm$ S.D. of independent triplicate experiments. B, β 2M mAb also inhibited total AR, nuclear AR (NE, nuclear extracts), and PSA protein expression in a dose-dependent pattern (0–10 μ g/ml) in LNCaP and C4-2B cells assayed by Western blot. The inhibitory effect was abrogated by preincubation of β 2M mAb with β 2M protein. Control IgG (10 μ g/ml) did not change AR and PSA protein expression. β -Actin was used as an internal loading control.

ize extracellular β 2M in an attempt to interrupt β 2M downstream signaling. As shown in Figs. 2, β 2M mAb (0 to 10 μ g/ml) significantly decreased both steady-state mRNA levels and protein amounts of AR and PSA in LNCaP and C4-2B cells in a dose-dependent pattern determined by semi-quantitative RT-

PCR and Western blot. Considering the specificity of the inhibitory effect of β 2M mAb, purified β 2M protein could rescue AR and PSA inhibition by β 2M mAb in prostate cancer cells. Control IgG did not affect AR and PSA expression. In addition to decreasing endogenous total AR protein, β 2M mAb also inhibited nuclear AR levels in LNCaP and C4-2B cells (Fig. 2B). These data suggested that antagonizing extracellular β 2M by β 2M mAb reduced AR and PSA transcription and translation in prostate cancer cells.

β 2M mAb Blocked AR Transcription via Down-regulation of Sterol Regulatory Element-binding Protein-1 Activity—AR gene transcription was studied in LNCaP and C4-2B cells by transfecting these cells with either the full-length or the deletion constructs of the 5'-flanking region of hAR promoter luciferase reporters (*i.e.* from –5400 to +580). Consistent with previous RT-PCR and Western blot results (Fig. 2), β 2M mAb (0–10 μ g/ml) significantly decreased the full-length hAR promoter luciferase activity in a concentration-dependent pattern (Fig. 3A), and purified β 2M protein was shown to restore the inhibition. Additionally, isotype-matched control IgG failed to decrease the hAR promoter luciferase activity in LNCaP and C4-2B cells. To further identify the responsible *cis*-acting element in the hAR promoter region, we conducted a hAR promoter deletion study. Three deletion constructs of hAR promoter fragment (Δ A, Δ B, and Δ C; Fig. 3B) were generated and confirmed the DNA sequence. After transfection into LNCaP and C4-2B cells, β 2M mAb significantly inhibited the activities of the full-length hAR, Δ B (deletion of –1100 to –600), and Δ C (deletion of –1600 to –1100) promoter luciferase constructs (Fig. 3B). However, β 2M

mAb did not affect Δ A promoter luciferase activity (deletion of –600 to –40; Fig. 3B). Among the three deletion constructs, decreased basal AR promoter luciferase activity was observed only in the Δ A construct when tested in LNCaP and C4-2B cells. Control IgG did not significantly change the promoter activities of

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway




FIGURE 3. SREBP-1 binding site within the 5'-flanking promoter region of hAR gene is responsible for AR transcriptional activity mediated by β 2M mAb. *A*, β 2M mAb decreased the full-length hAR promoter (−5400 to +580) luciferase activity with a concentration-dependent pattern (0–10 μ g/ml) in LNCaP and C4-2B cells. Purified β 2M protein restored the inhibitory effect of hAR promoter activity regulated by β 2M mAb. Control IgG did not suppress hAR promoter reporter activity. *B*, region A (−600 to −40) is responsible for hAR promoter luciferase activity mediated by β 2M mAb in LNCaP and C4-2B cells. β 2M mAb (10 μ g/ml) significantly decreased the promoter luciferase activities of the deleted region B (Δ B, −1100 to −600) and C (Δ C, −1600 to −1100) in hAR promoter report constructs but did not affect the luciferase activity of the Δ A (−600 to −40) construct. Isotype control IgG (10 μ g/ml) did not significantly change the promoter reporter activities of all deletion constructs. *C*, the DNA sequence of region A (560 bp) contains an EGR-1 binding site (−181 to −170), a SREBP-1 binding site (−347 to −336), and an AP-1 binding site (−475 to −465). Among the three deletion constructs (Δ EGR-1, Δ SREBP-1, and Δ AP-1 binding sites), the promoter luciferase activities of Δ EGR-1 and Δ AP-1 binding site constructs were significantly inhibited by β 2M mAb in LNCaP and C4-2B cells. Only a slight decrease of promoter luciferase activity was observed in a Δ SREBP-1 binding site construct treated with β 2M mAb in prostate cancer cells. Control IgG did not change the promoter reporter activities of these three deletion constructs. All of the promoter luciferase activity data (in Fig. 3) were normalized by internal control β -galactosidase activity and expressed as the means $\pm$ S.D. of three independent duplicate experiments. **, $p < 0.005$.

all these vector constructs. These results suggested that a potential *cis*-acting element mediating AR transcriptional activity by β 2M mAb may reside at region A. Because the full-length hAR promoter reporter construct is ~6 kb in length (from −5400 to +580), we further used a restriction enzyme, SacI, to generate a shorter promoter luciferase construct, a hAR/SacI vector (2 kb only, deletion of −4700 to −740), and tested this new reporter vector activity in LNCaP and C4-2B cells exposed to either β 2M mAb or IgG. The basal luciferase activity of the truncated hAR/SacI promoter vector decreased slightly when compared with the full-length hAR promoter activity assayed in LNCaP and C4-2B cells (Fig. 3, *B* and *C*). These results indicated that the *cis*-elements spanning from −4700 to −740 of the hAR promoter were not responsible for AR transcriptional regulation in human prostate cancer cells.

To determine the precise *cis*-acting elements in region A of the AR promoter responsible for β 2M mAb-mediated regulation, we searched the data base and predicted three potential *cis*-acting elements in this region: the EGR-1 binding site (−181 to −170), SREBP-1 binding site (−347 to −336), and AP-1 binding site

(−475 to −465) (Fig. 3C). Subsequently, we generated three respective deletion constructs and tested their luciferase reporter activities in prostate cancer cells exposed to either β 2M mAb or control IgG. In resemblance to the truncated hAR/SacI luciferase construct, we found that β 2M mAb inhibited Δ EGR-1 and Δ AP-1 but not Δ SREBP-1 binding site deletion construct activities in LNCaP and C4-2B cells (Fig. 3C). Decreased basal promoter luciferase activity was noted in the Δ SREBP-1 binding site deletion construct (Fig. 3C). Control IgG did not significantly change the promoter reporter activities of all deletion constructs. These data, taken together, demonstrated that the SREBP-1 binding site located within the 5'-flanking hAR promoter region is important for hAR promoter activity regulated by β 2M mAb in AR-positive human prostate cancer cells.

Confirmation of Nuclear SREBP-1 Protein Interaction with the SREBP-1 Binding Site within the hAR Promoter by EMSA and ChIP Assay—We conducted EMSA to further validate whether the SREBP-1 transcription factor is a key protein regulating AR transcriptional activity through β 2M mAb in prostate cancer cells. As shown in Fig. 4A, nuclear extracts prepared

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway




FIGURE 4. β 2M mAb inhibited the interaction between SREBP-1 and its binding *cis*-acting element located in the 5'-flanking hAR promoter region in prostate cancer cells. **A**, EMSA. LNCaP cells were exposed to β 2M mAb, control IgG (5 μ g/ml), or vehicle for 24 h in serum-free conditions. The cells were harvested, and nuclear extracts were prepared. EMSA was performed by incubating nuclear extracts with the 32 P-labeled SREBP-1 oligo-DNA probe and Novex TBE system (Invitrogen). *Lane 1*, no nuclear extracts (NE) added. *Lanes 2 and 3*, nuclear SREBP-1-oligo-DNA complex, not affected by control IgG. *Lane 4*, β 2M mAb greatly inhibited this nuclear protein-DNA complex formation. *Lane 5*, purified β 2M protein could rescue the inhibitory binding effect of β 2M mAb. *Lane 6*, this complex was competed off by adding 100-fold 32 P-unlabeled specific SREBP-1 oligo-DNA probe. **B**, ChIP assay. LNCaP and C4-2B cells were treated with or without β 2M mAb (5 μ g/ml) for 24 h. The chromatin and nuclear proteins were cross-linked by formaldehyde and sheared by enzymatic shearing mixture (ChIP-IT kit, Active Motif, Inc.) and subjected to immunoprecipitation (IP) assay by anti-SREBP-1 antibody or without antibody as a negative control. In the PCR product from chromatin DNA fragments immunoprecipitated by anti-SREBP-1 antibody as templates, a predicted single DNA band (116 bp, *top panel*) was amplified and visualized in LNCaP and C4-2B cells. This PCR product was decreased by treatment with β 2M mAb in prostate cancer cells. In addition, quantitative real time PCR of ChIP was performed (*bottom panel*). The results of quantitative PCR showed that β 2M mAb significantly down-regulated the interaction between SREBP-1 and the SREBP-1 binding site in the 5'-flanking hAR promoter, with a 13.3 and 11.0-fold decrease in LNCaP and C4-2B cells, respectively. The quantitative PCR data were normalized by input and assigned as 1.0-fold without β 2M mAb for each prostate cancer cell line. **, $p < 0.005$.

from β 2M mAb-treated LNCaP cells showed a greatly decreased the 32 P-oligo-DNA and SREBP-1 binding complex (*lane 4*) compared with the DNA-nuclear protein complex without β 2M mAb or control IgG treatment in LNCaP cells (*lanes 2 and 3*). Purified β 2M protein was shown to abrogate the complex formation decreased by β 2M mAb exposure (*lane 5*). The specificity of the binding of 32 P-oligo-DNA probe with nuclear SREBP-1 protein in LNCaP cells was shown by the effective competition of 32 P unlabeled oligo-DNA for this binding complex (*lane 6*).

We performed ChIP to investigate whether the interaction between nuclear SREBP-1 and its binding *cis*-acting element is affected by β 2M mAb in the chromatin environment in prostate cancer cells. An expected single DNA band (116 bp) was detected by a PCR primer set to amplify a SREBP-1 binding region within hAR promoter in LNCaP and C4-2B cells, whereas the chromatin DNA fragments immunoprecipitated by anti-SREBP-1 antibody were used as templates (Fig. 4B, *top panel*). The levels of this amplified PCR product were decreased by exposure to β 2M mAb in prostate cancer cells (Fig. 4B). In addition, utilizing real time quantitative PCR as a readout, we observed that β 2M mAb caused significant reduction of the interaction between SREBP-1 and its binding site within the AR promoter region, with 13.3- and 11.0-fold decreases in LNCaP

and C4-2B cells, respectively (Fig. 4B, *bottom panel*). In summary, EMSA and ChIP data confirmed that β 2M mAb inhibited the interaction between nuclear SREBP-1 and its *cis*-acting element within the AR promoter region, which accounts for the down-regulated AR transcriptional activity in prostate cancer cells.

A Triad Relationship among β 2M, SREBP-1, and AR Is Involved in the Regulation of Fatty Acid Levels and Cell Viability in Prostate Cancer Cells—We evaluated the potential triad relationship among the expression of β 2M, SREBP-1, and AR in prostate cancer cells. As shown in Fig. 5A, β 2M mAb (0 to 10 μ g/ml) specifically inhibited expression of precursor (125 kDa) and mature nuclear (68 kDa) SREBP-1 proteins in a concentration-dependent manner but did not affect expression of SREBP-2, which is a SREBP-1 isoform. Purified β 2M protein rescued the inhibitory effect of endogenous SREBP-1 expression by β 2M mAb. Control IgG did not affect SREBP-1 and SREBP-2 expression. A fatty acid biosynthetic oncogene, FAS, which is a downstream target gene of SREBP-1, was also shown to be decreased by β 2M mAb in a dose-dependent inhibition (Fig. 5A). In addition, we observed that β 2M mAb (5 μ g/ml, 24 h treatment) significantly decreased fatty acid levels in LNCaP ($20.8 \pm 2.9\%$) and C4-2B ($26.6 \pm 2.1\%$) cells. To investigate the role of SREBP-1 in regulating AR expression, we conducted functional

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway




FIGURE 5. A triad relationship among β 2M, SREBP-1, and AR expression in prostate cancer cells. *A*, β 2M mAb decreased the expression of precursor SREBP-1 (125 kDa), mature nuclear SREBP-1 (68 kDa), and FAS in a concentration-dependent pattern (0–10 μ g/ml) in LNCaP and C4-2B cells as determined by Western blot. The inhibitory effect of β 2M mAb on endogenous SREBP-1 and FAS expression was restored by preincubation of β 2M mAb with purified β 2M protein. Isotype control IgG (10 μ g/ml) did not affect expression of these proteins. β 2M mAb did not change the expression of SREBP-2, which is an isoform of SREBP-1. β -Actin was used as a loading control. *B*, a sequence-specific siRNA of SREBP-1 decreased the expression of precursor and nuclear SREBP-1 proteins in LNCaP and C4-2B cells. Because of down-regulation of SREBP-1 by SREBP-1 siRNA, total AR, and nuclear AR (NE) expression was also inhibited in LNCaP and C4-2B cells. To test the specificity of SREBP-1 siRNA, SREBP-2 expression was not changed by this siRNA. Control nonspecific siRNA (Con) did not affect SREBP-1, AR, and SREBP-2 expression. *C*, overexpressing SREBP-1 by a SREBP-1 expression vector (EV) increased the expression of precursor and nuclear SREBP-1 as well as endogenous AR protein in LNCaP and C4-2B cells. SREBP-1 EV did not affect SREBP-2 expression. Con, control empty expression vector. *D*, overexpressing or knocking down SREBP-1 by SREBP-1 EV or SREBP-1 siRNA significantly increased or decreased LNCaP and C4-2B cell viability after a 4-day transfection. Empty expression vector (pcDNA 3.1; Invitrogen) or nonspecific siRNA (Con) were used as control groups. The cell numbers (cell viability) were measured by a hemacytometer (Hauser Scientific, Horsham, PA) and crystal violet staining method. *, $p < 0.05$; **, $p < 0.005$, significant differences from the control groups. The data represent the means $\pm$ S.D. of two independent experiments replicated three times.

studies to knock down and overexpress SREBP-1 in prostate cancer cells. A sequence-specific siRNA of SREBP-1 caused a marked decrease of both precursor and mature nuclear SREBP-1 proteins in LNCaP and C4-2B cells (Fig. 5B). Down-regulation of SREBP-1 by SREBP-1 siRNA also inhibited the expression of total AR and nuclear AR proteins in LNCaP and C4-2B cells (Fig. 5B). In testing the specificity of SREBP-1 siRNA, SREBP-2 expression was shown not to be affected by this siRNA. Control nonspecific siRNA did not inhibit expression of SREBP-1, SREBP-2, and AR. Conversely, overexpressing SREBP-1 by a SREBP-1 expression vector increased expression of precursor and nuclear SREBP-1 as well as AR but not SREBP-2 in LNCaP and C4-2B cells (Fig. 5C). In addition, overexpressing or knocking down SREBP-1 significantly increased or decreased cell viability (Fig. 5D) and fatty acid levels (data not shown) in prostate cancer cells. These data are consistent with previous reports and in aggregate reveal that β 2M is a pleiotropic signaling molecule (6, 7, 36) and has a triad relationship with SREBP-1 and AR, which determines the growth and survival of prostate cancer cells. β 2M-mediated signaling is important for the maintenance of SREBP-1 expression and SREBP-1 regulates AR expression and prostate cancer cell growth and survival (Fig. 5, B–D). Likewise, a reciprocal relationship has been reported between AR and β 2M in which androgens and AR

regulated β 2M expression (8) and β 2M mediated AR expression and prostate cancer cell growth and survival (Figs. 1 and 2) (10).

The Involvement of the MAPK/ERK Signaling Pathway in β 2M mAb Inhibition of SREBP-1 and AR Expression in Prostate Cancer Cells—To determine the signaling mechanism by which β 2M mAb inhibited SREBP-1 and AR expression in prostate cancer cells, we examined the β 2M mAb-mediated status of MAPK/ERK and PI3K/Akt signaling pathways, which have been reported to regulate SREBP-1 and AR expression (25, 37, 38). The expression levels of both the phosphorylated and total proteins of Akt and MAPK were analyzed by Western blot. As shown in Fig. 6A, β 2M mAb decreased expression of phospho-Akt (Ser⁴⁷³) and phospho-MAPK (Thr²⁰²/Tyr²⁰⁴) proteins in LNCaP and C4-2B cells within 2 h in a time-dependent pattern. However, β 2M mAb did not suppress total Akt and MAPK expression. We also observed that AR and SREBP-1 proteins were not decreased by β 2M mAb within 2 h of treatment (Fig. 6A). To further confirm the responsible signaling pathways regulating SREBP-1 and AR expression, we evaluated the effect of U0126 (a MAPK selective inhibitor), LY294002 (a PI3K selective inhibitor), and H-89 (a PKA selective inhibitor) on prostate cancer cells. β 2M mAb and U0126 greatly inhibited expression of SREBP-1 and AR in LNCaP cells (Fig. 6B). FAS was also

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway




FIGURE 6. β 2M mAb inhibited SREBP-1 and AR expression through a MAPK/ERK signaling pathway in prostate cancer cells. *A*, β 2M mAb (10 μ g/ml) decreased phospho-Akt (p-Akt, Ser⁴⁷³) and phospho-MAPK (p-MAPK, Thr²⁰²/Tyr²⁰⁴) protein expression but did not inhibit total Akt and MAPK (p44/p42) protein expression in LNCaP and C4-2B cells at different time points (0, 15, 30, 60, and 120 min) as assayed by Western blot. AR and SREBP-1 protein expression was not affected by β 2M mAb within 2 h of treatment. *B*, β 2M mAb (5 μ g/ml) and U0126 (10 μ M, a MAPK selective inhibitor) greatly inhibited expression of endogenous SREBP-1, AR, and FAS proteins in LNCaP cells with 24 h of treatment as determined by Western blot. H-89 (1 μ M, a PKA selective inhibitor) slightly decreased SREBP-1, AR, and FAS expression. However, LY294002 (20 μ M, a PI3K selective inhibitor) appeared to increase SREBP-1 and AR expression. β -Actin was used as an internal loading control.

decreased by U0126 treatment. H-89 slightly decreased the expression of these proteins. However, LY294002 appeared to enhance SREBP-1 and AR expression in LNCaP cells. Interrupting the PI3K/Akt signaling pathway with LY294002 increased AR, and PSA protein expression in LNCaP cells has been reported (25). Similar results were observed when C4-2B cells were treated with these signaling pathway inhibitors (data not shown). These data suggested that a MAPK/ERK signaling pathway may play a dominant role in the regulation of SREBP-1 and AR expression through β 2M mAb in prostate cancer cells.

DISCUSSION

This study investigated the pleiotropic signaling functions of β 2M, which confer growth, survival, and metastasis benefits to prostate cancer cells. We focus here on the critical role of β 2M in the regulation of AR through SREBP-1, which plays a key role in lipid homeostasis, regulating its downstream target genes, such as FAS, and accumulation of fatty acids and lipid droplets (supplemental Fig. S1) by Oil Red O staining (39) in prostate cancer cells. We identified a triad relationship among β 2M, SREBP-1, and AR. In response to the β 2M-mediated cell signaling, SREBP-1 regulated AR expression by altering AR gene transcription. Conversely, androgens and AR were also found to mediate SREBP-1 expression reciprocally in androgen-responsive prostate cancer cells (40, 41). These data, taken together with previous reports from our laboratory and others that documented the regulatory role of β 2M on AR expression (10) and also androgens and AR regulation of β 2M expression (8), support the triad relationship among β 2M, SREBP-1, and AR.

There are several important clinical implications of this triad relationship. 1) β 2M could be an important driver modulating SREBP-1 and AR expression in prostate cancer cells. It has been shown that upon androgen-refractory progression of human prostate cancer, dysregulated expression of β 2M (2, 7, 42), SREBP-1 (34), and AR (43, 44) are observed. By employing β 2M mAb as a therapeutic agent, we and others found that this antibody caused massive cell death in human prostate and renal cancers (10, 11), multiple myeloma, leukemia, and lymphoma (9) without affecting the growth of normal cells. We propose that the inhibitory action of β 2M mAb could act via inhibition of SREBP-1 expression, which is linked to FAS expression and lipogenic pathways that are known to regulate cell membrane integrity, energy metabolism, lipid raft-mediated signaling in cancer cells (27, 29, 45, 46), and AR expression, which is regarded as a growth and survival factor for human prostate cancer cells (43). 2) The pleiotropic cell signaling network activated by β 2M could have a mediatory action on lipid metabolism and storage and lipid raft-directed cell signaling pathways. In this study, we showed that β 2M mAb inhibited a large number of cell signaling networks, including MAPK, SREBP-1, AR, and PI3K/Akt. It is conceivable that these signaling networks are interconnected through lipid raft complexes (47, 48). The inhibitory action exerted by β 2M mAb could affect the lipid composition of the raft structures, hence altering domain interactions and how downstream cell signal networks can be assembled and interact in a coordinated manner (45, 49). 3) β 2M-regulated downstream signaling is highly dynamic and could affect the function of cells without AR expression. β 2M mAb was shown to block the growth and downstream signaling of cancer but not normal cells regardless of their endogenous levels of AR (10), implying that AR function is not obligatory for the triad relationship. These observations also suggest that AR function could be bypassed by other redundant cell signaling networks mediated by soluble factors such as insulin-like growth factor I, epidermal growth factor, keratinocyte growth factor, and interleukin-6 (19, 20, 50, 51). Additional studies may be warranted to define how the triad relationship would function in normal *versus* cancer cells and in clinical prostate cancer, which characteristically contains cells with heterogeneous arrays of AR, including AR gene amplification and mutation and AR protein overexpression and silencing (52–54). In addition, the data presented in this communication are collected from the study of established AR-positive human prostate cancer cell lines, and further investigation of this concept in primary prostate cancer cells might be of importance.

The precursors of SREBP family proteins are endoplasmic reticulum membrane-anchored with the mature amino-terminal forms being translocated to the cell nucleus responsible for the activation of their target genes (27). Blockade of SREBP-1, a crucial transcriptional regulator for fatty acid and lipid biosynthesis, by β 2M mAb inhibited expression of both precursor and mature nuclear SREBP-1 and FAS (Fig. 5A) in prostate cancer cells and interrupted cell growth and promoted apoptosis (10). We also observed that β 2M mAb significantly decreased fatty acid contents and lipid droplet accumulation in LNCaP and C4-2B cells. However, the signaling mechanism regulating SREBP-1 and its downstream target gene expression by β 2M

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway




FIGURE 7. Proposed mechanism for β 2M mAb inhibition of AR and cell proliferation, survival, and progression through a MAPK/SREBP-1 signaling pathway in prostate cancer cells. β 2M mAb inhibited p-MAPK and p-AKT expression and decreased SREBP-1 expression in LNCaP and C4-2B cells. By the addition of U0126, a MAPK selective inhibitor, we observed down-regulation of SREBP-1 expression (see Fig. 6B). In addition, β 2M mAb inhibited AR and FAS expression; the latter is known to regulate fatty acid biosynthesis, cell membrane integrity, energy metabolism, and lipid raft-regulated cell signaling (31, 32, 46), which ultimately control prostate cancer cell proliferation, survival, and progression.

mAb in prostate cancer is still unclear. Reports in the literature suggest that SREBP-1 is induced by PI3K/Akt and/or MAPK signaling pathways in liver cells, macrophages, and mammary epithelial and breast cancer cells (38, 55, 56). The inhibitors of MAPK and PI3K signaling pathways were demonstrated to down-regulate SREBP-1 and FAS expression and inhibit fatty acid synthesis in MCF-7 and HCT166 carcinoma cells (38). In LNCaP and C4-2B prostate cancer cells, we observed that U0126 (a MAPK selective inhibitor) inhibited SREBP-1, FAS, and AR protein expression (Fig. 6B), similar to the action of β 2M mAb. However, LY294002 (a PI3K selective inhibitor) increased SREBP-1 and AR protein expression in AR-positive prostate cancer cells. Blocking the PI3K/Akt signaling pathway by LY294002 in LNCaP cells resulted in induced AR expression through activation of a Forkhead transcription factor, FOXO3a (25). H-89, a PKA selective inhibitor, did not significantly affect SREBP-1 expression (Fig. 6B). In addition, we showed that β 2M mAb decreased p-MAPK protein expression in LNCaP and C4-2B cells (Fig. 6A), and this inhibition coincides with decreased SREBP-1 and AR expression (Figs. 2 and 5A). Our data collectively suggest that MAPK signaling is the dominant pathway that positively controls SREBP-1, AR, and FAS expression in AR-positive prostate cancer cells and that this cell signaling network may be different in cancer and normal cells.

In conclusion, these results show for the first time that: 1) interrupting intracellular or extracellular β 2M using sequence-specific siRNA or mAb resulted in decreased expression of AR and PSA at the transcriptional and translational levels in AR-positive prostate cancer cells; 2) β 2M mAb inhibited AR expression by blocking a MAPK signaling pathway, decreasing the expression of precursor and nuclear SREBP-1 and reducing the binding between SREBP-1 and its *cis*-element binding site in the hAR promoter region in LNCaP and C4-2B cells; 3) β 2M mAb decreased fatty acid and lipid accumulation by inhibition

of SREBP-1 and FAS expression and may disrupt cell membrane integrity, intracellular lipid raft-mediated cell signaling and energy metabolism; and 4) functional studies of SREBP-1 demonstrated that SREBP-1 regulated endogenous AR expression and cell viability in LNCaP and C4-2B cells. Collectively, these results suggest that β 2M mAb is a novel therapeutic antibody capable of inhibiting the pleiotropic cell signaling network converging on β 2M in prostate cancer cells (Fig. 7). This study also expands our understanding of the key regulatory role of β 2M, which controls intracellular signaling pathways through the regulation of transcription factors such as cAMP-responsive element-binding protein (7, 57), hypoxia inducible factor-1 α (HIF1- α) (57), and SREBP-1.

Acknowledgments—We thank Dr. Donald J. Tindall (Mayo Clinic, Rochester, MN) for providing the hAR promoter luciferase reporter vector and Gary Mawyer for editing the manuscript.

REFERENCES

1. Townsend, A. R., Rothbard, J., Gotch, F. M., Bahadur, G., Wraith, D., and McMichael, A. J. (1986) *Cell* **44**, 959–968
2. Abdul, M., and Hoosain, N. (2000) *Urol. Oncol.* **5**, 168–172
3. Klein, T., Levin, I., Niska, A., Koren, R., Gal, R., Schachter, J., Kfir, B., Narinski, R., Warchaizer, S., and Klein, B. (1996) *Eur. J. Immunogenet.* **23**, 417–423
4. Bataille, R., Grenier, J., and Combes, T. (1988) *Cancer Invest.* **6**, 271–277
5. Tsimberidou, A. M., Kantarjian, H. M., Wen, S., O'Brien, S., Cortes, J., Wierda, W. G., Koller, C., Pierce, S., Brandt, M., Freireich, E. J., Keating, M. J., and Estey, E. H. (2008) *Clin. Cancer Res.* **14**, 721–730
6. Nomura, T., Huang, W. C., Zhau, H. E., Wu, D., Xie, Z., Mimata, H., Zayzafoon, M., Young, A. N., Marshall, F. F., Weitzmann, M. N., and Chung, L. W. (2006) *Clin. Cancer Res.* **12**, 7294–7305
7. Huang, W. C., Wu, D., Xie, Z., Zhau, H. E., Nomura, T., Zayzafoon, M., Pohl, J., Hsieh, C. L., Weitzmann, M. N., Farach-Carson, M. C., and Chung, L. W. (2006) *Cancer Res.* **66**, 9108–9116
8. Yoon, H. G., and Wong, J. (2006) *Mol. Endocrinol.* **20**, 1048–1060
9. Yang, J., Qian, J., Wozniak, M., Wang, S., Lin, P., Wang, M., Yaccoby, S., Kwak, L. W., Barlogie, B., and Yi, Q. (2006) *Cancer Cell* **10**, 295–307
10. Huang, W. C., Havel, J. J., Zhau, H. E., Qian, W. P., Lue, H. W., Chu, C. Y., Nomura, T., and Chung, L. W. (2008) *Clin. Cancer Res.* **14**, 5341–5347
11. Nomura, T., Huang, W. C., Seo, S., Zhau, H. E., Mimata, H., and Chung, L. W. (2007) *J. Urol.* **178**, 292–300
12. Koller, B. H., Marrack, P., Kappler, J. W., and Smithies, O. (1990) *Science* **248**, 1227–1230
13. Chen, Y., Clegg, N. J., and Scher, H. I. (2009) *Lancet Oncol.* **10**, 981–991
14. Attard, G., Reid, A. H., Olmos, D., and de Bono, J. S. (2009) *Cancer Res.* **69**, 4937–4940
15. Molife, L. R., Attard, G., Fong, P. C., Karavasili, V., Reid, A. H., Patterson, S., Riggs, C. E., Jr., Higano, C., Stadler, W. M., McCulloch, W., Dearnaley, D., Parker, C., and de Bono, J. S. (2010) *Ann. Oncol.* **21**, 109–113
16. Taplin, M. E. (2008) *Expert Rev. Anticancer Ther.* **8**, 1495–1508
17. Chang, C. Y., and McDonnell, D. P. (2005) *Trends Pharmacol. Sci.* **26**, 225–228
18. Zilliacus, J., Wright, A. P., Carlstedt-Duke, J., and Gustafsson, J. A. (1995) *Mol. Endocrinol.* **9**, 389–400
19. Culig, Z., Hobisch, A., Cronauer, M. V., Radmayr, C., Trapman, J., Hittmair, A., Bartsch, G., and Klocker, H. (1994) *Cancer Res.* **54**, 5474–5478
20. Hobisch, A., Eder, I. E., Putz, T., Horninger, W., Bartsch, G., Klocker, H., and Culig, Z. (1998) *Cancer Res.* **58**, 4640–4645
21. Kasbohm, E. A., Guo, R., Yowell, C. W., Bagchi, G., Kelly, P., Arora, P., Casey, P. J., and Daaka, Y. (2005) *J. Biol. Chem.* **280**, 11583–11589
22. Sadar, M. D. (1999) *J. Biol. Chem.* **274**, 7777–7783
23. Takane, K. K., and McPhaul, M. J. (1996) *Mol. Cell. Endocrinol.* **119**, 83–93
24. Mizokami, A., Yeh, S. Y., and Chang, C. (1994) *Mol. Endocrinol.* **8**, 77–88

β 2M mAb Inhibited AR through a MAPK/SREBP-1 Pathway

25. Yang, L., Xie, S., Jamaluddin, M. S., Altuwaijri, S., Ni, J., Kim, E., Chen, Y. T., Hu, Y. C., Wang, L., Chuang, K. H., Wu, C. T., and Chang, C. (2005) *J. Biol. Chem.* **280**, 33558–33565
26. Yang, X., Chen, M. W., Terry, S., Vacherot, F., Bemis, D. L., Capodice, J., Kitajewski, J., de la Taille, A., Benson, M. C., Guo, Y., and Buttyan, R. (2006) *Oncogene* **25**, 3436–3444
27. Brown, M. S., and Goldstein, J. L. (1997) *Cell* **89**, 331–340
28. Hua, X., Wu, J., Goldstein, J. L., Brown, M. S., and Hobbs, H. H. (1995) *Genomics* **25**, 667–673
29. Shimano, H. (2001) *Prog Lipid Res.* **40**, 439–452
30. Shimano, H., Shimomura, I., Hammer, R. E., Herz, J., Goldstein, J. L., Brown, M. S., and Horton, J. D. (1997) *J. Clin. Invest.* **100**, 2115–2124
31. Menendez, J. A., Decker, J. P., and Lupu, R. (2005) *J. Cell. Biochem.* **94**, 1–4
32. Baron, A., Migita, T., Tang, D., and Loda, M. (2004) *J. Cell. Biochem.* **91**, 47–53
33. Swinnen, J. V., Heemers, H., van de Sande, T., de Schrijver, E., Brusselmans, K., Heyns, W., and Verhoeven, G. (2004) *J. Steroid Biochem. Mol. Biol.* **92**, 273–279
34. Ettinger, S. L., Sobel, R., Whitmore, T. G., Akbari, M., Bradley, D. R., Gleave, M. E., and Nelson, C. C. (2004) *Cancer Res.* **64**, 2212–2221
35. Huang, W. C., Xie, Z., Konaka, H., Sodek, J., Zhau, H. E., and Chung, L. W. (2005) *Cancer Res.* **65**, 2303–2313
36. Chung, L. W., Huang, W. C., Sung, S. Y., Wu, D., Odero-Marrah, V., Nomura, T., Shigemura, K., Miyagi, T., Seo, S., Shi, C., Moliterno, J., Elmore, J., Anderson, C., Isotani, S., Edlund, M., Hsieh, C. L., Wang, R., Shehata, B., and Zhau, H. E. (2006) *Clin. Genitourin. Cancer* **5**, 162–170
37. Yang, L., Wang, L., Lin, H. K., Kan, P. Y., Xie, S., Tsai, M. Y., Wang, P. H., Chen, Y. T., and Chang, C. (2003) *Biochem. Biophys. Res. Commun.* **305**, 462–469
38. Yang, Y. A., Han, W. F., Morin, P. J., Chrest, F. J., and Pizer, E. S. (2002) *Exp. Cell Res.* **279**, 80–90
39. Havel, P. J. (2000) *Proc. Nutr. Soc.* **59**, 359–371
40. Ngan, S., Stronach, E. A., Photiou, A., Waxman, J., Ali, S., and Buluwela, L. (2009) *Oncogene* **28**, 2051–2063
41. Heemers, H. V., Verhoeven, G., and Swinnen, J. V. (2006) *Mol. Endocrinol.* **20**, 2265–2277
42. Gross, M., Top, I., Laux, I., Katz, J., Curran, J., Tindell, C., and Agus, D. (2007) *Clin. Cancer Res.* **13**, 1979–1986
43. Heinlein, C. A., and Chang, C. (2004) *Endocr. Rev.* **25**, 276–308
44. Ruizeveld de Winter, J. A., Janssen, P. J., Sleddens, H. M., Verleun-Mooijman, M. C., Trapman, J., Brinkmann, A. O., Santerse, A. B., Schröder, F. H., and van der Kwast, T. H. (1994) *Am. J. Pathol.* **144**, 735–746
45. Di Vizio, D., Solomon, K. R., and Freeman, M. R. (2008) *Tumori* **94**, 633–639
46. Di Vizio, D., Adam, R. M., Kim, J., Kim, R., Sotgia, F., Williams, T., Demichelis, F., Solomon, K. R., Loda, M., Rubin, M. A., Lisanti, M. P., and Freeman, M. R. (2008) *Cell Cycle* **7**, 2257–2267
47. Oh, H. Y., Lee, E. J., Yoon, S., Chung, B. H., Cho, K. S., and Hong, S. J. (2007) *Prostate* **67**, 1061–1069
48. Cinar, B., Mukhopadhyay, N. K., Meng, G., and Freeman, M. R. (2007) *J. Biol. Chem.* **282**, 29584–29593
49. Freeman, M. R., and Solomon, K. R. (2004) *J. Cell. Biochem.* **91**, 54–69
50. Wu, J. D., Haugk, K., Woodke, L., Nelson, P., Coleman, I., and Plymate, S. R. (2006) *J. Cell. Biochem.* **99**, 392–401
51. Orio, F., Jr., Térouanne, B., Georget, V., Lumbroso, S., Avances, C., Siatka, C., and Sultan, C. (2002) *Mol. Cell. Endocrinol.* **198**, 105–114
52. Golias, Ch., Iliadis, I., Peschos, D., and Charalabopoulos, K. (2009) *Exp. Oncol.* **31**, 3–8
53. Devlin, H. L., and Mudryj, M. (2009) *Cancer Lett.* **274**, 177–186
54. Brooke, G. N., and Bevan, C. L. (2009) *Curr. Genomics* **10**, 18–25
55. Fuhrman, B., Nitzan, O., Karry, R., Volkova, N., Dumler, I., and Aviram, M. (2007) *Atherosclerosis* **195**, e108–e116
56. Fleischmann, M., and Iynedjian, P. B. (2000) *Biochem. J.* **349**, 13–17
57. Wu, D., Zhau, H. E., Huang, W. C., Iqbal, S., Habib, F. K., Sartor, O., Cvitanovic, L., Marshall, F. F., Xu, Z., and Chung, L. W. (2007) *Oncogene* **26**, 5070–5077