

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 04-04-2011		2. REPORT TYPE Journal Article		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE “Pairing heterocyclic cations with <i>closo</i> -dodecafluorododecaborate(2–). Synthesis of binary heterocyclium(1+) salts and a Ag ₄ (heterocycle) ₈ ⁴⁺ salt of B ₁₂ F ₁₂ ^{2–} ” (Preprint)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) John L. Belletire, Stefan Schnider, Scott Shackelford, Dmitry V. Peryshkov,. and Steven H. Strauss				5d. PROJECT NUMBER	
				5f. WORK UNIT NUMBER 23030423	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RZSP 10 E. Saturn Blvd. Edwards AFB CA 93524-7680				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-RZ-ED-JA-2011-115	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RZS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-RZ-ED-JA-2011-115	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited (PA #10989).					
13. SUPPLEMENTARY NOTES For publication in the AIAA Journal of Fluorine Chemistry.					
14. ABSTRACT Eight binary salts that pair triazolium(1+), imidazolium(1+), pyrimidinium(1+), or puri-nium(1+) cations with the icosahedral <i>closo</i> -dodecafluorododecaborate(2–) anion (B ₁₂ F ₁₂ ^{2–}) were synthesized using open-air benchtop metathesis reactions in water or acetonitrile. The scale of the reactions varied from just milligrams to nearly one gram of the starting material K ₂ B ₁₂ F ₁₂ . Other reaction conditions, the scope of the reaction, and the solubilities for the new salts are discussed. Five of the [heterocyclium]2[B ₁₂ F ₁₂] salts, which were obtained in yields ranging from 84% to 99%, displayed significantly higher densities than the corresponding previously-reported analogous [heterocyclium]2[B ₁₂ H ₁₂] and [heterocyclium][CB ₁₁ H ₁₂] salts. A ninth high-density salt consisted of B ₁₂ F ₁₂ ^{2–} paired with an Ag ₄ (triazole) ₈ ⁴⁺ cation. The structures of eight of the nine new compounds were determined by single-crystal X-ray diffraction analysis. The density of five [heterocyclium]2[B ₁₂ F ₁₂] salts was found to increase approximately linearly as the distance between the five-membered-ring heterocyclium(1+) cation centroids decreased. This work demonstrates additional flexibility for the rational design of ionic structures with predictable properties, which will ultimately permit the tailoring of ingredient-response behavior.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Mr. Wayne M. Kalliomaa
Unclassified	Unclassified	Unclassified	SAR	40	19b. TELEPHONE NUMBER (include area code) N/A

**Pairing heterocyclic cations with *closo*-dodecafluorododecaborate(2[−]).
Synthesis of binary heterocyclium(1⁺) salts and
a Ag₄(heterocycle)₈⁴⁺ salt of B₁₂F₁₂^{2−}**

John L. Belletire^a, Stefan Schneider^b, Scott A. Shackelford^{b*},
Dmitry V. Peryshkov,^c Steven H. Strauss^{c**}

^a ERC, Inc., AFRL/RZSP, 10 East Saturn Blvd., Edwards AFB, 93524-7680

^b Air Force Research Laboratory, AFRL/RZSP, 10 East Saturn Blvd., Edwards AFB, CA 92524-7680

^c Department of Chemistry, Colorado State University, Ft. Collins, CO 80523

Abstract

Eight binary salts that pair triazolium(1⁺), imidazolium(1⁺), pyrimidinium(1⁺), or purinium(1⁺) cations with the icosahedral *closo*-dodecafluorododecaborate(2[−]) anion (B₁₂F₁₂^{2−}) were synthesized using open-air benchtop metathesis reactions in water or acetonitrile. The scale of the reactions varied from just milligrams to nearly one gram of the starting material K₂B₁₂F₁₂. Other reaction conditions, the scope of the reaction, and the solubilities for the new salts are discussed. Five of the [heterocyclium]₂[B₁₂F₁₂] salts, which were obtained in yields ranging from 84% to 99%, displayed significantly higher densities than the corresponding previously-reported analogous [heterocyclium]₂[B₁₂H₁₂] and [heterocyclium][CB₁₁H₁₂] salts. A ninth high-density salt consisted of B₁₂F₁₂^{2−} paired with an Ag₄(triazole)₈⁴⁺ cation. The structures of eight of the nine new compounds were determined by single-crystal X-ray diffraction analysis. The density of five [heterocyclium]₂[B₁₂F₁₂] salts was found to increase approximately linearly as the distance between the five-membered-ring heterocyclium(1⁺) cation centroids decreased. This work demonstrates additional flexibility for the rational design of ionic structures with predictable properties, which will ultimately permit the tailoring of ingredient-response behavior.

Keywords: heterocyclium; dodecafluorododecaborate; perfluoroborane; icosahedral borane; icosahedral carborane; metathesis

*Corresponding Author. Tel. +1 661 718 2665; fax arranged by e-mail or telecom
E-mail: scott.shack@roadrunner.com (S. A. Shackelford).

**Corresponding Author. Tel +1 970 491 5104; fax +1 970 491 1801
E-mail: steven.strauss@colostate.edu (S. H. Strauss).

1. Introduction

Eight new binary salts that pair the icosahedral *closo*-dodecafluorododecaborate(2−) anion ($\text{B}_{12}\text{F}_{12}^{2-}$) with heterocyclium(1+) cations and one new salt of $\text{B}_{12}\text{F}_{12}^{2-}$ with a tetrameric $\text{Ag}_4(\text{triazole})_6^{4+}$ cation were synthesized and characterized. All nine salts were prepared using open-air metathesis reactions in which stoichiometric amounts of a heterocyclium(1+) halide and $\text{K}_2\text{B}_{12}\text{F}_{12}$ were mixed in either water or acetonitrile.

The $\text{B}_{12}\text{F}_{12}^{2-}$ dianion is analogous to the isostructural $\text{B}_{12}\text{H}_{12}^{2-}$ anion that was first reported by Pitochelli and Hawthorne in 1960 [1]. The $\text{B}_{12}\text{H}_{12}^{2-}$ anion has been called a "super-aromatic" polyhedral-shaped structure, with 26 delocalized valence electrons in its σ -bonded framework, and is conceptually similar to the delocalized π -systems of planar aromatic hydrocarbons such as benzene [2–4]. The clustering of 12 B atoms into an icosahedral cage provides a polycenter of delocalized boron orbital bonding, which, like aromatic hydrocarbon rings, is very stable to chemical attack [5]. The 12 B–H vertexes of $\text{B}_{12}\text{H}_{12}^{2-}$, like the C–H vertexes in benzene, can undergo a wide variety of aromatic substitution reactions [2–4]. This characteristic permitted the synthesis of the $\text{B}_{12}\text{F}_{12}^{2-}$ dianion by electrophilic attack of supercritical HF in 1992 (it was isolated as $\text{Cs}_2(\text{H}_2\text{O})\text{B}_{12}\text{F}_{12}$ in 38% yield) [6]. Since then, the structures of nine other salts of $\text{B}_{12}\text{F}_{12}^{2-}$ have been reported. The first of the nine was the structure of $[\text{CPh}_3]_2[\text{B}_{12}\text{F}_{12}]$ (also reported was an improved synthesis of $\text{K}_2\text{B}_{12}\text{F}_{12}$ in 72% yield) [7]. The next three were salts contained bridged heterocyclium(2+) cations, where the functional unsaturated sites, centered in the alkyl-based bridge structure, have heterocyclium(1+) moieties tethered at each alkyl bridge terminus [8]. More recent examples include a detailed X-ray crystal structure of $\text{K}_2\text{B}_{12}\text{F}_{12}$ [9], the reactant used in the metathesis reactions described herein, and in the synthesis of the heterocyclium(2+) salts [8]. Finally, the structures of $\text{K}_2(\text{H}_2\text{O})_2\text{B}_{12}\text{F}_{12}$ [10], $\text{K}_2(\text{H}_2\text{O})_2\text{B}_{12}\text{F}_{12}$ [10], $\text{K}_3(\text{AsF}_6)\text{B}_{12}\text{F}_{12}$ [11], and $\text{Cs}_3(\text{AsF}_6)\text{B}_{12}\text{F}_{12}$ [11] have just been reported (the latter two are rare examples of salts containing fluoroanions with different shapes and charges).

The binary $[\text{heterocyclium}]_2[\text{B}_{12}\text{F}_{12}]$ salts containing two heterocyclium(1+) cations reported in this work represent a new type of $\text{B}_{12}\text{F}_{12}^{2-}$ salt. These salts, and all previously

reported [heterocyclium]₂[B₁₂H₁₂] [12,13] and [heterocyclium][CB₁₁H₁₂] salts [12–16], have a unique common characteristic. For each type, a planar aromatic π -electron-delocalized heterocyclium(1+) cation is stoichiometrically paired with a polyhedral (icosahedral or pseudo-icosahedral) aromatic σ -electron-delocalized dianion in the same lattice [12,13]. Examples of each of the three types of salt we have studied are shown in Fig. 1.

The previously reported binary heterocyclium(1+) salts containing B₁₂H₁₂²⁻ and CB₁₁H₁₂⁻ can be divided into two general types based on their solubilities. Some are only sparingly soluble in H₂O and precipitated from the aqueous metathesis-reaction medium during the open-air benchtop synthesis [12]. Others were very soluble in H₂O and could be isolated in pure form only by repeated extraction of the crude solid reaction product with hot CH₃CN, followed by subsequent removal of the solvent [13].

The new [heterocyclium]₂[B₁₂F₁₂] salts, which are shown in Fig. 2, were of the same two types: some were essentially insoluble in H₂O and others were quite soluble. Therefore, they were isolated and purified by procedures parallel to those discussed above. Salts **2**, **5**, and **6** could be synthesized using either H₂O or CH₃CN as the solvent.

The new salts exhibit significantly higher densities than those of the aforementioned heterocyclium(1+) salts of B₁₂H₁₂²⁻ and CB₁₁H₁₂⁻ [12]. Reaction conditions, product yields, density values, certain salt melting/decomposition behaviors, and the general scope of the synthesis of binary heterocyclium salts of B₁₂F₁₂²⁻ are addressed in this paper.

2. Results and discussion

2.1. General comments

A long sought-after goal of synthetic chemistry has been the ability to rationally design chemical structures which will exhibit predictable chemical and physical properties for attaining desired chemical or physical behavioral responses [12,13,17]. Heterocyclic salts offer three different types of flexibility for achieving this goal. The cation ring structure or its pendant groups can be altered (e.g., 1-NH₂-3-H-1,2,3-triazolium(1+) vs. 4-NH₂-1-H-1,2,4-triazolium(1+))


[12,13] or 1-NH₂-3-H-1,2,3-triazolium(1+) vs. 1-NH₂-3-Me-1,2,3-triazolium(1+)) when paired with a common anion [12]. The chemical structure of the anion can also be changed (e.g., a polyhedral borane(2-) anion vs. a carborane(1-) anion) when paired with an identical cation [12,13]. Furthermore, a cation with a different composition (with an identical anion), or an entirely different anion (with a common cation), can be substituted in a given salt (e.g., imidazolium(1+) vs. triazolium(1+) cation [12], or nitrate(1-) vs. a polyhedral borane(2-) anion [17]).

Tailoring the thermochemical initiation response of two bridged heterocyclium salts of borane(2-) anions, and of bridged heterocyclium(2+) nitrate(1-) salts, to rapid self-sustained combustions in air by altering the cation chemical structure, or by substitution of the anion, has been demonstrated experimentally [17] and was explained by an updated high-energy-material initiation-sensitivity concept [18]. Like the previously-reported bridged heterocyclium(2+) salts of B₁₂F₁₂²⁻ [8], the binary heterocyclium(1+) salts of B₁₂F₁₂²⁻ described in this paper provide additional examples in the flexibility of the heterocyclium(*n*+) system for rational structural design, predictable property modification, and the resultant tailoring of the chemical/physical behavior of this important class of salts.

2.1. Synthesis of new salts

The nine salts shown in Fig. 2 were synthesized using metathesis reactions similar to the example shown in Scheme 1 for the synthesis of salt **2**. Most of the metatheses were carried out on a small scale initially (e.g., 44–101 mg K₂B₂F₁₂) using H₂O as the solvent. As with the pre-

Scheme 1



viously reported binary [heterocyclium]₂[B₁₂H₁₂] [12,13] or [heterocyclium][CB₁₁H₁₂] salts [12–16], the solubility of the particular B₁₂F₁₂²⁻ salt dictated whether H₂O or CH₃CN could be used as the solvent for a high-yield metathesis reaction [12,13]. It was found that salts **2**, **5**, and **6** could be prepared in good yield in either solvent. In contrast, salts **1** and **3** were best prepared using CH₃CN (even small-scale reactions in this solvent gave good yields). Finally, salts **4** and **7–9** were best prepared using H₂O as the solvent. The metathesis reaction solvent, reaction scale, yield, and melting/decomposition behavior of binary salts **1–4** and **6** are listed in Table 1.

The improved synthesis of K₂B₁₂F₁₂ reported in 2009 [19] made this reactant salt available in significantly larger quantities than if the original synthesis reported in 2003 had been used [7]. The larger-scale and shorter-reaction-time synthetic method was achieved by perfluorination of 10 g of K₂B₁₂H₁₂ to K₂B₁₂F₁₂ in 74% yield and 99.5+% purity by continuously bubbling 20/80 F₂/N₂ through a CH₃CN solution of commercially-available K₂B₁₂H₁₂ using ordinary glassware [19]. The enhanced availability of K₂B₁₂F₁₂ permitted relatively large-scale metathesis reactions to be conducted in this work using 0.453–0.902 g quantities of K₂B₁₂F₁₂. This allowed salts **2**, **4**, and **6** to be prepared in quantities exceeding 1 gram in this work, as was also the case in the work reported in ref. [8].

Accomplishing these large-scale syntheses permitted reliable product yields and spectroscopic characterization data to be obtained for salts **2**, **4**, and **6**. Additionally, a direct comparison between two identical metathesis reactions was made using the same amount of K₂B₁₂F₁₂ and either H₂O or CH₃CN as the solvent to prepare salt **2** (see Table 1). The lower yield obtained from the metathesis reaction in H₂O (i.e., 84% vs. 99% when CH₃CN was the solvent) may be due to the fact that salt **2** is slightly soluble in H₂O. After isolating the precipitated salt **2** (84% yield) by filtration (Scheme 1), evaporating the H₂O filtrate resulted in a white solid residue that was partially soluble in DMSO-d₆, leaving behind some insoluble KCl byproduct. Proton and ¹³C NMR analyses of this clear, colorless DMSO-d₆ solution revealed the presence of the 1-Me-3-H-imidazolium(1+) cation, and an FTIR spectrum taken on this same white solid filtrate residue confirmed the presence of the B₁₂F₁₂²⁻ dianion by displaying two strong peaks at

1220 and 719 cm^{-1} [19]. Both samples of salt **2**, whether synthesized in H_2O or CH_3CN , gave identical spectroscopic data consistent with the composition $[\text{1-Me-3-H-imidazolium}]_2[\text{B}_{12}\text{F}_{12}]$.

Salt **4** was prepared by reacting a stoichiometric amount of $[\text{4-NH}_2\text{-1-H-1,2,4-triazolium}][\text{I}]$ with 0.902 g $\text{K}_2\text{B}_{12}\text{F}_{12}$. The digested solid was filtered and dried and afforded 1.03 grams of a fine white powder of **4** (90.0% yield). Spectroscopic analyses of this sample of **4** were consistent with the single-crystal X-ray determined structure of **4** determined using crystals prepared with a small-scale metathesis reaction. Salt **6** was prepared by reacting a stoichiometric amount of $[\text{1-NH}_2\text{-3-H-1,2,3-triazolium}][\text{I}]$ with 0.901 g $\text{K}_2\text{B}_{12}\text{F}_{12}$ under the same conditions just described to prepare salt **4**. This afforded, after filtering and drying, 1.06 grams of **6** as an off-white solid (92.3% yield). Proton and ^{13}C NMR and FTIR spectroscopic data for this sample of **6** were consistent with the single-crystal X-ray structure of **6** determined from crystals prepared using a small-scale milligram (87.2 mg) metathesis reaction. The spectroscopic data between the large-scale and small-scale prepared samples of **6** also matched.

To minimize the presence of residual KX byproduct in these larger scale reactions ($\text{X} = \text{Cl}, \text{Br}$) [8], the aqueous metatheses were conducted by dropwise addition of an aqueous solution of $\text{K}_2\text{B}_{12}\text{F}_{12}$ to a refluxing aqueous solution of the heterocyclium(1+) halide. After refluxing for 5–8 minutes, during which precipitation of the desired $[\text{heterocyclium}]_2[\text{B}_{12}\text{F}_{12}]$ perfluoroborane salt product **2**, **4**, or **6** occurred, the reaction suspension was cooled to room temperature prior to final isolation by filtration and subsequent vacuum drying. Samples of salt **2** prepared using either H_2O or CH_3CN as the metathesis solvent were analyzed for the presence of Cl^- by ion chromatography. In both cases the mass% of Cl^- was only 0.01%, demonstrating the level of purity that is possible using the metathesis reactions shown in Scheme 1.

The attempted synthesis of $[\text{4-NH}_2\text{-1-H-1,2,4-triazolium}]_2[\text{B}_{12}\text{F}_{12}]$ instead produced $[\text{4-NH}_2\text{-1-H-1,2,4-triazolium}]_2[\text{B}_{12}\text{F}_{12}]\cdot\text{4-NH}_2\text{-1,2,4-triazole}$ (**5**). The inclusion of the neutral 4- NH_2 -1,2,4-triazole molecule in salt **5** was first thought to result from an equilibrium between the reagent $[\text{4-NH}_2\text{-1-H-1,2,4-triazolium}][\text{Cl}]$ and the free-base triazole in H_2O . However, when the same reaction was carried out in CH_3CN , the product was also salt **5**. Both of these perplexing


but reproducible results were confirmed by single crystal X-ray analysis. One might expect a higher density for salt **5** were it devoid of this neutral triazole molecule.

Small-scale aqueous metathesis reactions were used to prepare salts **7** and **8**, which crystallized with three or two H₂O molecules per formula unit in their respective lattices (see Fig. 2). The presence of H₂O molecules in the lattices of these salts precludes a detailed comparison of the packing of the B₁₂F₁₂²⁻ anions and the six-membered-ring pyrimidinium(1+) cations in **7**·3H₂O or the bicyclic purinium(1+) cations in **8**·2H₂O with the five-membered-ring heterocyclium(1+) cation salts **1–4** and **6**. In this regard the lattice H₂O molecules in **7**·3H₂O and **8**·2H₂O are acting like the "extra" 1-NH₂-1,2,3-triazole molecule in **5**, taking up what is presumably empty space in the lattice of B₁₂F₁₂²⁻ anions and protonated heterocyclium(1+) cations.

Although the expected binary salts **1–4** and **6** and the hydrated salts **7** and **8** could be readily formed using the metathesis reactions in Scheme 1, we were not able to isolate binary [heterocyclium]₂[B₁₂F₁₂] salts using halide salts of the protonated heterocyclium(1+) cations **10⁺–13⁺**, shown in Fig. 3, as metathesis reagents. For cations **10⁺–12⁺**, it appears that the putative [heterocyclium]₂[B₁₂F₁₂] salts are more soluble in H₂O than K₂B₁₂F₁₂. For cation **13⁺**, only the mixed-cation crystalline "half-salt" product [K][**13**][B₁₂F₁₂]·2H₂O, which was characterized by a preliminary X-ray structure (not shown), was obtained using H₂O as the solvent. (Note that a similar "half-salt" product with the B₁₂H₁₂²⁻ anion, [K][4-NH₂-1-H-1,2,4-triazolium][B₁₂H₁₂], was recently reported [13].)

The previously reported CH₃CN-trituration procedure [13], successfully used for water-soluble binary heterocyclium(1+) salts of B₁₂H₁₂²⁻ and CB₁₁H₁₂⁻, also was attempted, but without success. A different metathesis reaction was also tried in order to prepare the elusive binary salt [1-NH₂-3-H-1,2,3-triazolium]₂[B₁₂F₁₂] (i.e., [**13**][B₁₂F₁₂]) by driving this metathesis reaction to completion through aqueous AgCl precipitation [13]. This metathesis reaction between [**13**][Cl] and [Ag₂(CH₃CN)₄][B₁₂F₁₂] in H₂O led not to the desired product but instead to the isolation of crystalline [Ag₄(1-NH₂-1,2,3-triazole)₈][B₁₂F₁₂] (**9**) with a tetrameric tetracation,

which was characterized by single-crystal X-ray diffraction. Apparently the equilibrium shown in Rxn. 1 lies far to the left, precluding the precipitation of AgCl (tz = 1-NH₂-1,2,3-triazole).



The new salts **2**, **4**, and **6**, prepared with the large-scale metathesis reactions, were characterized by ¹H and ¹³C NMR and FTIR spectroscopy. In one case, salt **6** prepared using a small-scale reaction had spectra that matched those recorded for a sample of **6** which was synthesized using a significantly larger-scale metathesis reaction. Samples of salt **2** prepared using either solvent were analyzed for the presence of Cl[−] by ion chromatography. In both cases the mass% of Cl[−] was only 0.01%, demonstrating the level of purity that is possible using the metathesis reactions shown in Scheme 1. Most of the new salts prepared using small-scale metathesis reactions were also compositionally characterized by single-crystal X-ray diffraction analysis (see below). The single-crystal X-ray structure of salt **7** was not of sufficient quality for publication, but analysis of a low-quality data set confirmed the trihydrate composition shown in Fig. 2.

2.2. X-ray structures of the new salts

Crystallographic data and structure refinement parameters for new B₁₂F₁₂^{2−} heterocyclium(1+) salts **1–6**, **8**, and **9** are listed in Table 2. Thermal ellipsoid plots of all eight formula units are shown in the Supplementary data. Portions of the structures of salts **1–6** that are relevant to the discussion below are shown in Figs. 4 and 5. These figures depict all atoms as spheres of arbitrary size for clarity. In addition, Figs. 4 and 5 depict the centroid of each B₁₂F₁₂^{2−} anion (i.e., the B₁₂ centroid, for which the symbol ⊙ is used) as a small sphere connected by dashed lines to other B₁₂ centroids in order to show the idealized rhombs formed by eight B₁₂F₁₂^{2−} anions in salts **1–6**. Selected solid-state structure parameters for B₁₂F₁₂^{2−} salts **1–6** and

for related $\text{B}_{12}\text{H}_{12}^{2-}$ salts with the same cations, including the $\odot\cdots\odot$ distances that define the anion rhombs, are listed in Table 3.

The recently published structures of $[\text{K}_2(\text{H}_2\text{O})_2][\text{B}_{12}\text{F}_{12}]$ and $[\text{K}_2(\text{H}_2\text{O})_4][\text{B}_{12}\text{F}_{12}]$ revealed a structural motif for $\text{B}_{12}\text{F}_{12}^{2-}$ salts consisting of a rhomb of anions containing a pair of hydrated K^+ cations. The $\odot\cdots\odot$ distances in those rhombs ranged from 6.915 to 8.598 Å, which is comparable to the range of $\odot\cdots\odot$ distances for salts **1–6**, 7.122–10.675 Å. In addition, the distance between the K^+ cations within the rhombs in $[\text{K}_2(\text{H}_2\text{O})_2][\text{B}_{12}\text{F}_{12}]$ and $[\text{K}_2(\text{H}_2\text{O})_4][\text{B}_{12}\text{F}_{12}]$ are ca. 4.3 Å, which is the middle of the range of cation $\cdots$ cation distances in salts **1–6**, 3.589–5.708 Å (these distances, also listed in Table 3, are the shortest distances between the centroids of the five-membered rings of the heterocyclium(1+) cations in salts **1–6** and will hereinafter be referred to as cat $\cdots$ cat distances).

2.2.1. Imidazolium(1+) salts **1**, **2**, and **3**

A drawing of the rhombs of $\text{B}_{12}\text{F}_{12}^{2-}$ centroids ($\odot$) and their pairs of heterocyclium(1+) cations for each of these three salts is shown in Fig. 4. The rhomb for salt **1** contains parts of four 1-Et-3-Me-imidazolium(1+) cations that add up to two complete cations; the rhombs for salts **2** and **3** contain two complete 1-H-3-Me-imidazolium(1+) or H(imidazolium)(1+) cations, respectively, in the middle of their rhombs. The data in Table 3 demonstrate that as the imidazolium(1+) substituents change from Et/Me to H/Me to H/H the formula unit volumes and the $\odot\cdots\odot$ and cat $\cdots$ cat distances all decrease and the density increases. The increase in density for the hypothetical transformation **1** $\rightarrow$ **2** $\rightarrow$ **3** is from 1.576 to 1.739 to 1.829 g cm $^{-3}$, respectively.

Unlike the cation orientations in the structure of **1**, the five-membered rings of the two cations within the anion rhombs in **2** and **3** are nearly parallel: the dihedral angles of their least-squares planes with respect to one another are 3.6 and 2.5°, respectively. Furthermore, the perpendicular distances of the atoms of one five-membered cation ring to the least-squares plane of the other are 3.3–3.5 Å in both cases, producing a graphite-like separation of the two cations despite the repulsion of the positive charges on each ring.

2.2.1. Aminotriazolium(1+) salts **4**, **5**, and **6**

A drawing of the rhombs of $\text{B}_{12}\text{F}_{12}^{2-}$ centroids ($\odot$) and their heterocyclium(1+) cations for each of these three salts is shown in Fig. 5. The structures of salts **4** and **6**, like those of salts **2** and **3**, contain two complete heterocyclium(1+) cations within the rhombs, cat...cat distances of 4.647 Å in **4** and 5.287 Å in **6**. However, unlike the nearly parallel orientations of the cation five-membered rings in salts **2** and **3**, the dihedral angles between the least-squares cation rings in **4** and **6** are 56.7 and 52.7°, respectively.

The composition of salt **5**, and its solid-state structure, are different than those of salts **1–4** and **6**. In addition to the two 4-NH₂-1-H-1,2,4-triazolium(1+) cations paired with each $\text{B}_{12}\text{F}_{12}^{2-}$ anion, there is a molecule of neutral 4-NH₂-1,2,4-triazole in the formula unit. There is a five-membered ring approximately centered on each of the six faces of the anion rhomb, four 4-NH₂-1-H-1,2,4-triazolium(1+) cations and two 4-NH₂-1,2,4-triazoles (only two cations and one neutral triazole are shown in Fig. 5 for clarity). Not surprisingly, the shortest cat...cat distance in salt **5**, which is 5.445 Å, is longer than the shortest distance between a cation centroid and the five-membered ring centroid of the neutral triazole molecule, which is 4.936 Å. Also not surprisingly, the interactions between the triazolium(1+) cations and the neutral triazole molecule in salt **5** are N–H...N hydrogen bonds ((N)H...N = 1.911 Å, N–H...N = 166.5°, N...N = 2.747 Å; (N)H...N = 2.025 Å, N–H...N = 168.5°, N...N = 3.001 Å; neither of these hydrogen bonds involve NH₂ groups).

The density of each of the [heterocyclium]₂[B₁₂F₁₂] salts **1**, **2**, **4**, and **6** can be compared to the density of the corresponding [heterocyclium]₂[B₁₂H₁₂] salt (see Table 3). As with perfluorocarbon versus analogous hydrocarbon organic compounds, the heterocyclium(1+) B₁₂F₁₂^{2–} salts are more dense than the corresponding heterocyclium(1+) B₁₂H₁₂^{2–} and CB₁₁H₁₂[–] salts (for example, the densities of benzene and hexafluorobenzene are 0.877 and 1.61 g cm^{–3}, respectively, and the densities of *n*-hexane and perfluoro-*n*-hexane are 0.655 and 1.67 g cm^{–3}, respectively). A comparison of the structures of [1-H-3-Me-imidazolium]₂[B₁₂F₁₂] (**2**) and

[1-H-3-Me-imidazolium]₂[B₁₂H₁₂] [12] is especially appropriate. Interestingly, the anion rhombs are very similar in the two salts: the $\odot\cdots\odot$ distances are ca. 7.5, 8.0, and 9.4 Å in both salts, and the acute $\odot\cdots\odot\cdots\odot$ angles are 65.0, 77.8, and 84.9° in **2** and 63.9, 66.1, and 89.7° in the B₁₂H₁₂²⁻ salt, demonstrating that the relative sizes of the B₁₂F₁₂²⁻ and B₁₂H₁₂²⁻ anions are quite similar (typical B–F and B–H distances in these anions are ca. 1.38 and 1.12 Å and typical $\odot\cdots\text{F}$ and $\odot\cdots\text{H}$ distances are ca. 3.08 and 2.80 Å, respectively). In addition, the cat $\cdots$ cat distances are also similar (they only differ by 11%), 4.585 Å in **2** and 4.113 Å in the B₁₂H₁₂²⁻ salt. Nevertheless, the densities differ by a factor of 1.571 (1.739 vs. 1.107 g cm⁻³), mostly because the molar mass of the B₁₂F₁₂²⁻ and B₁₂H₁₂²⁻ anions differ by a factor of 2.52 (357.72 vs. 141.84 g mol⁻¹).

Comparison of imidazolium(1+) salts **1–3** and 1,2,4-triazolium(1+) salts **4** and **5** reveals that the salt density is higher when the cation is a protonated heterocyclium(1+) cation instead of an alkylated heterocyclium(1+) cation. In fact, there is a nearly linear relationship between the salt density and the shortest cat $\cdots$ cat distance for these five B₁₂F₁₂²⁻ salts, as shown in Fig. 6. Interestingly, there is no such correlation for the B₁₂H₁₂²⁻ salts: the data in Table 3 show that the densities, and the shortest cat $\cdots$ cat distances, are nearly the same for [1-Et-3-Me-imidazolium][B₁₂H₁₂] (1.101 g cm⁻³ and 4.119 Å) and [1-H-3-Me-imidazolium][B₁₂H₁₂] (1.107 g cm⁻³ and 4.113 Å), despite the fact that the 1.576 g cm⁻³ density of [1-Et-3-Me-imidazolium][B₁₂F₁₂] (**1**) is more than 10% lower than the 1.739 g cm⁻³ density of [1-H-3-Me-imidazolium][B₁₂F₁₂] (**2**).

2.2.3. Salt **8**

The packing of 2,6-(NH₂)₂-1-H-purinium(1+) cations and H₂O molecules inside the rhomb of eight B₁₂F₁₂²⁻ anions in the structure of [2,6-(NH₂)₂-1-H-purinium]₂[B₁₂F₁₂]·2H₂O (**8**) is shown in Fig. 7. The planes of the two purinium(1+) cations are nearly parallel (dihedral angle of only 0.2°) and the C and N atoms of the fused-ring system of one cation are ca. 3.4 Å from the least-squares plane of the C and N atoms that make up the fused-ring system of the other cation,

once again a graphite-like interaction of the two cations. The H₂O molecules form (N)H $\cdots$ O hydrogen bonds with an H $\cdots$ O distances of 1.78 Å and N–H $\cdots$ O angles of 154° and two (O)H $\cdots$ F hydrogen bonds with H $\cdots$ F distances of 2.12 and 2.21 Å. The $\odot\cdots\odot$ distances in the structure of **8** are 7.423 ($\times 2$) and 12.241 Å.

Fig. 7 also shows a drawing of the structure of salt **3** for comparison. These drawings emphasize the dovetail-like packing of the B₁₂F₁₂²⁻ anions. The $\odot\cdots\odot$ distances for the two anions shown are 7.423 Å for **8** and 7.122 Å for **3**. The angles between the two highlighted F–B–B–F planes in each drawing are 78.7° for **8** and 90° for **3**. The F $\cdots$ F distances shown are 2.779, 3.193, 3.689, and 3.796 Å in **8** and 2.908 ($\times 2$) and 3.108 Å ($\times 2$) in **3** (note that twice the van der Waals radius of an F atom is ca. 3.0 Å). This dovetail-like packing was also observed in the structures of [K(H₂O)_n]₂[B₁₂F₁₂] (*n* = 1, 2) [10] and in heterocyclium(1+) salts of the B₁₂H₁₂²⁻ and CB₁₁H₁₂⁻ anions [12,13].

2.2.4. Salt **9**

The packing of the discrete Ag₄(1-NH₂-1,2,3-triazole)₈⁴⁺ cations and B₁₂F₁₂²⁻ anions in the structure of [Ag₄(1-NH₂-1,2,3-triazole)₈][B₁₂F₁₂]₂ (**9**) is complicated and will not be discussed here. Instead, we describe the structure of the tetrameric tetracation, shown in Fig. 8, and compare it to two similar Ag₄(triazole)_n⁴⁺ tetracations [20,21]. Relevant structural parameters are listed in Table 4. The four Ag⁺ ions in each tetracation in **9** are rigorously coplanar, with Ag $\cdots$ Ag distances of 3.764(2) and 3.858(2) Å and Ag $\cdots$ Ag $\cdots$ Ag angles of 65.8(1) and 114.2(1)°. These are similar to the corresponding parameters in the structures of [Ag₄(4-NH₂-1,2,4-triazole)₇][ClO₄] [20] and [Ag₄(4-(3-C₆H₄OH)-1,2,4-triazole)₆][NO₃]₄·4H₂O [21].

An interesting feature of the structure of the tetrameric Ag₄(1-NH₂-1,2,3-triazole)₈⁴⁺ cation in **9** is that the Ag1 cations are only bonded to the more basic N3 atoms of three μ -(1-NH₂-1,2,3-triazole)- κ N2: κ N3 ligands, with Ag1–N3 distances of 2.211(3), 2.238(3), and 2.322(4) Å. The Ag2 cations, on the other hand, are bonded to the less basic N2 atoms of the three bridging triazole ligands, with Ag2–N2 distances of 2.263(3), 2.404(3), and 2.492(4) Å. Probably because

of the weaker bonding between Ag₂ and the three bridging triazole ligands, each Ag₂ cation is also coordinated to a terminal 1-NH₂-1,2,3-triazole ligand through N3, with a Ag₂–N3 distance of 2.255(3) Å. If not for this terminal triazole ligand, the sum of bond-valences [22] for Ag₂ would be significantly less than 1.0 (an Ag–N distance of 2.255 Å corresponds to a bond valence of 0.335 [22]).

3. Conclusions

Five new binary [heterocyclium]₂[B₁₂F₁₂] salts **1-4** and **6** were synthesized in 84–99% isolated yield. In addition, three salts **5**, **7** and **8** that also contained neutral molecules, either a triazole or H₂O, were also prepared. These are the first salts that contain heterocyclium(1+) cations paired with the icosahedral B₁₂F₁₂²⁻ perfluorododecaborate dianion. Open-air benchtop double-displacement (metathesis) reactions using common laboratory glassware were employed, in which a stoichiometric amount of heterocyclium(1+) halide salt was treated with K₂B₁₂F₁₂ in either H₂O or CH₃CN solvent. Depending on the particular salt, the synthesis was carried out at a temperature that ranged from room temperature to 100 °C and on a scale that ranged from 44 mg of K₂B₁₂F₁₂ to nearly 1 g of this salt.

Compared with previously reported analogous heterocyclium cation(1+) salts of the icosahedral B₁₂H₁₂²⁻ borane and CB₁₁H₁₂⁻ anions that were synthesized in a similar manner, the scope of the reaction for binary [heterocyclium]₂[B₁₂F₁₂] perfluoroborane salts was found to be more limited. Four different protonated heterocyclium(1+) cations each form the previously reported [heterocyclium]₂[B₁₂H₁₂] and the [heterocyclium][CB₁₁H₁₂] salts, but only one analogous [heterocyclium]₂[B₁₂F₁₂] salt **5** was isolated. But unlike the protonated [heterocyclium]₂[B₁₂H₁₂] and [heterocyclium]₂[CB₁₁H₁₂] analogs of **5**, [heterocyclium]₂[B₁₂F₁₂] product salt **5** also incorporated a neutral heterocycle molecule in its structure. The reduction in reaction scope resulted from the lower solubility of K₂B₁₂F₁₂ relative to several putative [heterocyclium]₂[B₁₂F₁₂] product salts that could not be isolated. In these cases, the unreacted K₂B₁₂F₁₂ reactant salt was usually the solid recovered.

Larger protonated heterocyclic cations, a six-membered pyrimidinium and a fused bicyclic purinium species, gave new sparingly-soluble [heterocyclium]₂[B₁₂F₁₂] salts **7** and **8**, respectively, that could be isolated from the aqueous metathesis reaction mixture, although both contained solvent molecules (H₂O) in the crystalline salts. In one case an isolated mixed cation [heterocyclium][K][B₁₂F₁₂] perfluoroborane "half salt" product was isolated instead of the expected [heterocyclium]₂[B₁₂F₁₂] salt product. This apparently resulted from this "half salt" product being less soluble in the H₂O reaction solvent than were either the K₂B₁₂F₁₂ reactant salt or the desired [heterocyclium]₂[B₁₂F₁₂] product. One "half salt" [heterocyclium]₂[B₁₂H₁₂] product recently has appeared in the literature. Three binary [heterocyclium]₂[B₁₂F₁₂] perfluoroborane salt products (**2**, **4**, **6**) were isolated from three different methylated heterocyclium(1+) cations using either H₂O or CH₃CN solvent, and either solvent could be used for salt **2**. One binary dialkylated [heterocyclium]₂[B₁₂F₁₂] product salt **1** was obtained. All four alkylated product salts **1**, **2**, **4**, and **6** have corresponding binary [heterocyclium][B₁₂H₁₂] and [heterocyclium][CB₁₁H₁₂] product analogs that have been reported.

There were two anomalous observations made during the syntheses described in this work. First, when attempting to synthesize [4-NH₂-1-H-1,2,4-triazolium]₂[B₁₂F₁₂] in either H₂O or CH₃CN as the solvent, the neutral triazole molecule [4-NH₂-1,2,4-triazole] was incorporated into the product salt structure (i.e., the composition was [4-NH₂-1-H-1,2,4-triazolium]₂[4-NH₂-1,2,4-triazole][B₁₂F₁₂] (**5**)). Second, an attempt was made to drive an aqueous metathesis reaction to completion with AgCl precipitation. Reacting the [4-NH₂-1-H-1,2,4-triazolium][Cl] salt with [Ag(CH₃CN)₂]₂[B₁₂F₁₂] as the source of the B₁₂F₁₂²⁻ anion in place of K₂B₁₂F₁₂, resulted in the Ag⁺ ions coordinating to the N atoms of neutral triazole molecules to give a complex tetracation that formed a unique high density salt [Ag₄(1-NH₂-1,2,3-triazole)₈][B₁₂F₁₂]₂ (**9**).

The density of individual [heterocyclium]₂[B₁₂F₁₂] perfluorododecaborane salts were significantly higher than are the recently-published analogous [heterocyclium]₂[B₁₂H₁₂] and [heterocyclium][CB₁₁H₁₂] salts. Additionally, protonated [heterocyclium]₂[B₁₂F₁₂] salts displayed a higher density than did analogous alkylated [heterocyclium]₂[B₁₂F₁₂] salts. In fact, an

interesting linear correlation between salt density and cat···cat separation was discovered for a set of five $B_{12}F_{12}^{2-}$ salts **1-4**, and **6** that does not appear to be followed for $B_{12}H_{12}^{2-}$ salts.

4. Experimental

4.1. General Comments.

Caution! While no special protective equipment or handling procedures were used with salts (**1-9**), these are high energy materials, and the analogous binary heterocyclium salts of $B_{12}H_{12}^{2-}$ and $CB_{11}H_{12}^{-}$ can be initiated to rapid energy-releasing phenomena. Thermal initiation can occur on a hot plate surface ($>275\text{ }^{\circ}\text{C}$) to rapid combustion with protonated binary heterocyclium(1+) salts [17] and bridged heterocyclium(2+) salts of $B_{12}H_{12}^{2-}$ [17]. Recent standard impact, friction, and electrostatic discharge (ESD) hazards testing conducted on the binary salt $[1\text{-NH}_2\text{-3-Me-1,2,3-triazolium}]_2[B_{12}H_{12}]$, the analogue of the new salt $[1\text{-NH}_2\text{-3-Me-1,2,3-triazolium}]_2[B_{12}F_{12}]$, gave a 100% impact non-initiation ("no-go") response at 156 kg cm, a 21.6 kg cm friction minimum non-initiation response, but initiated electrostatically at 0.040 J on an ESD test [13]. These two 100% "No-Go" impact initiation values compare with known explosives for which the following more sensitive 50% initiation kg cm values are available: CL-20 at 33; PETN at 67; HMX at 115; and RDX at 117 kg cm. One bridged heterocyclium(2+) salt of $B_{12}F_{12}^{2-}$, the only salt of $B_{12}F_{12}^{2-}$ tested to date, gave a minimum non-initiation impact at 147 kg cm (100% "no-go" non-initiation value), a minimum non-initiation friction response at 21.6 kg cm, but initiated at 0.090 J in the ESD test [13].

All water used was deionized water obtained from a Millipore MILL-Q Reagent Grade Water System at the 18 M Ω cm level of purity. All organic solvents were commercially available as either Reagent Grade or HPLC purity and were used as received.

With one exception, the neutral heterocyclic compounds used to synthesize $B_{12}F_{12}^{2-}$ salts **1-9** were commercially available and were used as received. The exception is the compound 1-NH₂-1,2,3-triazole, which was prepared using a published procedure [23]. Conversion of each neutral heterocycle to the corresponding hydrochloride or *N*-methylated iodide salt was carried

out as previously described (see Supplemental Information) [12] and detailed below. The heterocyclium(1+) hydrochloride salts were obtained from the commercial free bases via treatment with concentrated aqueous HCl in ethanol at 25 °C, followed by removal of all volatiles by rotary evaporation and drying under vacuum (≤ 0.1 mTorr). The remaining solids were recrystallized from a minimum amount of hot 2-propanol. In some cases the addition of diethyl ether after the onset of crystallization resulted in a slightly higher yield. The microcrystalline products were filtered, washed with diethyl ether, and dried under vacuum.

An alternative method is possible for the synthesis of the hydrochloride of 4-NH₂-1,2,4-triazole. This salt can also be synthesized in a 250 mL round-bottom flask by adding 2.50 g (29.7 mmol) of 4-NH₂-1,2,4-triazole and 2.50 mL (29.9 mmol) of 37% aqueous HCl to 50 mL MeOH and letting the mixture stand at room temperature for two days. High vacuum Removal of the solvent under vacuum afforded 3.53 g of a thick syrup that was recrystallized from hot MeOH and Et₂O. Filtration, rinsing with a small portion of Et₂O, and vacuum drying afforded 2.24 g of microcrystalline [4-NH₂-1-H-1,2,4-triazolium][Cl] (62.6% yield). The methylated heterocyclium(1+) iodide salts were obtained by reaction of the neutral free-base heterocycle with excess CH₃I in EtOH at room temperature for at least 12 h followed by removal of all volatiles by rotary evaporation (the reaction vessel was wrapped with foil to protect the reaction mixture from direct light). If necessary, some of these salts were recrystallized from MeOH. The preparation of [H(imidazolium)][Cl] for the synthesis of salt **3** was achieved by bubbling dry HCl gas (10–15 mL/min flow rate) through an anhydrous CHCl₃ solution of imidazole for 30 min. The white needle-shaped crystals that formed were filtered, washed with anhydrous CHCl₃, and dried under vacuum. The preparation of [1-Et-3-Me-imidazolium][Br] for the synthesis of salt **1** was achieved by reacting EtBr (23.1 mmol) and 1-Me-imidazole (11.2 mmol) in refluxing anhydrous benzene (20 mL) for 2 h. Filtering the cooled reaction mixture, washing the white crystalline product, and drying under vacuum afforded [1-Et-3-Me-imidazolium][Br] in 64% yield. The salt [Ag(CH₃CN)₂]₂[B₁₂F₁₂] was prepared as previously described [24].

4.2. Spectroscopy, X-ray crystallography, ion chromatography, and melting points

A Bruker Avance 400 Digital NMR instrument was used to obtain both ^1H and ^{13}C spectra. FTIR spectra were taken as powder samples in air using a Nicolet 6700 Spectrometer equipped with an HATR optical system. Only the significant peaks that were observed are listed below. The FTIR spectrum of $\text{K}_2\text{B}_{12}\text{F}_{12}$ salt exhibits only two strong peaks at 1221 cm^{-1} ($\nu(\text{BF})$) and 719 cm^{-1} ($\nu(\text{BB})$) [19]. The presence of $\text{B}_{12}\text{F}_{12}^{2-}$ was confirmed in [heterocyclium] $_2[\text{B}_{12}\text{F}_{12}]$ salts **2**, **4**, and **6** by the presence of two strong peaks at ca. 1220 and ca. 720 cm^{-1} in their respective FTIR spectra.

Single-crystal X-ray diffraction data for salts **2**, **4–6**, **8**, and **9** were collected at Edwards AFB using a Bruker 3-circle-platform diffractometer equipped with a SMART APEX 2 detector with the χ -axis fixed at 54.74° using $\text{Mo K}\alpha$ or $\text{Cu K}\alpha$ radiation from a fine-focus tube. The goniometer head, equipped with a Nylon Cryoloop and magnetic base, was used to mount the crystals using perfluoropolyether oil. Diffraction data for salts **1** and **3** were collected at Colorado State University using a Bruker Kappa APEX II CCD diffractometer at $110(2)\text{ K}$ employing $\text{Mo K}\alpha$ radiation (graphite monochromator). Unit cell parameters were obtained from least-squares fits to the angular coordinates of all reflections, and intensities were integrated from a series of frames (ω and φ rotation) covering more than a hemisphere of reciprocal space. Absorption and other corrections were applied using SADABS [25]. The structures were solved using direct methods and refined (on F^2 using all data) by a full-matrix, weighted least-squares process. Standard Bruker control and integration software (APEX II) was employed [26], and Bruker SHELXTL software was used for structure solution, refinement, and molecular graphics [27]. Crystal data and refinement details for salts **1–6**, **8**, and **9** are listed in Table 2.

For ion chromatography Cl^- analyses, each sample was weighed to 0.01 mg and diluted to 25 mL in a class B centrifuge tube with $18\text{ M}\Omega\text{ cm}$ deionized water. The samples that were not readily soluble were heated to 80°C with a reflux cap. All solution transfers and injections were accomplished with sterile syringes and syringe filters (IC Millex LG filter unit $0.2\text{ }\mu\text{m}$). A four-point linear calibration curve was generated using a blank deionized water solution, followed by

a 0.1, 1, and 5 ppm ($\mu\text{g g}^{-1}$) NIST traceable Cl^- standard where the peak area was determined for each chloride standard solution. Lower detection limits (LDL) and lower quantitation limits (LQL) were determined using seven individually prepared blanks consisting only of deionized H_2O (LDL = 3 times and LQL = 5 times the standard deviation of the seven blank runs). Concentrations were calculated by comparing peak area response versus the linear calibration graph and are corrected for dilution.

Melting points were determined using a Stanford Research Systems OptiMelt MPA100-Automated Melting Point Apparatus equipped with digital image video playback software.

4.3. Small-scale CH_3CN solvent synthesis procedure

The heterocyclium halide salt was dissolved in CH_3CN and mixed with a CH_3CN solution of $\text{K}_2\text{B}_{12}\text{F}_{12}$ at room temperature to form a white KX precipitate ($\text{X} = \text{Cl}, \text{Br}$). The solution was filtered and dried to give the desired binary $[\text{heterocyclium}]_2[\text{B}_{12}\text{F}_{12}]$ salt, which was subsequently recrystallized from CH_3CN .

4.3.1. $[1\text{-Et-3-Me-imidazolium}]_2[\text{B}_{12}\text{F}_{12}]$ (**1**)

Solutions of $[1\text{-Et-3-Me-imidazolium}][\text{Br}]$ (0.088 g, 0.45 mmol) in 2 mL CH_3CN and $\text{K}_2\text{B}_{12}\text{F}_{12}$ (0.10 mg, 0.22 mmol) in 2 mL CH_3CN were mixed at room temperature, affording 0.13 g of salt **1** (90% yield). Single-crystal X-ray diffraction analysis confirmed the composition of this salt.

4.3.2. $[\text{H}(\text{imidazolium})]_2[\text{B}_{12}\text{F}_{12}]$ (**3**)

Solutions of $[\text{H}(\text{imidazolium})][\text{Cl}]$ (0.048 g, 45 mmol) in 2 mL CH_3CN and $\text{K}_2\text{B}_{12}\text{F}_{12}$ (0.10 mg, 0.22 mmol) in 2 mL CH_3CN were mixed at room temperature, affording 0.10 g of salt **3** (93% yield). Single-crystal X-ray diffraction analysis confirmed the composition of this salt.

4.3.3. $[4\text{-NH}_2\text{-1-H-1,2,4-triazolium}]_2[\text{triazole}][\text{B}_{12}\text{F}_{12}]$ (**5**)

The compounds [4-NH₂-1-H-1,2,4-triazolium][Cl] (20.1 mg, 0.167 mmol) and K₂B₁₂F₁₂ (36.4 mg, 0.0835 mmol) together which were triturated with 5 × 8 mL of boiling CH₃CN. The solution was cooled and the solvent removed under vacuum to yield 48.9 mg (95.7% yield) of off-white solid (95.7%). The solid was dissolved in a mixture of 1 mL CH₃CN and 40 mL Et₂O, and after standing overnight in a dessicator this solution deposited crystals of **5** suitable for single crystal X-ray diffraction analysis which revealed that one neutral 4-NH₂-1-H-1,2,4-triazole molecule occupied the formula unit.

4.3.4. [1-NH₂-3-Me-1,2,3-triazolium]₂[B₁₂F₁₂] (**6**)

The compounds [1-NH₂-3-Me-1,2,3-triazolium][I] (0.0904 g, 0.400 mmol) and K₂B₁₂F₁₂ (87.2 mg, 0.200 mmol) were each dissolved in a minimum amount of CH₃CN at room temperature. The two solutions were mixed and the solvent was removed under vacuum to yield 0.179 g of a solid containing some KI byproduct. The crude solid dissolved in 1.24 mL H₂O, refluxed, and cooled, producing crystals of **6** used for single crystal X-ray diffraction analysis. ¹H NMR (400 MHz, DMSO-d₆ (δ 2.50)): δ 8.74 (d_{ovlap.}, 2H); 8.06 (d_{ovlap.}, 2H); 8.82 (s, 4H); 4.21 (s, 6H). ¹³C NMR (100 MHz, DMSO-d₆ (δ 39.51)): δ 132.54, 126.84, 39.63 (shoulder on DMSO-d₆ peak δ 39.72); CD₃CN (δ 1.39), δ 132.22, 129.58, 41.12. HATR-FTIR: 3375, 3309, 3185, 3166, 1624, 1535, 1434, 1396, 1330, 1222 (B₁₂F₁₂²⁻), 1091, 1029, 948, 797, 725 (B₁₂F₁₂²⁻), 667, 630 cm⁻¹. These spectroscopic data are identical to those listed for salt **6** in Section 4.6.3, below.

4.4. Small-scale H₂O solvent synthesis procedure

The [heterocyclium][halide] salt and K₂[B₁₂F₁₂] were mixed as solids into one flask. The mixture was dissolved in a minimum amount of H₂O and warmed if necessary, usually below the boiling point. The solution was cooled to room temperature and placed into a refrigerator (3.5 °C) overnight or until crystals formed. This generally produced crystals suitable for single-

crystal X-ray diffraction analysis. Otherwise suitable crystals were obtained with solvent removal followed by recrystallization.

4.4.1. *[4-NH₂-1-Me-1,2,4-triazolium]₂[B₁₂F₁₂] (4)*

The compounds *[4-NH₂-1-Me-1,2,4-triazolium][I]* (42.5 mg, 0.329 mmol) and K₂B₁₂F₁₂ (43.6 mg, 0.100 mmol) were dissolved in 0.54 mL H₂O and heated. Cooling gave crystals of **4** suitable for single-crystal X-ray diffraction analysis.

4.4.2. *[4-NH₂-1H-1,2,4-triazolium]₂[4-NH₂-1,2,4-triazole][B₁₂F₁₂] (5)*

The compounds *[4-NH₂-1-H-1,2,4-triazolium][Cl]* (17.2 mg, 0.143 mmol) and K₂B₁₂F₁₂ (43.6 mg, 0.100 mmol) were dissolved in 5 mL H₂O and heated to 95 °C for 15 min. The solvent was removed under vacuum (ca. 18 mTorr) for 12 h and the resultant solid was redissolved in 1 mL H₂O and placed in a refrigerator (3.5 °C) overnight. This afforded crystals of **5** suitable for single-crystal X-ray diffraction analysis which revealed that one neutral 4-NH₂-1-H-1,2,4-triazole molecule occupied the formula unit.

4.4.3. *[2,4,6-(NH₂)₃-1-H-pyrimidinium]₂[B₁₂F₁₂] (7)*

The compounds *[2,4,6-(NH₂)₃-1-H-pyrimidinium][Cl]* (64.6 mg, 0.400 mmol) and K₂B₁₂F₁₂ (87.2 mg, 0.200 mmol) were dissolved in 3 mL H₂O, brought to a boil, and cooled. The cooled solution deposited crystals of **7**·3H₂O that were used for a preliminary X-ray diffraction study.

4.4.4. *[2,6-diamino-1-H-purinium]₂[B₁₂F₁₂] (8)*

The compounds *[2,6-(NH₂)₂-1-H-purinium][Cl]* (74.6 mg, 0.400 mmol) and K₂B₁₂F₁₂ (87.2 mg, 0.200 mmol) were dissolved in 3 mL H₂O and brought to a boil. When cooled, this solution deposited crystals of **8**·2H₂O suitable for single crystal X-ray diffraction analysis.

4.4.5. *[Ag₄(1-NH₂-1,2,3-triazole)₈][B₁₂F₁₂]₂ (9)*

The compounds [1-NH₂-3-H-1,2,3-triazolium][Cl] (24.1 mg, 0.280 mmol) and [Ag(CH₃CN)₂]₂[B₁₂F₁₂] (73.7 mg, 0.0999 mmol) were dissolved in 7 mL water and filtered through Celite. Solvent removal under high vacuum gave 42.4 mg (38.8% yield) of a solid which was dissolved in EtOH, filtered, and recrystallized from EtOH to give crystals of **9** suitable for single-crystal X-ray diffraction analysis.

4.5. Large-scale CH₃CN solvent synthesis of [1-Me-3-H-imidazolium]₂[B₁₂F₁₂] (**2**)

In a reaction flask containing a Teflon-coated magnetic stirring bar, [H(MeIm)][Cl] (0.254 g, 2.15 mmol) was dissolved in 5 mL CH₃CN at room temperature. K₂B₁₂F₁₂ (0.454 g, 1.04 mmol) was dissolved in 4 mL CH₃CN at room temperature in a second glass vessel. The K₂B₁₂F₁₂ solution was added during 4 min to the stirred reaction flask solution using a disposable pipette, forming a precipitate. The second glass vessel was rinsed with 2 × 0.5 mL portions CH₃CN, each of which was added to the stirred reaction flask during 1 min for a total addition time of 6 min. The suspension was stirred at room temperature for 17 h, then filtered using Celite to remove the white solid KCl byproduct. The CH₃CN solvent filtrate was removed by rotary evaporation and gave an off-white solid. Removal of the volatiles under high vacuum at 65 °C (24 h) in an Electrothermal ChemDry[®] apparatus gave 0.540 g (98.9% yield) of a light tan solid. ¹H NMR (400 MHz; DMSO-d₆ (δ 2.50)): δ 14.17 (bd s, 2H); 9.03 (s, 2H); 7.69–7.68 (m, 2H); 7.66 (m, 2H); 3.86 (s, 6H). ¹³C NMR (100 MHz; DMSO-d₆ (δ 39.51)): δ 135.82, 123.16, 119.78, 35.42. HATR-FTIR: 3397, 3355, 3178, 3122, 3078, 1640, 1585, 1443, 1329, 1306, 1218 (B₁₂F₁₂²⁻), 1142, 1104, 1082, 1008, 874, 844, 763, 723 (B₁₂F₁₂²⁻), 697, 616 cm⁻¹. An aqueous synthesis of this salt **2** conducted on the same scale is described in paragraph 4.6.1.

4.6. Large-scale H₂O solvent synthesis procedure

The [heterocyclium][halide] salt was dissolved in a minimum volume of H₂O in a 25 mL 14/20 single-necked recovery flask reaction vessel containing a Teflon-coated stirring bar. The flask was connected to a water-cooled reflux condenser and the solution was stirred and heated

to reflux in a 107–108 °C oil bath for 5 min. At this time, an aqueous solution of $\text{K}_2\text{B}_{12}\text{F}_{12}$ that had been prepared in a minimum amount of warm H_2O in a 15 mL single-necked pear-shaped flask was added dropwise to the reaction flask, during 5–7 min, through the center of the reflux condenser using a disposable pipette, at which point a precipitate was observed. The pear-shaped flask that previously contained the $\text{K}_2\text{B}_{12}\text{F}_{12}$ solution was rinsed with 2×1 mL portions of H_2O , each of which were added dropwise to the refluxing reaction mixture. After an additional 5 min, the cloudy reaction mixture was cooled to room temperature and placed in a 3.5 °C refrigerator overnight. Vacuum filtration, rinsing the solid cake with 2×1 mL of 3.5 °C H_2O and air-drying in the Büchner funnel at room temperature for ca. 1 h gave a semi-dry solid. The solid was placed into a 4 dram bottle and dried at 65 °C under vacuum (ca. 10^{-4} Torr) in a Electrothermal ChemDry[®] apparatus for 18–55 h to yield the desired anhydrous salt.

4.6.1. [1-Me-3-H-imidazolium]₂[B₁₂F₁₂] (2)

[H(MeIm)][Cl] (0.254 g, 2.14 mmol) in 1 mL H_2O and the $\text{K}_2\text{B}_{12}\text{F}_{12}$ (0.453 g, 1.04 mmol) in 4 mL H_2O was added dropwise over 7 min., then cooled 24 h in the refrigerator, filtered, and vacuum dried 24 h. Yield: 0.460 g (84.4%) of a slightly off-white solid. ¹H NMR (400 MHz, DMSO-d₆ (δ 2.50)): δ 14.15 (bd. s, 2H); 9.03 (s, 2H), 7.68 (m, 2H); 7.66 (m, 2H); 3.86 (s, 6H). ¹³C NMR (100 MHz, DMSO-d₆ (δ 39.51)): δ 135.81, 123.15, 119.77, 35.42. HATR-FTIR: 3398, 3355, 3179, 3124, 3087, 1640, 1585, 1552, 1442, 1328, 1307, 1216 (B₁₂F₁₂²⁻), 1141, 1104, 1080, 1008, 877, 843, 763, 719 (B₁₂F₁₂²⁻), 695, 615 cm⁻¹.

4.6.2. [4-NH₂-1-Me-1,2,4-triazolium]₂[B₁₂F₁₂] (4)

[4-NH₂-1-Me-1,2,4-triazolium][I] (0.964 g, 4.27 mmol) in 2 mL H_2O and $\text{K}_2\text{B}_{12}\text{F}_{12}$ (0.902 g, 2.07 mmol) in 8 mL H_2O was added dropwise over 5 min., then cooled 21.5 h in the refrigerator, filtered, and vacuum dried 18.5 h. Yield: 1.03 g (90.0%) of a slightly off-white solid. ¹H NMR (400 MHz, DMSO-d₆ (δ 2.50)): δ 10.06 (s, 2H); 9.16 (s, 2H); 6.95 (s, 4H); 4.02 (s, 6H). ¹³C NMR (100 MHz, DMSO-d₆ (δ 39.51)): δ 145.04, 142.95, unknown (3rd peak masked

by DMSO and CD₃CN internal standard), 146.13, 143.95, 118.41 (CD₃CN), 40.24. HATR-FTIR: 3386, 3317, 3159, 3120, 1620, 1582, 1567, 1434, 1407, 1220 (B₁₂F₁₂²⁻), 1170, 1072, 993, 980, 942, 891, 722 (B₁₂F₁₂²⁻), 659, 621 cm⁻¹.

4.6.3. [1-NH₂-3-Me-1,2,3-triazolium]₂[B₁₂F₁₂] (**6**)

[1-NH₂-3-Me-1,2,3-triazolium][I] (0.964 g, 4.26 mmol) in 2 mL H₂O and K₂B₁₂F₁₂ (0.901 g, 2.07 mmol) in 7 mL H₂O was added dropwise over 5 min., then cooled 18.5 h in the refrigerator, filtered, and vacuum dried for 54.5 h. Yield: 1.06 g (92.3%) of a light tan solid. ¹H NMR (400 MHz, DMSO-d₆ (δ 2.50): δ 8.74 (d_{ovlap.}, 2H); 8.60 (d_{ovlap.}, 2H); 8.28 (s, 4H); 4.21 (s, 6H). ¹³C NMR (100 MHz, DMSO-d₆ (δ 39.51)): δ 131.52, 126.83, unknown (3rd peak masked by DMSO and CD₃CN internal standard), 132.23, 129.59, 41.13. HATR-FTIR: 3375, 3309, 3185, 3166, 1622, 1533, 1434, 1400, 1329, 1218 (B₁₂F₁₂²⁻), 1090, 1028, 947, 798, 722 (B₁₂F₁₂²⁻), 663, 629 cm⁻¹. These spectroscopic data are identical to those listed for salt **6** in Section 4.3.4.

Supplementary data

Figures showing the formula units of salts **1-6**, **8**, and **9** as thermal ellipsoid plots. These can be found in a document located at doi:10.1016/J.fluchem.2011.XY.XYZ. Crystallographic data for salts **1-6**, **8**, and **9** have been deposited with the Cambridge Crystallographic Data Centre (deposition numbers CCDC **XYZXYZ-XYZXYZ**). Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK, fax: +44 1223 336033; email: deposit@ccdc.cam.ac.uk.

Acknowledgment

The authors thank the AFRL Space and Missiles Division (AFRL/RZS), Edwards AFB, CA, and ERC, Inc., Huntsville, AL, for funding support, Dr. Jerry A. Boatz (AFRL/RZSP) and Prof. Herman L. Ammon (Department of Chemistry and Biochemistry, University of Maryland) for

helpful technical discussions, and Mr. Brett A. Wight (AFRL/RZSP) for the Cl^- ion chromatography analyses of samples of salt **2**.

Table 1

Distribution A. Approved for public release; distribution unlimited.

Metathesis reaction solvent, scale, yield, and melting/decomposition behavior for **1–4** and **6**.^a

Salt	Solvent	Rxn. scale (g K ₂ B ₁₂ F ₁₂)	Yield	Melting/decomposition behavior
1	CH ₃ CN	0.100	90%	Not determined
2	CH ₃ CN	0.454	99%	Slowly converted to a dark-brown solid with some liquification and gas evolution up to 398 °C.
2	H ₂ O	0.453	84%	Slowly converted to a dark-brown solid with some liquification and gas evolution up to 398 °C.
3	CH ₃ CN	0.101	93%	Not determined
4	H ₂ O	0.436	90%	Sintered at 190.1–191.2 °C; liquified at 191.3–193.2 °C.
6	H ₂ O	0.436	92%	Sintered at 201.3–202.2 °C; liquified at 202.3–204.2 °C.

^a The five salts listed in this table are as follows: **1**, [1-Et-1-Me-imidazolium]₂[B₁₂F₁₂]; **2**, [1-H-3-Me-imidazolium]₂[B₁₂F₁₂]; **3**, [H(imidazolium)]₂[B₁₂F₁₂]; **4**, [4-NH₂-1-Me-1,2,4-triazolium]₂[B₁₂F₁₂]; **6**, [1-NH₂-3-Me-1,2,3-triazolium]₂[B₁₂F₁₂]. Drawings of the cations in these salts are shown in Fig. 2.

Table 2Crystallographic data and structure refinement parameters for B₁₂F₁₂²⁻ salts **1–6**, **8**, and **9**.^a

	1	2	3	4	5	6	8	9
Molecular formula	C ₁₂ H ₂₂ B ₁₂ F ₁₂ N ₄	C ₈ H ₁₄ B ₁₂ F ₁₂ N ₄	C ₆ H ₁₀ B ₁₂ F ₁₂ N ₄	C ₆ H ₁₄ B ₁₂ F ₁₂ N ₈	C ₆ H ₁₄ B ₁₂ F ₁₂ N ₁₂	C ₆ H ₁₄ B ₁₂ F ₁₂ N ₈	C ₁₀ H ₁₈ B ₁₂ F ₁₂ N ₁₂ O ₂	C ₈ H ₁₆ Ag ₂ B ₁₂ F ₁₂ N ₁₆
Formula weight	580.06	523.95	495.90	555.97	612.01	555.97	606.08	909.83
Crystal system	Monoclinic	Monoclinic	Monoclinic	Monoclinic	Orthorhombic	Monoclinic	Monoclinic	Monoclinic
Space group	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ / <i>c</i>	<i>C</i> 2/ <i>m</i>	<i>P</i> 2 ₁ / <i>c</i>	<i>P</i> 2 ₁ 2 ₁	<i>C</i> 2/ <i>c</i>	<i>C</i> 2/ <i>c</i>	<i>P</i> 2 ₁ / <i>n</i>
<i>Z</i>	4	2	4	2	4	4	4	4
Color of crystal	White	White	White	White	White	White	White	White
Unit cell dims.								
<i>a</i> , Å	15.6537(6)	7.5255(2)	14.245(2)	7.6121(9)	10.882(1)	19.065(2)	25.148(1)	8.9933(5)
<i>b</i> , Å	9.0441(3)	17.0776(4)	14.556(2)	17.941(2)	13.047(1)	9.2222(8)	11.7631(5)	11.7111(6)
<i>c</i> , Å	17.2734(6)	7.9652(2)	10.025(1)	8.494(1)	16.194(2)	14.806(2)	9.0568(4)	28.124(1)
<i>α</i> , deg	90	90	90	90	90	90	90	90
<i>β</i> , deg	90.980(2)	102.234(1)	119.971(9)	113.247(2)	90	120.639(4)	104.634(2)	97.807(2)
<i>γ</i> , deg	90	90	90	90	90	90	90	90
Unit cell vol., Å ³	2445.1(2)	1000.42(4)	1800.7(5)	1065.9(2)	2299.1(4)	2239.7(4)	2592.2(2)	2934.6(3)
Temperature, K	100(2)	173(2)	100(2)	173(2)	173(2)	173(2)	173(2)	296(2)
Final <i>R</i> indices	<i>R</i> ₁ = 0.046	<i>R</i> ₁ = 0.039	<i>R</i> ₁ = 0.086	<i>R</i> ₁ = 0.034	<i>R</i> ₁ = 0.032	<i>R</i> ₁ = 0.069	<i>R</i> ₁ = 0.037	<i>R</i> ₁ = 0.038
[<i>I</i> > 2σ(<i>I</i>)]	<i>wR</i> ₂ = 0.144	<i>wR</i> ₂ = 0.100	<i>wR</i> ₂ = 0.217	<i>wR</i> ₂ = 0.097	<i>wR</i> ₂ = 0.085	<i>wR</i> ₂ = 0.196	<i>wR</i> ₂ = 0.103	<i>wR</i> ₂ = 0.105
GOF on <i>F</i> ²	1.048	1.124	1.101	1.089	1.069	1.086	1.064	1.030

^a The eight salts listed in this table are as follows: **1**, [1-Et-1-Me-imidazolium]₂[B₁₂F₁₂]; **2**, [1-H-3-Me-imidazolium]₂[B₁₂F₁₂]; **3**, [H(imidazolium)]₂[B₁₂F₁₂]; **4**, [4-NH₂-1-Me-1,2,4-triazolium]₂[B₁₂F₁₂]; **5**, [4-NH₂-1-H-1,2,4-triazolium]₂[B₁₂F₁₂]-4-NH₂-1,2,4-triazole; **6**, [1-NH₂-3-Me-1,2,3-triazolium]₂[B₁₂F₁₂]; **8**, [2,6-(NH₂)₂-1-H-purinium]₂[B₁₂F₁₂]-2H₂O; **9**, [Ag₄(1-NH₂-1,2,3-triazole)₈][B₁₂F₁₂]. Drawings of the cations in these salts are shown in Fig. 2.

Table 3**Distribution A. Approved for public release; distribution unlimited.**

Solid-state parameters for B₁₂F₁₂²⁻ and B₁₂H₁₂²⁻ salts of heterocyclium(1+) cations.^a

Compound	ref ^b	formula unit vol., Å ³	density, ^c g cm ⁻³	⊖⋯⊖, Å	cation⋯cation, Å ^d
[1-Et-3-Me-imidazolium] ₂ [B ₁₂ F ₁₂] (1)	tw	611.3(1)	1.576	9.003, 9.044, 10.675	5.708
[1-Et-3-Me-imidazolium] ₂ [B ₁₂ H ₁₂]	[12]	549.4(2)	1.101 ^e	8.268, 8.673, 8.846	4.119
[1-H-3-Me-imidazolium] ₂ [B ₁₂ F ₁₂] (2)	tw	500.21(2)	1.739	7.525, 7.965, 9.422	4.585
[1-H-3-Me-imidazolium] ₂ [B ₁₂ H ₁₂]	[12]	461.9(4)	1.107 ^f	7.535, 8.167, 9.348	4.113
[H(imidazolium)] ₂ [B ₁₂ F ₁₂] (3)	tw	450.2(1)	1.829	7.122, 7.278, 8.938	3.589
[4-NH ₂ -1-Me-1,2,4-triazolium] ₂ [B ₁₂ F ₁₂] (4)	tw	533.0(1)	1.732	7.612, 8.891, 9.925	4.647
[4-NH ₂ -1-Me-1,2,4-triazolium] ₂ [B ₁₂ H ₁₂]	[12]	454.3(1)	1.243 ^g	7.036, 8.140, 8.140	4.124
[4-NH ₂ -1-H-1,2,4-triazolium] ₂ [B ₁₂ F ₁₂]-4-NH ₂ -1,2,4-triazole (5)	tw	574.8(1)	1.768	8.100, 8.480, 8.712	5.445 ^h
[1-NH ₂ -3-Me-1,2,3-triazolium] ₂ [B ₁₂ F ₁₂] (6)	tw	559.9(1)	1.649	7.609, 9.222, 10.692	5.287
[1-NH ₂ -3-Me-1,2,3-triazolium] ₂ [B ₁₂ H ₁₂]	[12]	620.1(1)	1.19 ^{i,j}	8.689, 8.959, 10.324	3.785

^a See also Table 2. ^b tw = this work. ^c X-ray diffraction-derived densities at 173(2) or 100(2) K. ^d The shortest distance between the centroids of the five-membered rings of the heterocyclium(1+) cations. ^e The density of [1-Et-1-Me-imidazolium][CB₁₁H₁₂] at 173 K is 1.03 g cm⁻³. ^f The density of [1-H-3-Me-imidazolium][CB₁₁H₁₂] at 173 K is 1.11 g cm⁻³ [12]. ^g The density of [4-NH₂-1-Me-1,2,4-triazolium][CB₁₁H₁₂] at 173 K is 1.12 g cm⁻³ [12]. ^h This is the shortest distance between cation centroids. The shortest distance between a cation centroid and the five-membered ring centroid of the neutral molecule of 4-NH₂-1,2,4-triazole is 4.936 Å. ⁱ The density of [1-NH₂-3-Me-1,2,3-triazolium][CB₁₁H₁₂] at 173 K is 1.10 g cm⁻³ [12]. ^j The density reported here for [1-NH₂-3-Me-1,2,3-triazolium]₂[B₁₂H₁₂] was determined at 25 °C by pycnometry on a vacuum-dried sample because crystals of this compound contained two molecules of CH₃CN per formula unit [12]. The X-ray density of [1-NH₂-3-Me-1,2,3-triazolium]₂[B₁₂H₁₂]-2CH₃CN is 1.132 g cm⁻³ [12].

Table 4Structural results for tetrameric silver(I) triazole compounds.^a

parameter	[Ag ₄ (tz-1) ₈][B ₁₂ F ₁₂] ₂	[Ag ₄ (tz-2) ₇][ClO ₄] ₄	[Ag ₄ (tz-3) ₆][NO ₃] ₄ ·4H ₂ O
reference	this work	[20]	[21]
Ag1–N	2.211(3), 2.238(3), 2.322(4) Å	2.179(4), 2.200(4), 2.314(5) Å	2.230(3), 2.247(3), 2.317(3)
Ag1–F	2.960(3) Å		
Ag2–N	2.255(3), 2.263(3), 2.404(3), 2.492(4) Å	2.283(4), 2.283(4), 2.326(5), 2.369(4)	2.253(3), 2.255(3), 2.311(3) Å
Ag2–F	2.806(3) Å		
Ag1 bond- valence sum	1.05	1.09	0.98
Ag2 bond- valence sum	1.13	1.14	0.96
Ag1⋯Ag2	3.764(2) Å	3.156(1) Å	3.714(2) Å
Ag1⋯Ag2'	3.858(2) Å	3.632(1) Å	3.905(2) Å
Ag2⋯Ag1⋯Ag2	114.2(1)°	156.41°(3)	92.8(1)°
Ag1⋯Ag2⋯Ag1	65.8(1)°	132.68°(3)	87.2(1)°

^a tz-1 = 1-NH₂-1,2,3-triazole; tz-2 = 4-NH₂-1,2,4-triazole; tz-3 = 4-(3-C₆H₄OH)-1,2,4-triaz




Fig. 1. Examples of binary heterocyclium(1+) salts of $B_{12}F_{12}^{2-}$ (top), $B_{12}H_{12}^{2-}$ (middle), and $CB_{11}H_{12}^{-}$ (bottom). Each vertex of the icosahedral anions bears an F atom (top) or a H atom (middle and bottom). The black sphere in the drawing of the $CB_{11}H_{12}^{-}$ anion is the C-H vertex. The anion drawings also show the numbering of the vertexes.




Fig. 2. The compositions of salts **1–9** studied in this work. The positive charge in salt **8** is delocalized over the entire fused-ring system. The positive charges in salt **9** are delocalized to some extent over the eight triazole molecules, which are coordinated to the Ag^+ cations.




Fig. 3. Heterocyclium(1⁺) cations that did not readily form binary [heterocyclium]₂[B₁₂F₁₂] salts using the metathesis reactions shown in Scheme 1.




Fig. 4. The packing of pairs of heterocyclium(1⁺) cations inside rhombs of eight $B_{12}F_{12}^{2-}$ anions in the structures of [1-Et-1-Me-imidazolium]₂[B₁₂F₁₂] (**1**, top), [1-H-3-Me-imidazolium]₂[B₁₂F₁₂] (**2**, bottom left), and [H(imidazolium)]₂[B₁₂F₁₂] (**3**, bottom right). The blue spheres are N atoms. The eight corners of the rhombs are composed of B₁₂ centroids. Each rhomb contains two complete cations or, in the case of **1**, parts of four cations that yield two complete cations. Structural drawings of these cations in salts **1-3** are seen in Fig. 2.




Fig. 5. The packing of pairs of heterocyclium(1+) cations inside rhombs of eight $B_{12}F_{12}^{2-}$ anions in the structures of $[4-NH_2-1-Me-1,2,4-triazolium]_2[B_{12}F_{12}]$ (**4**, top), $[4-NH_2-1-H-1,2,4-triazolium]_2[B_{12}F_{12}] \cdot 4-NH_2-1,2,4-triazole$ (**5**, middle), and $[1-NH_2-3-Me-1,2,3-triazolium]_2[B_{12}F_{12}]$ (**6**, bottom). The blue spheres are N atoms. The eight corners of the rhombs are composed of B_{12} centroids. Each rhomb contains two complete cations in **4** and **6**. In **5**, each rhomb contains parts of four triazolium cations and two triazole molecules that yield two complete cations and one complete triazole molecule (each five-membered ring is nearly centered on each of the six faces of the anion rhomb). Structural drawings of these cations in salts **4-6** are seen in Fig. 2.




Fig. 6. Plot of salt density and the shortest cat··cat distance for the five $B_{12}F_{12}^{2-}$ salts **1–4** and **6**. The line is a linear least-squares fit to the five points.




Fig. 7. The packing of 2,6-(NH₂)₂-1-H-purinium(1+) cations and H₂O molecules inside the rhomb of eight B₁₂F₁₂²⁻ anions in the structure of [2,6-(NH₂)₂-1-H-purinium]₂[B₁₂F₁₂]·2H₂O (**8**) and the packing of H(imidazolium)(1+) cations inside the rhomb of B₁₂F₁₂²⁻ anions in the structure of [H(imidazolium)]₂[B₁₂F₁₂] (**3**) (the corners of the rhombs are composed of B₁₂ centroids). The blue, red, green, and yellow spheres are N, O, B, and F atoms, respectively. These drawings emphasize the dovetail-like packing of the B₁₂F₁₂²⁻ anions. The ⊙⋯⊙ distances for the two anions shown are 7.423 Å for **8** and 7.122 Å for **3**. The angles between the two highlighted F–B–B–F planes in each drawing are 78.7° for **8** and 90° for **3**. The F⋯F distances shown are 2.779, 3.193, 3.689, and 3.796 Å in **8** and 2.908 (× 2) and 3.108 Å (× 2) in **3**. Structural drawings of these cations in salts **3** and **8** are seen in Fig. 2.




Fig. 8. The centrosymmetric $\text{Ag}_4(1\text{-NH}_2\text{-}1,2,3\text{-triazole})_8^{4+}$ tetracation in the structure of $[\text{Ag}_4(1\text{-NH}_2\text{-}1,2,3\text{-triazole})_8][\text{B}_{12}\text{F}_{12}]$ (**9**). The blue and purple atoms are N atoms and Ag^+ cations, respectively. The four Ag^+ ions are rigorously coplanar with $\text{Ag}\cdots\text{Ag}$ distances of 3.764(2) and 3.858(2) Å and $\text{Ag}\cdots\text{Ag}\cdots\text{Ag}$ angles of 65.8(1) and 114.2(1)°.

References

- [1] A. R. Pitochelli, M. F. Hawthorne, *J. Am. Chem. Soc.* 82 (1960) 3228–3229.
- [2] R. N. Grimes, *Angew. Chem., Int. Ed.* 42 (2003) 1198–1200, and references therein.
- [3] J. Aihara, *J. Am. Chem. Soc.* 100 (1978) 3339–3342.
- [4] R. B. King, *Chem. Rev.* 101 (2001) 1119–1152.
- [5] R.T. Sanderson, *Inorganic Chemistry*, 1st ed., Reinhold, New York, 1967.
- [6] K. A. Solntsev, A. M. Mebel, N. A. Votnova, N. T. Kuznetsov, O. P. Charkin, *Koord. Khim.* 18 (1992) 296–317.
- [7] S. V. Ivanov, S. M. Miller, O. P. Anderson, K. A. Solntsev, S. H. Strauss, *J. Am. Chem. Soc.* 125 (2003) 4694–4695.
- [8] S. A. Shackelford, J. L. Belletire, J. A. Boatz, S. Schneider, A. K. Wheaton, B. A. Wight, H. L. Ammon, D. V. Peryshkov, S. H. Strauss, *Org. Lett.* 12 (2010) 2714–2717.
- [9] D. V. Peryshkov, S. H. Strauss *J. Fluorine Chem.* 131 (2010) 1252–1256.
- [10] D. V. Peryshkov, A. A. Popov, S. H. Strauss, *J. Am. Chem. Soc.* 132 (2010) 13902–13913.
- [11] D. V. Peryshkov, E. Goresnik, Z. Mazej, S. H. Strauss, *J. Fluorine Chem.* 131 (2010) 1225–1228.
- [12] S. A. Shackelford, J. L. Belletire, J. A. Boatz, S. Schneider, A. K. Wheaton, B. A. Wight, L. M. Hudgens, H. L. Ammon, S. H. Strauss, *Org. Lett.* 11 (2009) 2623–2626.
- [13] J. L. Belletire, S. Schneider, B. A. Wight, S. L. Strauss, S. A. Shackelford, *Syn. Comm.* (2011) in press.
- [14] A. S. Larsen, J. D. Holbrey, F. S. Tham, C. A. Reed, *J. Am. Chem. Soc.* 122 (2000) 7264–7272.
- [15] Y. Zhu, C. Ching, K. Carpenter, R. Xu, S. Selvaratnam, N. S. Hosmane, J. A. Maguire, *Appl. Organometal. Chem.* 17 (2003), 346–350.
- [16] M. Nieuwenhuyzen, K. R. Seddon, F. Teixidor, A. V. Puga, *Inorg. Chem.* 48 (2009) 889–901.

- [17] (a) S. A. Shackelford, J. L. Belletire, *Centr. Europ. J. Energ. Mater.* 6 (2009) 219–237, (b) S. A. Shackelford, J. L. Belletire, 12th New Trends in Research on Energetic Materials (NTREM) International Seminar, Pardubice, Czech Republic, April 1–3, 2009.
- [18] (a) S. A. Shackelford, *Centr. Europ. J. Energ. Mater.* 5 (2008) 75–101, (b) S. A. Shackelford, 10th New Trends in Research on Energetic Materials (NTREM) International Seminar, Pardubice, Czech Republic, April 25–27, 2007.
- [19] D. V. Peryshkov, A. A. Popov, S. H. Strauss, *J. Am. Chem. Soc.* 131 (2009) 18393–18403.
- [20] Q.-L. Wang, H. Xu, H.-W. Hou, G. Yang, *Z. Naturforsch.* 64b (2009) 1143–1146.
- [21] B. Liu, L. Xu, G.-C. Guo, J.-S. Huang *Inorg. Chem. Commun.* 9 (2006) 687–690.
- [22] N. E. Brese, M. O'Keeffe *Acta Cryst B* 47 (1991) 192–197.
- [23] G. Kaplan, G. Drake, K. Tollison, L. Hall, T. Hawkins, *J. Heterocyclic Chem.* 42 (2005) 19–27.
- [24] D. V. Peryshkov, Ph.D. Dissertation, Colorado State University, 2011.
- [25] G. M. Sheldrick, SADABS, V. 2.10-A program for area detector absorption corrections; Bruker AXS: Madison, WI, 2003.
- [26] G. M. Sheldrick, APEX2, V. 2.0-2, Bruker AXS, Madison, WI, 2006.
- [27] G. M. Sheldrick, SHELXTL, V. 6.12 UNIX, Bruker AXS, Madison, WI, 2001.

Graphical Abstract

Pairing heterocyclic cations with *closo*-dodecafluorododecaborate(2⁻). Synthesis of binary heterocyclium(1⁺) salts and a Ag₄(heterocycle)₈⁴⁺ salt of B₁₂F₁₂²⁻

John L. Belletire^a, Stefan Schneider^b, Scott A. Shackelford^b, Dmitry V. Peryshkov^c, Steven H. Strauss^c

^aERC, Inc., AFRL/RZSP, Edwards AFB, CA 93524-7680, USA

^bAir Force Research Laboratory, AFRL/RZSP, Edwards AFB, CA 93524-7680, USA

^cDepartment of Chemistry, Colorado State University, Ft. Collins, CO 80523, USA

