

AWARD NUMBER: W81XWH-10-1-0181

TITLE: Induced Pluripotent Stem Cells as Potential Therapeutic Agents in NF1

PRINCIPAL INVESTIGATOR: Jonathan Chernoff, Ph.D.

CONTRACTING ORGANIZATION: Institute for Cancer Research
Philadelphia, PA 19111

REPORT DATE: May 2011

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 1 May 2011		2. REPORT TYPE Annual		3. DATES COVERED 1 May 2010 – 30 Apr 2011	
4. TITLE AND SUBTITLE Induced Pluripotent Stem Cells as Potential Therapeutic Agents in NF1				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-10-1-0181	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jonathan Chernoff, Ph.D. E-Mail: jonathan.chernoff@fcc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute for Cancer Research Philadelphia, PA 19111				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT In this project, we are seeking to use induced pluripotent stem (iPS) cell technology as a potential therapy in NF1. In the first year, we have successfully produced iPS cells from fibroblasts from Nf1+/- mice and characterized the properties of these cells, which include growth in clusters, expression of stem cell markers, normal karyotype, and the ability to form teratomas in mice. We have also created a targeting "gene-repair" vector to replace the defective Nf1 allele in these iPS cells. Thus, we are on schedule for this project, having achieved our aims in the expected time frame.					
15. SUBJECT TERMS NF1, iPS, Schwann cells, gene targeting					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

Introduction	4
Body	4
Key Research Accomplishments	6
Reportable Outcomes	6
Conclusion	6
References	6
Appendices	6

INTRODUCTION:

The goal of this project was to use induced pluripotent stem (iPS) cells and homologous recombination as the basis for therapy in neurofibromatosis 1 (NF1). Our hypothesis was that we could reprogram skin cells derived from NF1 heterozygous to become iPS cells, then repair the damaged NF1 allele, differentiate these iPS cells to hematopoietic precursors, and reintroduce such cells into irradiated *Krox20-Cre; Nf1^{fllox/-}* mice. The theory behind these experiments is that NF1-related tumors require not only *Nf1^{-/-}* Schwann cells, but also *Nf1^{+/-}* mast cells (1), and these could be replaced by the repaired iPS cells.

BODY: We set ourselves four specific tasks. These were:

Task 1. **Create iPS cells from *Nf1^{+/-}* keratinocytes.** In this step, we will obtain keratinocytes from *Nf1^{+/-}*; Oct4-GFP mice, which should express GFP when converted to iPS cells.

- 1a. Cross *Nf1^{+/-}* mice with Oct4-GFP mice to obtain *Nf1^{+/-}*; Oct4-GFP mice (months 1-4).
- 1b. Isolate and expand keratinocytes from 1-2 month old mice (months 4-6).
- 1c. Infect with four adenoviruses (encoding Oct4, Klf4, Sox2, and c-Myc, isolate and characterize iPS cells (months 6-12).

Task 2. **Repair the damaged *Nf1* allele.** We will use standard methods of homologous recombination to repair the damaged *Nf1* allele in *Nf1^{+/-}* iPS cells.

- 2a. Construct *Nf1* (re)targeting vector (months 1-6).
- 2b. Transfect *Nf1^{+/-}* iPS cells and identify targeted, *Nf1^{+/+}* iPS cells. (months 12-15).

Task 3. **Convert the iPS cells to hematopoietic precursors.** To convert the undifferentiated iPS cells to a transplantable state, we will use HoxB4 to drive the cells towards a hematopoietic lineage.

- 3a. Infect iPS cells with GFP-HoxB4 retrovirus (month 15).
- 3b. Isolate embryoid bodies and grow cells on OP9 stromal feeder layer. (months 15-18).

Task 4. **Transplant NF1 mouse with iPS-derived *Nf1^{+/+}* or *Nf1^{+/-}* cells.** We will replace the bone marrow cells of *Krox20-Cre; Nf1^{fllox/-}* mice with marrow derived from repaired iPS cells and observe the animals for signs of disease.

- 4a. Irradiate *Krox20-Cre; Nf1^{fllox/-}* and transplant with hematopoietic precursors derived from iPS cells. (months 18).
- 4b. Observe mice for signs of disease. (months 18-24).

Progress

In the first year, we have achieved the first aim, and are well on our way to achieving aim two, as detailed below.

Task 1. Create iPS cells from *Nf1*^{+/-} keratinocytes. We made several technical changes in the course of achieving this aim.

- 1a. Cross *Nf1*^{+/-} mice with Oct4-GFP mice to obtain *Nf1*^{+/-}; Oct4-GFP mice (months 1-4). We omitted this step, as efficiencies of iPS production are now such we did not need to use an Oct4 reporter. We therefore did not carry out this cross.
- 1b. Isolate and expand keratinocytes from 1-2 month old mice (months 4-6) _____. We also omitted this step, as we found that we could efficiently reprogram mouse fibroblasts.
- 1c. Infect with four adenoviruses (encoding Oct4, Klf4, Sox2, and c-Myc, isolate and characterize iPS cells (months 6-12)). We used retroviruses instead, as adenoviral transduction was not efficient in our hands. We used both the classic four factor (Oct4, Klf4, Sox2, c-Myc) and a three factor (Oct4, Klf4, Sox2) combinations to induce iPS formation. We found that the three-factor combination, which omits the potential oncogene c-Myc, gave robust, stem-cell appearing colonies (Fig. 1). Colony 3F-11, shown here, has a normal karyotype (not shown). This clone showed reactivation of endogenous stem cell genes, and formed embryoid bodies in vitro and teratomas *in vivo* (not shown).

Task 2. Repair the damaged *Nf1* allele. We are currently midway through this task, using standard methods of homologous recombination to repair the damaged *Nf1* allele in *Nf1*^{+/-} iPS cells.

- 2a. Construct *Nf1* (re)targeting vector (months 1-6). We used recombineering to create a “repair” allele (Fig. 2). This allele uses hygromycin for selection, and G418 sensitivity for counter-selection. That is, correctly targeted alleles, in which the damaged, neo-containing NF1 allele are replaced with our floxed hygromycin cassette, will be identified as colonies of iPS cells that are hygromycin resistant but G418 sensitive.
- 2b. Transfect *Nf1*^{+/-} iPS cells and identify targeted, *Nf1*^{+/+} iPS cells. (months 12-15). This subtask is ongoing.

Task 3. Convert the iPS cells to hematopoietic precursors. We have not initiated this task yet, as it cannot be begun until task two is complete.

Task 4. Transplant NF1 mouse with iPS-derived *Nf1*^{+/+} or *Nf1*^{+/-} cells. We have not initiated this task yet, as it cannot be begun until task three is complete.

KEY RESEARCH ACCOMPLISHMENTS:

- o Constructed pluripotent iPS cell lines from *Nf1*^{+/-} cells.
- o Completed retargeting construct to repair NF1 allele in these cells.

REPORTABLE OUTCOMES:

None to date.

CONCLUSION:

We are on schedule to complete our tasks. We made a number of changes along the way (e.g., use of fibroblasts in place of keratinocytes, use of retroviruses in place of adenoviruses, use of three reprogramming factors in place of four) to achieve successful production of pluripotent *Nf1*^{+/-} iPS cells. In addition, we have successfully constructed a gene repair plasmid, which will be used to complete task 2.

The production of iPS cells from NF1 animals also allows us to differentiate these cells *in vitro* to Schwann cells. While not a part of this proposal, it is likely that such cells would be a useful resource for studying the role of NF1 in Schwann cell differentiation and in drug sensitivity studies.

REFERENCES:

1. Yang, F.C., Ingram, D.A., Chen, S., Zhu, Y., Yuan, J., Li, X., Yang, X., Knowles, S., Horn, W., Li, Y., Zhang, S., Yang, Y., Vakili, S.T., Yu, M., Burns, D., Robertson, K., Hutchins, G., Parada, L.F., and Clapp, D.W., Nf1-dependent tumors require a microenvironment containing Nf1+/- and c-kit-dependent bone marrow. *Cell*, 135:437-448, 2008.

APPENDICES:

- 1) **Figure 1. Creation of iPS cells from *Nf1*^{+/-} fibroblasts using three factors**
- 2) **Figure 2. NF1 gene repair vector**

Figure 1. Creation of iPS cells from *Nf1*^{+/-} fibroblasts using three factors





Figure 2. NF1 gene repair vector**Figure 2. NF1 gene repair vector.** The vector is designed to replace the disrupted NF1 allele in this mouse model, restoring exon 31.