

FACTORIZATIONS AND REPRESENTATIONS OF BINARY POLYNOMIAL RECURRENCES BY MATRIX METHODS

EMRAH KILIC AND PANTELIMON STĂNICĂ

ABSTRACT. In this paper we derive factorizations and representations of a polynomial analogue of an arbitrary binary sequence by matrix methods. It generalizes various results on Fibonacci, Lucas, Chebyshev and Morgan–Voyce polynomials.

1. INTRODUCTION

In [10], the divisibility properties of the Fibonacci polynomial sequence $\{f_n(x)\}$ was studied. The Fibonacci polynomial sequence is defined by the recursion

$$f_{n+2}(x) = xf_{n+1}(x) + f_n(x); \quad f_0(x) = 0, \quad f_1(x) = 1.$$

Five years later, Hoggatt and Long [2] considered the general Fibonacci type polynomial sequence $\{u_n(x, y)\}$ of two variables. This sequence is defined by

$$(1.1) \quad u_{n+2}(x, y) = xu_{n+1}(x, y) + yu_n(x, y)$$

where $u_0(x, y) = 0$ and $u_1(x, y) = 1$.

The authors of [2, 10] found the roots of these polynomials and then obtained the factorizations of their polynomials. In [2], the authors found that

$$u_n(x, y) = y^{(n-1)/2} \prod_{k=1}^{n-1} \left(\frac{x}{\sqrt{y}} - 2i \cos \frac{k\pi}{n} \right).$$

Further, in [1], the authors use the relationships between the determinants of certain tridiagonal matrices and the Fibonacci and Lucas numbers, and then by matrix methods, they obtained the factorizations and representations of these sequences. The factorization of Fibonacci numbers was initially proposed in [6], and the factorization of Lucas numbers was obtained in [11].

The (companion) generalized Lucas polynomial sequence $v_n(x, y)$ is defined by

$$v_{n+2}(x, y) = xv_{n+1}(x, y) + yv_n(x, y)$$

where $v_0(x, y) = 2$ and $v_1(x, y) = 1$.

Recently, in [5], the binary sequential analogues of the generalized Fibonacci and Lucas polynomial sequences was considered and factorizations and representations of these sequences was obtained. These sequences are defined by

$$\begin{aligned} U_{n+1} &= AU_n + BU_{n-1} \\ V_{n+1} &= AV_n + BV_{n-1} \end{aligned}$$

2000 *Mathematics Subject Classification.* 11B37, 11C20.

Key words and phrases. Second order recurrences, factorization, Chebyshev polynomials, tridiagonal matrix, matrix methods.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Factorizations and Representations of Binary Polynomial Recurrences by Matrix Methods				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Applied Mathematics, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Rocky Mountain Journal of Mathematics 41:4 (2011), 1247-1264.					
14. ABSTRACT In this paper we derive factorizations and representations of a polynomial analogue of an arbitrary binary sequence by matrix methods. It generalizes various results on Fibonacci, Lucas, Chebyshev and Morgan{Voyce polynomials.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 12	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

where $U_0 = 0, U_1 = 1$ and $V_0 = 2, V_1 = A$, respectively. Also, in [4], we gave the more general factorizations of second order linear recurrences $\{U_n\}$ and $\{V_n\}$ with indices in arithmetic progressions. Furthermore, we obtained the factorization of these general sequences by the matrix methods considering how these recurrences are related to the determinants of certain tridiagonal matrices.

As it can be seen from the above mentioned results, the most general cases of these polynomial and binary sequences have not been studied. Consequently, we define $\{A_n(a, b; p, q)(x)\}$ (we shall often drop the argument $(a, b; p, q)$ and simply write $\{A_n(x)\}$) to be a polynomial sequence satisfying

$$A_{n+1}(x) = p(x)A_n(x) + q(x)A_{n-1}(x),$$

and initial conditions $A_0 = a(x), A_1 = b(x)$, where a, b, p, q are polynomials of an indeterminate x with real coefficients. For easy notation, we shall sometimes write A_n, p, q, a, b for $A_n(x), p(x), q(x), a(x)$ and $b(x)$, respectively. We display some special cases of the sequence $\{A_n\}$ in the following table.

$p(x)$	$q(x)$	$a(x)$	$b(x)$	$A_n(a, b; p, q)(x)$	Polynomial Type
$2x$	-1	1	x	$T_n(x)$	1st kind Chebyshev
$2x$	-1	1	$2x$	$U_n(x)$	2nd kind Chebyshev
$x+1$	-1	1	$x+1$	$b_n(x)$	Morgan-Voyce
x	y	0	1	$u_n(x, y)$	generalized Fibonacci
x	y	2	1	$v_n(x, y)$	generalized Lucas
$2x$	-1	1	$2x+1$	$W_n(x)$	4th kind Chebyshev
$2x$	-1	1	$2x-1$	$V_n(x)$	3rd kind Chebyshev

Table 1

In Section 2 we present a recurrence, and define a tridiagonal matrix whose determinant is precisely $A_n(a, b; p, q)(x)$, with n in an arithmetic progression, $n \equiv c \pmod{k}$. In our main Section 3 we derive the factorization and representations of the sequence $\{A_n(a, b; p, q)\}$ (under some assumptions), by matrix methods, thus generalizing some results of [1]–[5] and others. As consequences, we obtain the factorizations for the Chebyshev's and generalized Lucas polynomials, among others.

2. A RECURRENCE FOR $A_n(a, b; p, q)(x)$, WHERE $n \equiv c \pmod{k}$

We start this section with the Binet formulas of both positively and negatively indexed terms of the sequence $\{A_n\}$, namely

$$(2.1) \quad A_{\pm n} = \left(\frac{b - a\beta}{\alpha - \beta} \right) \alpha^{\pm n} + \left(\frac{a\alpha - b}{\alpha - \beta} \right) \beta^{\pm n}$$

where

$$(2.2) \quad \alpha = \frac{p + \sqrt{p^2 + 4q}}{2}, \quad \beta = \frac{p - \sqrt{p^2 + 4q}}{2}.$$

First, we prove the following lemma.

Lemma 1. *For $k > 0, n > 0$, the sequence $\{A_n\}$ satisfies the following recursion*

$$(2.3) \quad A_{p(n+1, \pm k, c)} = y_{\pm k} A_{p(n, \pm k, c)} - z_{\pm k} A_{p(n-1, \pm k, c)}$$

where $y_{\pm k} = \alpha^{\pm k} + \beta^{\pm k}$, $z_{\pm k} = q^{\pm k}$ and $p(n, k, c) = nk + c$ (c constant).

Proof. From the definition of the sequence $\{A_n\}$, considering the case $A_0 = 2$ and $A_1 = p$, we write

$$y_{\pm k} = A_{\pm k}(2, p; p, q) = \alpha^{\pm k} + \beta^{\pm k}$$

where α and β are given by (2.2). Further, for the positive case, we note that

$$y_k = A_k(2, p; p, q) = \alpha^k + \beta^k.$$

Since the positively and negatively indexed terms cases are similar, we only consider the positively indexed terms case. By the Binet formula of the sequence $\{A_n\}$ and since $z_k = q^k = (\alpha\beta)^k$,

$$\begin{aligned} & y_k A_{p(n,k,c)} - z_k A_{p(n-1,k,c)} \\ &= (\alpha^k + \beta^k) \left[\left(\frac{b-a\beta}{\alpha-\beta} \right) \alpha^{kn+c} + \left(\frac{a\alpha-b}{\alpha-\beta} \right) \beta^{kn+c} \right] \\ & \quad - (\alpha\beta)^k \left[\left(\frac{b-a\beta}{\alpha-\beta} \right) \alpha^{k(n-1)+c} + \left(\frac{a\alpha-b}{\alpha-\beta} \right) \beta^{k(n-1)+c} \right] \\ &= \frac{1}{\alpha-\beta} ((b-a\beta) \alpha^{k(n+1)+c} + (a\alpha-b) \beta^{k(n+1)+c} \\ & \quad + (b-a\beta) \alpha^{kn+c} \beta^k + (a\alpha-b) \beta^{kn+c} \alpha^k \\ & \quad - (b-a\beta) \alpha^{kn+c} \beta^k - (a\alpha-b) \beta^{kn+c} \alpha^k) \\ &= \left(\frac{b-a\beta}{\alpha-\beta} \right) \alpha^{k(n+1)+c} + \left(\frac{a\alpha-b}{\alpha-\beta} \right) \beta^{k(n+1)+c} \\ &= A_{p(n+1,k,c)}, \end{aligned}$$

which proves the lemma. $\square$

Now we present a relationship between the terms $A_{p(\pm(n+1),k,c)}$ and the determinant of a certain tridiagonal matrix.

Define the $n \times n$ tridiagonal matrix $M_{n,k}$ as shown:

$$(2.4) \quad M_{n,\pm k} = \begin{bmatrix} A_{p(1,\pm k,c)} & A_{p(0,\pm k,c)} \sqrt{z_{\pm k}} & & & \\ \sqrt{z_{\pm k}} & y_{\pm k} & \sqrt{z_{\pm k}} & & \\ & \sqrt{z_{\pm k}} & y_{\pm k} & \ddots & \\ & & \ddots & \ddots & \sqrt{z_{\pm k}} \\ & & & \sqrt{z_{\pm k}} & y_{\pm k} \end{bmatrix}.$$

Theorem 1. For $n > 1$, we have

$$\det M_{n,\pm k} = A_{p(n,\pm k,c)}.$$

Proof. As before, we only consider the positively indexed terms. The other case can be similarly derived. Expanding $\det M_{n,k}$ using the cofactor expansion of a determinant with respect to the last column, we obtain

$$\det M_{n,k} = y_k \det M_{n-1,k} - z_k \det M_{n-2,k}.$$

Replacing $n = 1$ in equation (2.3), we obtain $A_{p(2,k,c)} = y_k A_{p(1,k,c)} - z_k A_{p(0,k,c)}$ and $\det M_{2,k} = y_k A_{p(1,k,c)} - z_k A_{p(0,k,c)}$, and so, $\det M_{2,k} = A_{p(2,k,c)}$. Obviously, $\det M_{1,k} = A_{p(1,k,c)}$. Since the recurrence relations (and initial conditions) of $\det M_{n,k}$ and the sequence $\{A_{p(n,k,c)}\}$ are the same, the conclusion follows from Lemma 1. $\square$

3. FACTORIZATIONS FOR $A_n(a, b; p, q)(x)$

Now we investigate all possible cases of our main general considerations and then we give their factorizations and representations. However, a few special cases could be treated separately. For compactness, we do not consider all these cases, but we shall point out, whenever appropriate some hints in treating those cases.

First, we consider the case $A_{p(1, \pm k, c)} = y_{\pm k}$ and $A_{p(0, \pm k, c)} = 1$. Under these assumptions, we label the matrix $M_{n, \pm k}$ and the sequence $A_{p(\pm n, k, c)}$ by $M_{n, \pm k}^{(1)}$ and $A_{p(n, \pm k, c)}^{(1)}$, respectively. By Theorem 1, the matrix $M_{n, \pm k}^{(1)}$ takes the following form:

$$M_{n, \pm k}^{(1)} = \begin{bmatrix} y_{\pm k} & \sqrt{z_{\pm k}} & & \\ \sqrt{z_{\pm k}} & \sqrt{z_{\pm k}} & \ddots & \\ & \ddots & \ddots & \sqrt{z_{\pm k}} \\ & & \sqrt{z_{\pm k}} & y_{\pm k} \end{bmatrix}$$

and we have that

$$(3.1) \quad \det M_{n, \pm k}^{(1)} = A_{p(n, \pm k, c)}^{(1)}.$$

Let Q_1 be the following $(n \times n)$ tridiagonal Toeplitz matrix

$$Q_1 = \begin{bmatrix} 0 & 1 & & \\ 1 & 0 & \ddots & \\ & \ddots & \ddots & 1 \\ & & 1 & 0 \end{bmatrix}.$$

The characteristic equation of the matrix Q_1 satisfies the following recursion, for $n > 2$

$$f_n(\lambda) = -\lambda f_{n-1}(\lambda) - f_{n-2}(\lambda),$$

where $f_1(\lambda) = -\lambda$ and $f_2(\lambda) = \lambda^2 - 1$.

Taking $\lambda \equiv -2x$, the family $\{f_n(\lambda)\}$ is reduced to the family $\{U_n(x)\}$ (Chebyshev polynomial of the second kind). From [9, 8, 7], the zeros of the Chebyshev polynomials are known, and so, the eigenvalues of Q_1 are of the form:

$$(3.2) \quad \lambda_r = -2 \cos \frac{\pi r}{n+1}, \quad r = 1, 2, \dots, n.$$

We can also write $M_{n, \pm k}^{(1)} = y_{\pm k} I_n + \sqrt{z_{\pm k}} Q_1$, where I_n is the $n \times n$ unit matrix.

Theorem 2. *Let $A_{p(1, \pm k, c)} = y_{\pm k}$ and $A_{p(0, \pm k, c)} = 1$. Then for $n > 1$,*

$$A_{p(n, \pm k, c)}^{(1)} = \prod_{r=1}^n \left(y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{\pi r}{n+1} \right).$$

Proof. Assume that λ_r 's are the eigenvalues of the matrix Q_1 with respect to the eigenvectors w_r . Since

$$\begin{aligned} M_{n, \pm k}^{(1)} w_r &= (y_{\pm k} I_n + \sqrt{z_{\pm k}} Q_1) w_r = y_{\pm k} I_n w_r + \sqrt{z_{\pm k}} Q_1 w_r \\ &= (y_{\pm k} + \sqrt{z_{\pm k}} \lambda_r) w_r, \end{aligned}$$

the $(y_{\pm k} + \sqrt{z_{\pm k}}\lambda_r)$'s are the eigenvalues of the matrix $M_{n,\pm k}^{(1)}$ with respect to the eigenvectors w_r . Thus, by (3.2) and (3.1), we have the conclusion:

$$\det M_{n,\pm k}^{(1)} = A_{p(n,\pm k,c)}^{(1)} = \prod_{r=1}^n (y_{\pm k} + \sqrt{z_{\pm k}}\lambda_r) = \prod_{r=1}^n \left(y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{\pi r}{n+1} \right).$$

□

Corollary 1. For $n > 1$,

$$A_{p(n,\pm k,c)}^{(1)} = \begin{cases} y_{\pm k} \prod_{r=1}^{n/2} \left(y_{\pm k}^2 - 4z_{\pm k} \cos^2 \frac{\pi r}{n+1} \right) & \text{if } n \text{ is even,} \\ \prod_{r=1}^{(n-1)/2} \left(y_{\pm k}^2 - 4z_{\pm k} \cos^2 \frac{\pi r}{n+1} \right) & \text{if } n \text{ is odd.} \end{cases}$$

Proof. Since $\cos(k\pi/n) = -\cos((n-k)\pi/n)$ for $1 \leq k \leq n/2$, the conclusion follows from Theorem 2. □

Now we give some applications of Theorem 2 in the following corollaries.

Corollary 2. Let U_n be n th Chebyshev polynomial of the second kind. For $n, k > 1$, then

$$(3.3) \quad U_{\pm(n+1)k-1} = 2^n U_{\pm k-1} \prod_{r=1}^n \left(T_{\pm k} - \cos \frac{\pi r}{n+1} \right).$$

where T_n is n th Chebyshev polynomial of the first kind.

Proof. When $p = 2x, q = -1$, then the first coefficient $y_{\pm k} = 2T_{\pm k}$, where T_k is the k th Chebyshev polynomials of the first kind (see Table 1). According to the first case, to satisfy $A_{p(1,\pm k,c)}^{(1)} = y_{\pm k} = 2T_{\pm k}$ and $A_{p(0,\pm k,c)}^{(1)} = 1$, we shall choose $a = 1$ and $b = 2x$. We see that for $c = -1$,

$$A_{p(1,\pm k,-1)}^{(1)} = \frac{U_{\pm 2k-1}}{U_{\pm k-1}} = 2T_{\pm k} \text{ and } A_{p(0,\pm k,-1)}^{(1)} = \frac{U_{\pm k-1}}{U_{\pm k-1}} = 1$$

and in general

$$A_{p(n,\pm k,-1)}^{(1)} = \frac{U_{\pm(n+1)k-1}}{U_{\pm k-1}}.$$

Then by Theorem 2, we get for $y_k = 2T_k$ and $z_k = 1$

$$(3.4) \quad A_{p(n,\pm k,c)}^{(1)} = \frac{U_{\pm(n+1)k-1}}{U_{\pm k-1}} = \prod_{r=1}^n \left(2T_{\pm k} - 2 \cos \frac{\pi r}{n+1} \right).$$

The required equation (3.3) follows immediately. □

Corollary 3. For $n, k > 1$, then

$$U_{\pm k(n+1)-1} = \begin{cases} 2^{n-1} U_{\pm 2k-1} \prod_{r=1}^{(n-1)/2} \left[T_{\pm k}^2 - \cos^2 \left(\frac{\pi r}{n+1} \right) \right] & \text{if } n \text{ is odd,} \\ 2^n U_{\pm k-1} \prod_{r=1}^{n/2} \left[T_{\pm k}^2 - \cos^2 \left(\frac{\pi r}{n+1} \right) \right] & \text{if } n \text{ is even.} \end{cases}$$

Proof. The proof follows from Corollary 1 and equation (3.4). □

Corollary 4. *Let $u_n(x, t)$ be the generalized Fibonacci polynomial. For $n > 1$ and $k > 0$,*

$$(3.5) \quad u_{\pm(n+1)k}(x, t) = u_{\pm k}(x, t) \prod_{r=1}^n \left(v_{\pm k}(x, t) - 2\sqrt{(-t)^{\pm k}} \cos \frac{\pi r}{n+1} \right).$$

Proof. When $p = x, q = t$, then the first coefficient $y_{\pm k} = v_{\pm k}(x, t)$ where v_k is the k th term of the generalized Lucas polynomial sequence (see Table 1). For our first case, to satisfy $A_{p(1, \pm k, c)}^{(1)} = y_{\pm k} = v_{\pm k}(x, t)$ and $A_{p(0, \pm k, c)}^{(1)} = 1$, we let $a = 0$ and $b = 1$. For $c = 0$, we get

$$A_{p(1, \pm k, 0)}^{(1)} = \frac{u_{\pm 2k}(x, t)}{u_{\pm k}(x, t)} = v_{\pm k}(x, t) \text{ and } A_{p(0, \pm k, 0)}^{(1)} = \frac{u_{\pm k}(x, t)}{u_{\pm k}(x, t)} = 1$$

and in general

$$A_{p(n, \pm k, 0)}^{(1)} = \frac{u_{\pm k(n+1)}}{u_{\pm k}}.$$

Then by Theorem 2, we obtain for $y_{\pm k} = v_{\pm k}(x, t)$ and $z_k = (-t)^k$,

$$(3.6) \quad A_{p(n, \pm k, 0)}^{(1)} = \frac{u_{\pm(n+1)k}}{u_{\pm k}} = \prod_{r=1}^n \left(v_{\pm k} - 2\sqrt{(-t)^{\pm k}} \cos \frac{\pi r}{n+1} \right)$$

and so, we have the required equation (3.5). $\square$

Indeed, when $k = 1$, then by the above corollary, we get that $u_1 = 1$, $v_1 = x$ and so

$$u_n(x, t) = t^{(n-1)/2} \prod_{r=1}^{n-1} \left(\frac{x}{\sqrt{t}} - 2i \cos \frac{r\pi}{n} \right)$$

which is given in [2]. Thus it can be seen that our result generalizes earlier work.

Corollary 5. *For $n > 1$ and $k > 0$,*

$$u_{(n+1)k} = \begin{cases} u_k \prod_{r=1}^{n/2} \left(v_k^2 - 4y \cos^2 \frac{\pi r}{n+1} \right) & \text{if } n \text{ is even,} \\ u_{2k} \prod_{r=1}^{(n-1)/2} \left(v_k^2 - 4y \cos^2 \frac{\pi r}{n+1} \right) & \text{if } n \text{ is odd.} \end{cases}$$

Proof. Since $u_{2n} = u_n v_n$, by Corollary 1 and equation (3.6), we have the conclusion. $\square$

Next, we consider both binary and polynomial sequences satisfying $A_{p(1, \pm k, c)} = y_{\pm k}$ and $A_{p(0, \pm k, c)} = 2$. In this case, we label the matrix $M_{n, \pm k}$ and the general sequence $A_{p(n, \pm k, c)}$ by $M_{n, \pm k}^{(2)}$ and $A_{p(n, \pm k, c)}^{(2)}$, respectively.

Theorem 1 implies the following facts. For

$$M_{n, \pm k}^{(2)} = \begin{bmatrix} y_{\pm k} & 2\sqrt{z_{\pm k}} & & & \\ \sqrt{z_{\pm k}} & y_{\pm k} & \sqrt{z_{\pm k}} & & \\ & \sqrt{z_{\pm k}} & y_{\pm k} & \ddots & \\ & & \ddots & \ddots & \sqrt{z_{\pm k}} \\ & & & \sqrt{z_{\pm k}} & y_{\pm k} \end{bmatrix},$$

we have

$$(3.7) \quad \det M_{n, \pm k}^{(2)} = A_{p(n, \pm k, c)}^{(2)}.$$

The $(n \times n)$ tridiagonal matrix Q_2 as shown:

$$Q_2 = \begin{bmatrix} 0 & 2 & & & \\ 1 & 0 & 1 & & \\ & 1 & 0 & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & 0 \end{bmatrix}.$$

The characteristic polynomial of matrix Q_n satisfies the following recurrence relation: for $n > 2$

$$t_n(\delta) = -\delta t_{n-1}(\delta) - t_{n-2}(\delta)$$

where $t_1(\delta) = -\delta$, $t_2(\delta) = \delta^2 - 2$.

Taking $\delta \equiv -2x$, the family $\{t_n(\delta) : n = 1, 2, \dots\}$ is reduced to the family of Chebyshev polynomials of the first kind $\{2T_n(x) : n = 1, 2, \dots\}$. From [9, 8, 7], since the zeros of Chebyshev polynomials of the first kind are known, we obtain

$$(3.8) \quad \delta_r = -2 \cos \frac{(2r-1)\pi}{2n}, \quad r = 1, 2, \dots, n.$$

We can also write $M_{n,k}^{(2)} = y_k I_n + \sqrt{z_k} Q_2$, where I_n is the unit matrix of order n .

Theorem 3. Let $A_{p(1,\pm k,c)} = y_{\pm k}$ and $A_{p(0,\pm k,c)} = 2$. For $n > 1$, $k > 0$,

$$A_{p(n,\pm k,c)}^{(2)} = \prod_{r=1}^n \left[y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{(2r-1)\pi}{2n} \right].$$

Proof. If the eigenvalues of Q_2 are δ_r 's with respect to the eigenvectors w_r , then

$$M_{n,k}^{(2)} w_r = (y_k I_n + \sqrt{z_k} Q_2) w_r = y_k I_n w_r + \sqrt{z_k} Q_2 w_r = (y_k + \sqrt{z_k} \delta_r) w_r.$$

Thus the eigenvalues of $M_{n,\pm k}^{(2)}$ are the $y_{\pm k} + \sqrt{z_{\pm k}} \delta_r$'s with respect to the eigenvectors w_r . By (3.8),

$$\det M_{n,\pm k}^{(2)} = \prod_{r=1}^n (y_{\pm k} + \sqrt{z_{\pm k}} \delta_r) = \prod_{r=1}^n \left[y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{(2r-1)\pi}{2n} \right].$$

By (3.7), the theorem is proven. $\square$

Corollary 6. For $n > 1$, $k > 0$,

$$A_{p(n,k,c)}^{(2)} = \begin{cases} y_k \prod_{r=1}^{(n-1)/2} \left[y_k^2 - 4z_k \cos^2 \left(\frac{(2r-1)\pi}{2n} \right) \right] & \text{if } n \text{ is odd,} \\ \prod_{r=1}^{n/2} \left[y_k^2 - 4z_k \cos^2 \left(\frac{(2r-1)\pi}{2n} \right) \right] & \text{if } n \text{ is even.} \end{cases}$$

Now we give some applications of the above results for some well known sequences.

Corollary 7. Let $v_n(x, t)$ be the generalized Lucas polynomial sequence. Then for $n > 1$, $k > 0$,

$$v_{\pm kn}(x, t) = \prod_{r=1}^n \left(v_{\pm k}(x, t) - 2\sqrt{(-t)^{\pm k}} \cos \frac{(2r-1)\pi}{2n} \right).$$

Proof. When $p = x, q = t$, then the first coefficient $y_{\pm k} = v_{\pm k}(x, t)$. For our second case and satisfying $A_{p(1, \pm k, c)}^{(2)} = y_{\pm k} = v_{\pm k}$ and $A_{p(0, \pm k, c)}^{(2)} = 2$, letting $a = 2$ and $b = x$, we see that for $c = 0$

$$A_{p(1, \pm k, 0)}^{(2)} = v_{\pm k} \text{ and } A_{p(0, \pm k, 0)}^{(2)} = x$$

and in general $A_{p(n, \pm k, c)}^{(2)} = v_{\pm kn}$. By Theorem 3, we have the conclusion: for $y_{\pm k} = v_{\pm k}$ and $z_{\pm k} = (-t)^{\pm k}$,

$$A_{p(n, \pm k, 0)}^{(2)} = v_{\pm kn}(x, t) = \prod_{r=1}^n \left(v_{\pm k}(x, t) - 2\sqrt{(-t)^{\pm k}} \cos \frac{(2r-1)\pi}{2n} \right).$$

□

Corollary 8. *Let $v_n(x, t)$ be the generalized Lucas polynomial sequence. Then for $n > 1, k > 0$,*

$$v_{\pm kn}(x, t) = \begin{cases} v_{\pm k}(x, t) \prod_{r=1}^{(n-1)/2} \left[v_{\pm k}(x, t)^2 - 4(-t)^{\pm k} \cos^2 \left(\frac{(2r-1)\pi}{2n} \right) \right] & \text{if } n \text{ is odd,} \\ \prod_{r=1}^{n/2} \left[v_{\pm k}(x, t)^2 - 4(-t)^{\pm k} \cos^2 \left(\frac{(2r-1)\pi}{2n} \right) \right] & \text{if } n \text{ is even.} \end{cases}$$

For the Chebyshev polynomial of the first kind, if we consider $v_n(2x, -1)$ in the above two corollaries, then we see that $v_n(2x, -1) = 2T_n$, so

$$2T_{\pm kn} = v_{\pm kn}(2x, -1) = \prod_{r=1}^n \left(2T_{\pm k} - 2 \cos \frac{(2r-1)\pi}{2n} \right)$$

and clearly

$$T_{\pm kn} = 2^{n-1} \prod_{r=1}^n \left(T_{\pm k} - \cos \frac{(2r-1)\pi}{2n} \right).$$

In this third case, we let $A_{p(1, \pm k, c)} = y_{\pm k} = -1$ and $A_{p(0, \pm k, c)} = 1$. Further, we label the matrix $M_{n, \pm k}$ and the sequence $A_{p(n, \pm k, c)}$ by $M_{n, \pm k}^{(3)}$ and $A_{p(n, \pm k, c)}^{(3)}$, respectively.

Theorem 1 implies the following facts. For the $n \times n$ tridiagonal matrix $M_{n, k}^{(3)}$:

$$M_{n, \pm k}^{(3)} = \begin{bmatrix} y_{\pm k} - 1 & \sqrt{z_{\pm k}} & & & \\ \sqrt{z_{\pm k}} & y_{\pm k} & \sqrt{z_{\pm k}} & & \\ & \sqrt{z_{\pm k}} & y_{\pm k} & \ddots & \\ & & \ddots & \ddots & \sqrt{z_{\pm k}} \\ & & & \sqrt{z_{\pm k}} & y_{\pm k} \end{bmatrix},$$

by Theorem 1, we have for $n > 1$

$$(3.9) \quad \det M_{n, \pm k}^{(3)} = A_{p(n, \pm k, c)}^{(3)}.$$

Define the $(n \times n)$ tridiagonal matrix Q_3 as shown:

$$Q_3 = \begin{bmatrix} -1 & 1 & & & \\ 1 & 0 & 1 & & \\ & 1 & 0 & \ddots & \\ & & \ddots & \ddots & 1 \\ & & & 1 & 0 \end{bmatrix}.$$

Thus the characteristic polynomial of the matrix Q_3 satisfies the following recurrence relation (for $n > 2$)

$$g_n(\mu) = -\mu g_{n-1}(\mu) - g_{n-2}(\mu)$$

where $g_1(\mu) = -\mu - 1$, $g_2(\mu) = \mu^2 + \mu - 1$.

If we take $\delta \equiv -2x$, then the family $\{g_n(\mu) : n = 1, 2, \dots\}$ is reduced to the family of Chebyshev polynomials of the third kind $\{V_n(x) : n = 1, 2, \dots\}$. From [9, 8, 7], the zeros of Chebyshev polynomials of the third kind are known, then we obtain

$$(3.10) \quad \mu_k = -2 \cos \frac{(k - \frac{1}{2})\pi}{n + \frac{1}{2}}, \quad k = 1, 2, \dots, n.$$

We can also write $M_{n,\pm k}^{(3)} = y_{\pm k}I_n + \sqrt{z_{\pm k}}Q_3$.

Theorem 4. Let $A_{p(1,\pm k,c)} - y_{\pm k} = -1$ and $A_{p(0,\pm k,c)} = 1$. For $n > 1$, $k > 0$,

$$A_{p(n,\pm k,c)}^{(3)} = \prod_{r=1}^n \left(y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{(2k-1)\pi}{2n+1} \right).$$

Proof. Let the eigenvalues of the matrix Q_3 to be μ_r with respect to the eigenvectors w_r , and write

$$M_{n,\pm k}^{(3)} w_r = (y_{\pm k}I_n + \sqrt{z_{\pm k}}Q_3) w_r = (y_{\pm k} + \sqrt{z_{\pm k}}\mu_r) w_r.$$

Thus the eigenvalues of $M_{n,\pm k}^{(3)}$ are $(y_{\pm k} + \sqrt{z_{\pm k}}\mu_r)$. By equations (3.10) and (3.9), we have

$$\det M_{n,\pm k}^{(3)} = A_{p(n,\pm k,c)}^{(3)} = \prod_{r=1}^n \left(y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{(2k-1)\pi}{2n+1} \right).$$

Thus the theorem is proven. $\square$

The following corollary follows directly from Theorem 4 since, for $1 \leq k \leq n/2$, we have $\cos \left(\frac{k\pi}{n} \right) = -\cos \left(\frac{(n-k)\pi}{n} \right)$.

Corollary 9. For $n > 1$, $k > 0$,

$$A_{p(n,\pm k,c)}^{(3)} = \begin{cases} y_{\pm k} \prod_{r=1}^{(n-1)/2} \left(y_{\pm k}^2 - 4z_{\pm k} \cos^2 \left(\frac{(2k-1)\pi}{2n+1} \right) \right) & \text{if } n \text{ is odd,} \\ \prod_{r=1}^{n/2} \left(y_{\pm k}^2 - 4z_{\pm k} \cos^2 \left(\frac{(2k-1)\pi}{2n+1} \right) \right) & \text{if } n \text{ is even.} \end{cases}$$

A particular case of the previous corollary can be found in literature.

Corollary 10. For $n > 1$,

$$(3.11) \quad T_{4n+2} = 2^n T_2 \prod_{k=1}^n \left(T_4 - \cos \frac{(2k-1)\pi}{2n+1} \right).$$

Proof. For $k = 2$, $p = 2x, q = -1$, the first coefficient is $y_4 = 2T_4$ (Chebyshev polynomial of the first kind). Also under the conditions $a = 1$ and $b = 2x$, we obtain $A_{p(1,4,-2)}^{(3)} = y_4 = 2T_4$ and $A_{p(0,4,-2)}^{(3)} = 1$,

$$A_{p(1,4,-2)}^{(3)} = \frac{T_6}{T_2} = 2T_4 - 1 \text{ and } A_{p(0,4,-2)}^{(3)} = \frac{T_2}{T_2} = 1$$

and in general

$$A_{p(n,4,-2)}^{(3)} = \frac{T_{4(n+1)-2}}{T_2}.$$

Then by Theorem 4, we get for $y_4 = 2T_4$ and $z_4 = 1$

$$(3.12) \quad A_{p(n,4,-2)}^{(3)} = \frac{T_{4(n+1)-1}}{T_2} = \prod_{r=1}^n \left(2T_4 - 2 \cos \frac{(2k-1)\pi}{2n+1} \right)$$

and so equation (3.11) is obtained. $\square$

By Corollary 9 and the above result, we have

$$T_{4n+2} = \begin{cases} 2^n T_4 T_2 \prod_{r=1}^{(n-1)/2} \left(T_4^2 - \cos^2 \left(\frac{(2k-1)\pi}{2n+1} \right) \right) & \text{if } n \text{ is odd,} \\ 2^n T_2 \prod_{r=1}^{n/2} \left(T_4^2 - \cos^2 \left(\frac{(2k-1)\pi}{2n+1} \right) \right) & \text{if } n \text{ is even.} \end{cases}$$

Further, we consider our final case $A_{p(1,\pm k,c)} - y_{\pm k} = 1$ and $A_{p(0,\pm k,c)} = 1$. In this case, we label the matrix $M_{n,\pm k}$ and the sequence $A_{p(n,\pm k,c)}$ by $M_{n,\pm k}^{(4)}$ and $A_{p(n,\pm k,c)}^{(4)}$, respectively. By Theorem 1, we have

$$M_{n,\pm k}^{(4)} = \begin{bmatrix} y_{\pm k} + 1 & \sqrt{z_{\pm k}} & & & \\ \sqrt{z_{\pm k}} & y_{\pm k} & \sqrt{z_{\pm k}} & & \\ & \sqrt{z_{\pm k}} & y_{\pm k} & \ddots & \\ & & \ddots & \ddots & \sqrt{z_{\pm k}} \\ & & & \sqrt{z_{\pm k}} & y_{\pm k} \end{bmatrix},$$

and for $n > 1$

$$(3.13) \quad \det M_{n,\pm k}^{(4)} = A_{p(n,\pm k,c)}^{(4)}.$$

We also define the $(n \times n)$ matrix Q_4 as follows:

$$Q_4 = \begin{bmatrix} 1 & 1 & & & \\ 1 & 0 & 1 & & \\ & 1 & 0 & \ddots & \\ & & 1 & \ddots & 1 \\ & & & 1 & 0 \end{bmatrix}.$$

Thus the characteristic polynomial of Q_4 satisfies the recurrence

$$h_{n+1}(\omega) = -\omega h_n(\omega) - h_{n-1}(\omega)$$

where $h_1(\omega) = 1 - \omega$ and $h_2(\omega) = \omega^2 + \omega + 1$.

Taking $\omega \equiv -2x$, the family $\{h_n(\omega)\}$ is converted into the family of Chebyshev polynomials of the fourth kind $\{W_n(x) : n = 1, 2, \dots\}$. By [9, 8, 7], the zeros of Chebyshev polynomials of the fourth kind are known, namely

$$(3.14) \quad \omega_k = -2 \cos \frac{(k-1)\pi}{n + \frac{1}{2}}, \quad k = 1, 2, \dots, n.$$

The matrix $M_{n,\pm k}^{(4)}$ can be expressed as $y_{\pm k}I_n + \sqrt{z_{\pm k}}Q_4$.

Theorem 5. Let $A_{p(1,\pm k,c)} - y_{\pm k} = 1$ and $A_{p(0,\pm k,c)} = 1$. For $n > 1$, $k > 0$,

$$A_{p(n,\pm k,c)}^{(4)} = \prod_{r=1}^n \left(y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{2(r-1)\pi}{2n+1} \right).$$

Proof. If ω_r 's are the eigenvalues of $M_{n,\pm k}^{(4)}$ according to the eigenvectors w_r , then

$$M_{n,\pm k}^{(4)} w_r = (y_{\pm k}I_n + \sqrt{z_{\pm k}}Q_4) w_r = y_{\pm k} + \sqrt{z_{\pm k}}\omega_r.$$

Thus the eigenvalues of $M_{n,\pm k}^{(4)}$ are $(y_{\pm k} + \sqrt{z_{\pm k}}\omega_r)$ and by (3.13), (3.14), we obtain the conclusion:

$$\det M_{n,\pm k}^{(4)} = \prod_{r=1}^n \left(y_{\pm k} - 2\sqrt{z_{\pm k}} \cos \frac{2(r-1)\pi}{2n+1} \right).$$

□

As a consequence of Theorem 5, we have the following corollary.

Corollary 11. For $n > 1$, $k > 0$,

$$A_{p(n,\pm k,c)}^{(4)} = \begin{cases} y_{\pm k} \prod_{r=1}^{(n-1)/2} \left(y_{\pm k}^2 - 4z_{\pm k} \cos^2 \left(\frac{2(r-1)\pi}{2n+1} \right) \right) & \text{if } n \text{ is odd,} \\ \prod_{r=1}^n \left(y_{\pm k}^2 - 4z_{\pm k} \cos^2 \left(\frac{2(r-1)\pi}{2n+1} \right) \right) & \text{if } n \text{ is even.} \end{cases}$$

REFERENCES

- [1] Cahill, N.D., D'Errico, J.R., Spence, J.P.: Complex factorizations of the Fibonacci and Lucas numbers, *Fibonacci Quart.* 41 (2003), 13–19.
- [2] Hoggatt, Jr.V., Long, C.: Divisibility properties of generalized Fibonacci polynomials, *Fibonacci Quart.* 12 (1974), 113–120.
- [3] Horadam, A.F.: Special properties of the sequence $W_n(a, b; p, q)$. *Fibonacci Quart.* 5 (1967), 424–434.
- [4] Kiliç, E., Stănică, P.: Factorizations and representations of second order linear recurrences with indices in arithmetic progressions, submitted for publication.
- [5] Kilic, E., Tasci, D.: Factorizations and representations of the backward second-order linear recurrences, *J. Comput. Appl. Math.* (1) 201 (2007), 182–197.
- [6] Lind, D.A.: Problem H-64, *Fibonacci Quart.* 3 (1965), 44.
- [7] Püschel, M., Moura Jose, M.F.: The algebraic approach to the discrete cosine and sine transforms and their fast algorithms, *SIAM J. Comput.* 32 (2003), 1280–1316.
- [8] Rivlin, T.: Chebyshev polynomials: From Approximation Theory to Algebra and Number Theory 2nd ed., New York, Wiley & Sons, 1990.
- [9] Strang, G.: The discrete cosine transform, *SIAM Rev.* 41 (1999), 135–147.
- [10] Webb, W.A., Parberry, E.A.: Divisibility properties of Fibonacci polynomials, *Fibonacci Quart.* 7 (1969), 457–463.
- [11] Zeitzlin, D.: Solution to problem H-64: Fibonacci numbers as a product of cosine terms, *Fibonacci Quart.* 5 (1967), 74–75.

TOBB ECONOMICS AND TECHNOLOGY UNIVERSITY, MATHEMATICS DEPARTMENT,
06560 SOGUTOZU, ANKARA, TURKEY
E-mail address: `ekilic@etu.edu.tr`

DEPARTMENT OF APPLIED MATHEMATICS, NAVAL POSTGRADUATE SCHOOL, MON-
TEREY, CA 93943
E-mail address: `pstanica@nps.edu`