

Final Technical Report

Grant No. N000140710076

Principal Investigator: John W. Frost

Performing Organization:

Draths Corporation
2367 Science Parkway, Ste. 2
Okemos, MI 48864

Grant Title:

Green Synthesis of Phloroglucinol: Exploiting
Pseudomonas fluorescens and Scale-Up

Grant Period:

October 04, 2006 through October 14, 2009

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2009		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Green Synthesis of Phloroglucinol: Exploiting Pseudomonas fluorescens and Scale-Up				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Draths Corporation 2367 Science Parkway, Ste. 2 Okemos, MI 48864				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Through a combination of metabolic engineering and reaction engineering, a new generation of phloroglucinol synthesizing microbe has been developed and evaluated using resin-based extractive fermentation process. Phloroglucinol synthase, phlD, from P. fluorescens PF-5 had been cloned and expressed in E. Coli PG1/pBC2.274. The isolated genes product has been unambiguously identified as phloroglucinol synthase using electrospray ionization tandem mass spectroscopy. The expression of phlD in E. Coli PG1/pBC2.274 under fermentor-control conditions generated inclusion bodies of inactive protein. By attenuating the expression level of phlD in E. coli, a more stable phloroglucinol production E. coli strain, PG1/pKIT10.080, has been developed and capable of producing 25 g/L of phloroglucinol under fermentor-controlled resin-based extractive cultivation conditions. The use of resin-based extractive technique facilitates the removal of toxic phloroglucinol product from the fermentation process as it is being formed allowing for maximum production of phloroglucinol in the E. coli host. In contrast, microbial synthesis of phloroglucinol by a wild-type P. fluorescens species under similar culturing condition produced only 0.1g/L phloroglucinol.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Abstract/Project Summary: Through a combination of metabolic engineering and reaction engineering, a new generation of phloroglucinol synthesizing microbe has been developed and evaluated using resin-based extractive fermentation process. Phloroglucinol synthase, *phlD*, from *P. fluorescens* PF-5 had been cloned and expressed in *E. Coli* PG1/pBC2.274. The isolated gene's product has been unambiguously identified as phloroglucinol synthase using electrospray ionization tandem mass spectroscopy. The expression of *phlD* in *E. Coli* PG1/pBC2.274 under fermentor-control conditions generated inclusion bodies of inactive protein. By attenuating the expression level of *phlD* in *E. coli*, a more stable phloroglucinol production *E. coli* strain, PG1/pKIT10.080, has been developed and capable of producing 25 g/L of phloroglucinol under fermentor-controlled resin-based extractive cultivation conditions. The use of resin-based extractive technique facilitates the removal of toxic phloroglucinol product from the fermentation process as it is being formed allowing for maximum production of phloroglucinol in the *E. coli* host. In contrast, microbial synthesis of phloroglucinol by a wild-type *P. fluorescens* species under similar culturing condition produced only 0.1g/L phloroglucinol.

Scientific Technical Objectives: A new generation of phloroglucinol synthesizing microbe has been evaluated using resin-based extractive fermentation at 1 L scale. Strategies to understand *in vivo* phloroglucinol synthase expression in this genetically engineered *E. coli* were examined. Improving phloroglucinol synthase activity is essential to increase the microbial phloroglucinol synthesis titer and yield. Strategies to identify novel phloroglucinol synthases were therefore explored. It is also believed that expression of phloroglucinol synthase *phlD* gene inside its native *Pseudomonas fluorescens* Pf-5 will likely deliver higher expression level and activity of this gene product. Constructing a phloroglucinol synthesizing *P. fluorescens* was therefore pursued.

Approach: Coupling metabolic engineering and reaction engineering, improved phloroglucinol synthesizing *E. coli* PG1/pKIT10.080 was evaluated under resin-based extractive fermentor-controlled conditions. Heterologous expression of *P. fluorescens* Pf-5 *phlD* gene in *E. coli* presents numerous complications. 2D protein gel electrophoresis and tandem mass spectroscopy were used to examine the relatively short catalytic lifetime and low activity of PhlD. In an attempt to identify better phloroglucinol synthases, the genetic diversity of *phlD* was explored. Bioinformatics analysis of potential *phlD* candidates in various organisms with significant amino acid sequence identity to our currently used Pf-5 *phlD* were examined. In parallel with these efforts, microbial synthesis of phloroglucinol in *P. fluorescens* was examined. Defined minimal salt medium was formulated to enable high density cultivation of *P. fluorescens* Pf1.162/pJA2.232 in the fermentor.

Accomplishments: Under optimized resin-based extractive fermentation, *E. coli* PG1/pKIT10.080 synthesized 25 g/L of phloroglucinol. Coupling 2D protein gel electrophoresis and tandem mass spectroscopy, it was determined that heterologously expressing PhlD-encoded phloroglucinol synthase under fermentor-controlled conditions led to significant formation of inclusion bodies. In attempts to identify phloroglucinol synthase alternatives using bioinformatics approach, five active phloroglucinol synthases were identified. High-cell density cultivation of *P. fluorescens* was achieved in fermentation vessels using minimal salts medium. Phloroglucinol production was observed by culturing *P. fluorescens* Pf1.162/pJA2.232 under defined fermentor-controlled conditions.

Conclusions: The production of phloroglucinol is easily achieved from glucose, and the remaining technical issue is now clear; expression of the phlD enzyme in its properly folded, catalytically active state. Draths is actively pursuing the necessary business issues for access to the Pfenex expression system, which is known to be capable of functional expression of pseudomonad proteins such as phlD.

Significance: Phloroglucinol continues to attract interest at commercial scale and Draths continues to receive inquiries for commercial supply. Production of phloroglucinol from biomass sugars enjoys no viable competition from the petrochemical industry. This commercially useful molecule is expected to be very attractive target for the bio-based chemicals industry.

Publications: None

Patent Information: Provisional Patent for : Phloroglucinal Synthases & Methods of Making and Using the Same by John W. Frost. Filed in United States.

Technology Transfer: None

Awards/Honors: None