

SCIENTIFIC NOTE

ANOPHELES BELENRAE, A POTENTIAL VECTOR OF *PLASMODIUM VIVAX* IN THE REPUBLIC OF KOREA

LEOPOLDO M. RUEDA^{1,4} CONG LI,¹ HEUNG CHUL KIM,² TERRY A. KLEIN,³ DESMOND H. FOLEY¹
AND RICHARD C. WILKERSON¹

ABSTRACT. The malarial parasite, *Plasmodium vivax*, was detected in 4 species of *Anopheles* (Hyrceanus Group) mosquitoes, namely *An. kleini*, *An. pullus*, *An. belenrae*, and *An. sinensis*, from Gyeonggi Province, Republic of Korea (ROK). This study confirmed that *An. belenrae* was infected by *P. vivax*, and implicated this species as a potential vector of *vivax* malaria in the ROK.

KEY WORDS *Anopheles belenrae*, *Plasmodium vivax*, Culicidae, Hyrcanus Group, Korea

Anopheles belenrae Rueda, a member of the *Anopheles* Hyrcanus Group (Culicidae, Diptera), was 1st described in 2005 (Li et al. 2005, Rueda 2005) from specimens collected in the Republic of Korea (ROK). It is presently known to occur in three Asian countries, namely, ROK (Gyeonggi Province; Rueda 2005, Rueda et al. 2006); Democratic People's Republic of Korea (Hwanghaebuk and Pyongyannam Provinces; Rueda and Gao 2008), and China (Liaoning and Shandong Provinces; Rueda et al. 2007).

In the ROK, *An. kleini* Rueda and *An. pullus* Yamada are considered the primary vectors of *Plasmodium vivax* Grassi and Feletti malaria, while *An. sinensis* Wiedemann is a secondary vector (Lee et al. 2007, Klein et al. 2008). Laboratory studies have also shown the potential for *An. lesteri* Baisas and Hu to be an efficient vector (Joshi et al. 2009) and is considered a primary vector (= *An. anthropophagus* Xu and Feng) in parts of China (Rueda et al. 2007). However, since populations of *An. lesteri* are very low, its role in malaria transmission in the ROK is questionable.

We collected about 7,000 adult anopheline mosquitoes by New Jersey light traps (Bioquip, Rancho Dominguez, CA), Mosquito Magnets® (Woodstream Corp., Lititz, PA), and resting catches (mouth aspirator with filter disk; Bio-

quip) at a variety of locations in the ROK in 2008, to better understand malaria transmission in the field. Each New Jersey light trap was suspended from a wooden pole, near barracks and larval habitats (e.g., rice paddy, irrigation ditch, pond, etc.), about 1.5 m above ground level. New Jersey light traps and Mosquito Magnets were set each afternoon between 1700 and 1730 h, and the collections were picked up the following morning between 0800 and 0830 h. Resting adults were caught using aspirators inside the cattle shed, particularly from the walls, roof, and other parts of the building, between 2000 and 2200 h. Collected specimens were removed, taken to the laboratory, sorted, initially identified morphologically (Tanaka et al. 1979, Rueda 2005), and stored in the freezer prior to polymerase chain reaction (PCR) assay. Only the head and thorax of mosquitoes were assayed by PCR for species identification and confirmation (Wilkinson et al. 2003, Li et al. 2005) and by single step and seminested multiplex-PCR to identify *P. vivax* sporozoite infections (Rubio et al. 1999, Hasan et al. 2009). The abdomen was separated from the mosquito body prior to assay to reduce the probability of detecting DNA from oocysts in the hind gut.

Four *P. vivax*-infected adult females of *An. belenrae* (4/87) were collected from Tongilchon and Ilsan, Gyeonggi Province in 2008. One infected female (1/73) was caught in a New Jersey light trap from Tongilchon (37.90517°N, 126.73367°E), Paju, Gyeonggi Province; with collection/serial no. 8.3435, August 6, 2008. Two infected females (2/9) were collected while resting in a cowshed, using an inhaling type aspirator (with filter disk; Bioquip) from Ilsan (37.67135°N, 126.69677°E), Goyang, Gyeonggi Province; with collection/serial no. 8.0758, July 7, 2008 and serial no. 8.3847, August 10, 2008. One infected female (1/5) was collected in a New Jersey light trap from Camp Casey, Dongduchon

¹ Walter Reed Biosystematics Unit, Division of Entomology, Walter Reed Army Institute of Research, 503 Robert Grant Avenue, Silver Spring, MD 20910-7500.

² 5th Medical Detachment, 168th Multifunctional Medical Battalion, 65th Medical Brigade, Unit 15247, APO AP 96205-5247.

³ Force Health Protection and Preventive Medicine, 65th Medical Brigade/USAMEDDAC-Korea, Unit 15281, APO AP 96205-5281.

⁴ Walter Reed Biosystematics Unit, Museum Support Center, Smithsonian Institution, 4210 Silver Hill Road, Suitland, MD 20646-2863.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Anopheles Belenrae, A Potential Vector of Plasmodium Vivax in the Republic of Korea				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Walter Reed Biosystematics Unit, 503 Robert Grant Avenue, Silver Spring, MD, 20910				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The malarial parasite, Plasmodium vivax, was detected in 4 species of Anopheles (Hyrceanus Group) mosquitoes, namely An. kleini, An. pullus, An. belenrae, and An. sinensis, from Gyeonggi Province, Republic of Korea (ROK). This study confirmed that An. belenrae was infected by P. vivax, and implicated this species as a potential vector of vivax malaria in the ROK.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

(37.928075°N, 127.05957°E), Gyeonggi Province, with collection/serial no. 8.2789, July 22, 2010. These data implicate *An. belenrae* as a potential vector of *P. vivax*. In addition, Chang et al. (unpublished data) noted that two specimens of *An. belenrae* collected from Paju on May 27, 2009 and Goseong County, ROK on June 22, 2009 were tested positive by PCR for *P. vivax*.

To incriminate a malaria vector in nature, it is necessary to demonstrate: 1) an association between the vector and human malaria cases; 2) evidence of direct human–vector contact; and 3) evidence that the vector harbors sporozoites in the salivary glands (Beier, 2002). In addition to evidence of sporozoites in *An. belenrae*, we should mention that in 2006–2007 human malaria cases had rapidly increased in the area where we collected malaria positive *An. belenrae* (Goyang) (Jun et al. 2009), but the role, if any, of this species in malaria transmission at this location is unknown.

Other *Anopheles* species that were also found positive for *P. vivax* from our collections in Gyeonggi Province, ROK, were *An. kleini*, *An. pullus*, and *An. sinensis*. Six infected females of *An. kleini* (6/199) were collected from the following sites: Warrior Base (37.92444°N, 126.73944°E), *n* = 1 infected female (F) (1/7), August 8, 2008, collection/serial no. 8.3923; Tongilchon, Paju (37.90517°N, 126.738411°E), *n* = 1 F (1/71), July 2, 2008, collection/serial nos. 8.0627, 8.0458; *n* = 2 F (2/119), July 29, 2009, collection/serial nos. 8.2817, 8.2821; and ROK Army Base (37.918832°N, 126.738411°E), *n* = 2 F (2/2), June 13 and July 14, 2008, collection/serial nos. 8.2294, 8.2476. Three infected females of *An. pullus* (3/42) were collected from Tongilchon, Paju, *n* = 3 F, July 2, 2008, collection/serial no. 8.0470, 8.0494, 0508. Fifteen infected females of *An. sinensis* (15/273), were found in Tongilchon, Paju: *n* = 5 F (5/63), July 2, 2008, collection/serial nos. 8.0610, 8.0628, 8.0520, 8.0548, 8.0597; *n* = 1 F (1/60), July 11, 2008, collection/serial no. 8.0966; *n* = 3 F (3/37), July 23, 2008, collection/serial nos. 8.0206, 8.0221, 8.0238; *n* = 5 F (5/107), July 29, 2009; collection/serial nos. 8.2714, 8.2756, 8.2785, 8.2826, 8.2832; and ROK Army Base, *n* = 1 F (1/6), June 13, 2008, collection/serial no. 8.2476.

Furthermore, we are continuing the molecular analysis of several thousand *Anopheles* females from different locations in the ROK, particularly to identify the mosquito species and their associated *Plasmodium* parasites.

Thanks to personnel of the 5th Medical Detachment, and staff of 65th Medical Brigade, U.S. Army, ROK, for field collections of mosquito specimens; and J. Pecor and Walter Reed Biosystematics Unit staff for curatorial help. We are grateful to Y. M. Huang, C. R. Summers, and B. P. Rueda for helpful reviews of

the manuscript. Funding for this work was provided by the Global Emerging Infections Surveillance and Response Systems of the Armed Forces Health Surveillance Center, Silver Spring, MD. This research was performed under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Department of the Army or the Department of Defense.

REFERENCES CITED

- Beier JC. 2002. Vector incrimination and entomological inoculation rates. *Methods Mol Med* 72:3–11.
- Hasan AU, Suguri S, Sattabongkot J, Fujimoto C, Amakawa M, Harada M, Ohmae H. 2009. Implementation of a novel PCR based method for detecting malaria parasites from naturally infected mosquitoes in Papua New Guinea. *Malaria J* 8(182):1–11.
- Joshi D, Choochote W, Park MH, Kim JY, Kim TS, Suwonkerd W, Min GS. 2009. The susceptibility of *Anopheles lesteri* to infection with Korean strain of *Plasmodium vivax*. *Malaria J* 8:42.
- Jun G, Yeom JS, Hong JY, Shin EH, Chang KS, Yu JR, Oh S, Chung H, Park JW. 2009. Resurgence of *Plasmodium vivax* malaria in the Republic of Korea during 2006–2007. *Am J Trop Med Hyg* 81:605–610.
- Klein TA, Kim HC, Lee WJ, Rueda LM, Sattabongkot J, Moore RG, Chong ST, Sames W, Pike JG, Wilkerson RC. 2008. Reemergence, persistence and surveillance of vivax malaria and its vectors in the Republic of Korea. In: Robinson WK, Bajoni D, eds. Proceedings of the 6th International Conference on Urban Pests. 2008 July 13–16; Budapest, Hungary: Oak Press, Kft. p 325–331.
- Lee WJ, Klein TA, Kim HC, Choi YM, Yoon SH, Chang KS, Chong ST, Lee IY, Jones JW, Jacobs JS, Sattabongkot J, Park JS. 2007. *Anopheles kleini*, *An. pullus*, and *An. sinensis*: Potential vectors of *Plasmodium vivax* in the Republic of Korea. *J Med Entomol* 44:1086–1090.
- Li C, Lee JS, Groebner JL, Kim HC, Klein TA, O'Guinn ML, Wilkerson RC. 2005. A newly recognized species in the *Anopheles* Hyrcanus Group and molecular identification of related species from the Republic of South Korea (Diptera: Culicidae). *Zootaxa* 939:1–8.
- Rubio JM, Benito A, Roche J, Berzosa PJ, Garcia ML, Mico M, Edu M, Alvar J. 1999. Semi-nested, multiplex polymerase chain reaction for detection of human malaria parasites and evidence of *Plasmodium vivax* infection in Equatorial Guinea. *Am J Trop Med Hyg* 60:183–187.
- Rueda LM. 2005. Two new species of *Anopheles* (*Anopheles*) Hyrcanus Group (Diptera: Culicidae) from the Republic of South Korea. *Zootaxa* 941: 1–26.
- Rueda LM, Gao Q. 2008. New records of *Anopheles belenrae* Rueda (Diptera: Culicidae) in North Korea. *Proc Entomol Soc Wash* 110:523–524.

- Rueda LM, Kim HC, Klein TA, Pecor JE, Li C, Sithiprasasna R, Debboun M, Wilkerson RC. 2006. Distribution and larval habitat characteristics of *Anopheles* Hyrcanus Group and related mosquito species (Diptera: Culicidae) in South Korea. *J Vector Ecol* 31:199–206.
- Rueda LM, Zhao T, Ma YJ, Gao Q, Guoding Z, Khuntirat B, Sattabongkot J, Wilkerson RC. 2007. Updated distribution records of the *Anopheles* (*Anopheles*) hyrcanus species-group (Diptera: Culicidae) in China. *Zootaxa* 1407:43–55.
- Tanaka K, Mizusawa K, Saugstad ES. 1979. A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Contrib Am Entomol Inst.* 16:1–987.
- Wilkerson RC, Li C, Rueda LM, Kim HC, Klein TA, Song GH, Strickman D. 2003. Molecular confirmation of *Anopheles* (*Anopheles*) *lesteri* from the Republic of South Korea and its genetic identity with *An. (Ano.) anthropophagus* from China (Diptera: Culicidae). *Zootaxa* 378:1–14.