

AD _____

Award Number: W81XWH-10-1-0666

TITLE: BMP7 Induces Dormancy of Prostatic Tumor Stem Cell in Bone

PRINCIPAL INVESTIGATOR: Aya Kobayashi, Ph.D.

CONTRACTING ORGANIZATION: Southern Illinois University, School of Medicine
Springfield, IL 62794

REPORT DATE: October 2011

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE October 2011		2. REPORT TYPE Annual		3. DATES COVERED 30 September 2010 – 29 September 2011	
4. TITLE AND SUBTITLE BMP7 Induces Dormancy of Prostatic Tumor Stem Cell in Bone				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-10-1-0666	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Aya Kobayashi, Ph.D. E-Mail: akobayashi@siuimed.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Southern Illinois University, School of Medicine Springfield, IL 62794				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Bone is the most common metastatic site for prostate cancer. The growth of the tumor cells in the bone is generally slow and they often become dormant until an appropriate microenvironment is established for their re-growth. The recent stem cell theory predicts that the metastatic cells are a small population of stem-like cells in the primary tumor. However, the precise mechanism of dormancy is virtually unknown, and identifying the responsible factors and understanding their underlining mechanism are crucial for developing a novel therapeutic approach. Our preliminary data indicate that (i) bone morphogenetic protein 7 (BMP7) which is secreted from bone marrow stromal cells is able to induce senescence to prostate tumor cell and (ii) this induction is mediated by activation of the tumor metastasis suppressor gene, N-myc downstream regulated gene 1 (NDRG1). These results strongly suggest that the BMP7-NDRG1 axis plays a critical role in dormancy of prostate tumor cells in the bone. The overall goal of this proposal is to elucidate the mechanism of BMP7-induced dormancy in tumor stem cells and explore a possibility of using BMP7 as an anti-metastatic drug for prostate cancer.					
15. SUBJECT TERMS Prostate cancer, dormancy, BMP7, tumor stem cell, bone metastasis					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	9	19b. TELEPHONE NUMBER (include area code)

Table of Contents

Introduction..... 4

Body..... 4

Key Research Accomplishments..... 8

Reportable Outcomes..... 8

Conclusions..... 9

References..... 9

INTRODUCTION

Bone is the most common metastatic site for prostate cancer which affects approximately 70% of patients with advanced disease (1). Despite this clinical importance, the exact molecular mechanism of the bone-specific metastasis has not been clearly defined. The growth of the tumor cells in the bone is generally slow and they often become dormant until an appropriate microenvironment is established for their re-growth. According to the recent cancer stem cell theory, which still remains a hypothesis, recurrent tumor must arise from a dormant tumor stem cell (2). However, the precise mechanism of dormancy is virtually unknown, and identifying the responsible factors and understanding the underlying mechanism are crucial for developing a novel therapeutic approach. Our recent preliminary data suggest that BMP7 (bone morphogenetic protein 7) which is secreted from bone stromal cells is able to induce senescence to prostate tumor cells and that this induction is mediated by activation of the tumor metastasis suppressor gene, NDRG1 (N-myc downstream regulated gene 1). Therefore, we hypothesize that a prostatic tumor stem cell becomes dormant in the bone through the BMP7-mediated activation of p38 and NDRG1.

BODY

Task 1. To clarify the signaling pathway of BMP7-induced senescence in tumor stem cell.

- (a) Examine whether BMP7 blocks Wnt pathway by activating p38-NDRG1 axis.
- (b) Test whether BMP7 blocks Wnt pathway by increasing the binding of NDRG1 to LRP6.
- (c) Examine whether the BMP7-induced senescence is dependent on Wnt signaling.
- (d) Examine whether BMP7-induced MET (mesenchymal-epithelial-transition) is dependent on Wnt signal and whether it is prerequisite for senescence.

To accomplish Task 1 (a)-(c), we first examined the effect of BMP7 and Wnt on the growth of prostate cancer cells. As shown in Figure 1, BMP7 significantly suppressed the growth, while Wnt3a significantly promoted the prostate cancer cell growth. Furthermore, up-regulation of NDRG1 counteracted the effect of Wnt3a on the cancer cell growth, suggesting that the balance of BMP and Wnt signalings may determine whether the tumor cells stay in growth arrest or proliferate. Next, we isolated cancer stem cells (CSCs, CD24⁻/CD44⁺/CD133⁺) from PC3mm cells that were labeled with the luciferase gene. Our result of serial dilution assay for tumor initiating ability in animals indicates that the CSC population was significantly more tumorigenic than non-stem cells (CD24⁺/CD44⁻/CD133⁻) (Figure 2A). We also found that BMP7 was able to activate p38, NDRG1 and p21 (Figure 2B-2D) in CSCs, and that the induction of NDRG1 and p21 was mediated by p38 (Figure 2C and 2D). In addition, BMP7 significantly inhibited the sphere forming ability of CSCs; however, these cells regained the growth ability after withdrawal of BMP7 from the medium (Figure 2E). Furthermore, BMP7 was also able to induce senescence in CSCs (Figure 2F; +/+) with concomitant activation of

NDRG1 and p21 (Figure 2G; +/+ and 2H; +/+). Of note, the BMP7-mediated induction of senescence as well as the activation of NDRG1 and p21 was reversed after withdrawal of BMP7 (Figure 2F; +/-, 2G; +/-, and 2H; +/-). Similarly, up-regulation of NDRG1 in CSCs by using PC3mm cell line which had tetracycline-inducible




Figure 1. NDRG1 counteracts the effect of Wnt on cancer cell growth. PC3mm/Tet-NDRG1 cells were cultured for 96 h with or without BMP7, Wnt3a or tetracycline, and the cell viability was measured by MTS assay. **, P<0.01, ***, P<0.001 versus -, ##, P<0.01 versus Wnt. n = 3.

NDRG1 significantly suppressed the sphere formation and induced senescence, while de-induction of NDRG1 by withdrawal of tetracycline reversed this effect (Figure 2I and 2J). These results suggest that BMP7 was able to induce reversible senescence in CSCs through activation of p38, p21 and NDRG1.




Figure 2. BMP7 induces reversible senescence in cancer stem-like cells (CSCs) through activation of p38, p21 and NDRG1. (A) CSCs isolated from PC3mm were injected subcutaneously into nude mice, and the growth of tumor was monitored by bioluminescence imaging (BLI). (B) The CSCs were treated with the HS5 (bone stromal cells)-CM or BMP7, and the expression of p-p38, NDRG1 and β-Tubulin was examined by Western blot. (C and D) The CSCs were treated with or without BMP7 and/or SB203580 (SB), and the expression of NDRG1 (C) and p21 (D) were examined by qRT-PCR. $n = 3$, *, $P < 0.05$, ***, $P < 0.001$. (E) Effect of BMP7 on the sphere forming ability of CSCs was measured. (-/-/-): no treatment throughout. (-/+/-): no treatment for 5 days, treated with BMP7 for 4 days, and no treatment thereafter. (+/+/-): treatment for 9 days, and no treatment thereafter ($n = 5$, *, $P < 0.001$ versus (-/-/-), #, $P < 0.001$ versus (-/-/-)). (F) CSCs were cultured in the presence (+/+), absence (-/-) or withdrawal after treatment (+/-) of BMP7, and SA-β-gal staining was performed. $n = 6$, ***, $P < 0.001$. Scale bar represents 10 μm. (G and H) The CSCs were treated with BMP7, and the expression of NDRG1 (G) and p21 (H) was measured by qRT-PCR. (-/-): no treatment control, (+/+): continuous treatment with BMP7 for 96 h, (+/-): 48 h-treatment followed by 48 h-withdrawal of BMP7. $n = 3$, *, $P < 0.05$, **, $P < 0.01$. (I) A similar experiment was performed as (E) for CSCs from PC3mm/Tet-NDRG1 with (+) or without (-) induction of NDRG1, followed by assaying sphere formation. (-/-/-): no induction throughout. (-/+/-): no induction for 5 days, induction for 3 days, and no induction thereafter. (+/+/-): induction for 8 days, and no treatment thereafter ($n = 5$, *, $P < 0.001$ versus (-/-/-), #, $P < 0.001$ versus (-/-/-)). (J) A similar experiment was performed as (F) for CSCs from PC3mm/Tet-NDRG1. They were cultured with (+/+) or without (-/-) induction or withdrawal after induction of NDRG1 (+/-), and SA-β-gal staining was performed ($n = 8$, ***, $P < 0.001$). The expression of NDRG1 and β-Tubulin was examined by Western Blot (upper panel).

To accomplish Task 1 (d), we examined whether BMP7 activates MET program in prostate cancer cells. We found that BMP7 significantly reduced vimentin (mesenchymal marker) expression while it significantly augmented E-cadherin (epithelial marker) expression (Figure 3A, B). Moreover, up-regulation of NDRG1 induced similar MET effect (Figure 3C, D), suggesting that BMP7-NDRG1 axis induced MET in prostate cancer cells.




Figure 3. BMP7 induces MET. (A and B) PC3mm cells were cultured with BMP7 or control vehicle for 48 h and the expression of vimentin (A) and E-cadherin (B) were examined by qRT-PCR. (C and D) PC3mm/Tet-NDRG1 cells were cultured with or without tetracycline for 48 h and the expression of vimentin (C) and E-cadherin (D) were examined by qRT-PCR.

Therefore, we now have identified that BMP7-p38-NDRG1 axis plays critical roles in inducing senescence and MET in CSCs. To accomplish Task 1, we will examine the effect of Wnt pathway on BMP7-induced senescence, MET and its signaling pathway. Therefore, Task 1 is still ongoing.

Task 2. To examine the effect of BMP7 on tumor stem cell *in vivo*

- Co-inject tumor stem cell and bone marrow stromal cell into nude mouse using calcium-phosphate scaffold and examine the effect of BMP7 on the tumor growth.
- Examine the effect of BMP7 on dormancy by injecting tumor stem cell via intracardiac followed by directly injecting BMP7 intravenously.

To accomplish Task 2, we isolated tumor stem cells from PC3mm labeled with luciferase gene and these cells were seeded onto CaP scaffolds with HS5 (bone stromal cells), and then these constructs were implanted into the upper back of nude mice. Recombinant human BMP7 was peritumorally injected daily after implantation. Tumor growth was monitored weekly by bioluminescent imaging. We found that BMP7 treatment significantly suppressed the tumor growth (Figure 4A and B). To further examine the effect of BMP7 on the tumor growth in bone, CSCs were injected into tibiae of mice with HS5 cells that had either scrambled shRNA (control) or shRNA for BMP7 (shBMP7). We found that the control HS5 was able to suppress tumor growth and activated p38 and NDRG1; however, HS5 with shBMP7 significantly abrogated such an effect on CSCs and failed to suppress tumor-

induced osteolysis in the tibiae (Figure 4C). These results suggest that BMP7 secreted from the bone stromal cells is indeed capable of inducing growth arrest of CSCs through activation of p38 and NDRG1 in the bone environment. We also assessed the effect of NDRG1 in the suppression of tumor growth by co-injecting HS5 and CSCs in which NDRG1 expression was knocked down by shRNA (shNDRG1), into mice tibiae. We found that knockdown of NDRG1 in CSCs enabled these cells to grow significantly greater than that of control CSCs even when they were co-injected with HS5 (Figure 4D). These results indicate that the BMP7-NDRG1 axis plays a critical role in the growth suppression of metastasized prostate tumor cells in bone, and also strongly suggest the potential therapeutic utility of BMP7 for metastatic disease.




Figure 4. BMP7 suppresses tumor growth of CSCs *in vivo*. (A and B) CSCs from PC3mm were seeded onto CaP scaffolds with HS5 and these constructs were implanted into nude mice, and then BMP7 was administrated daily. BLI of representative mice in each group six weeks after implantation (A). Normalized BLI signals after six weeks (B). -: control vehicle, $n=6$, BMP7: 100 g/kg BMP7 treatment, $n=10$. *; $P=0.0225$ by Mann-Whitney test. (C) CSCs isolated from PC3mm were co-injected into tibiae of nude mice with control (scramble, $n = 11$) or BMP7-knocked down (shBMP7, #1: $n = 14$, #2: $n = 10$) HS5 cells. Left panel: bioluminescence imaging (BLI) and x-ray radiography of bone lesions from representative mice in each group four weeks after inoculation. Osteolytic lesions are indicated by arrow heads. Center panel: Normalized BLI signals after four weeks. *; $P<0.05$ by Mann-Whitney test. Right panel: Representative images of immunohistochemical staining of tumors in the tibiae for p-p38 and NDRG1. Scale bar represents 50 μ m. (D) CSCs isolated from PC3mm/shNDRG1 ($n = 12$) or PC3mm/scramble ($n = 12$) were co-injected with HS5 into tibiae of nude mice. Left panel: BLI and x-ray radiography of representative mice in each group four weeks after inoculation. Osteolytic lesions are indicated by arrow heads. Center panel: BLI after four weeks. **; $P<0.01$, by Mann-Whitney test. Right panel: H&E staining of the tibiae from representative mice four weeks after inoculation. Asterisk indicates tumor. Scale bar represents 200 μ m.

Therefore, Task 2 (a) is accomplished and Task2 (b) is in progress.

Task 3. To examine whether serum level of BMP7 and Wnt ligand correlates with bone metastasis in patients with prostate cancer

(a) Examine the expression level of BMP7 and Wnt ligand in serum of patients with normal, primary and metastatic prostate tumor.

We are still in the process of collecting sera of prostate cancer patients. We expect that we will obtain the samples and conduct the experiment of Task 3 in the next fiscal year.

KEY RESEARCH ACCOMPLISHMENTS

1. We found that NDRG1 counteracts the growth promoting effect on prostate cancer cells by Wnt.
2. We found that BMP7 induces senescence by activating p38, p21 and NDRG1 in CSCs.
3. We found that BMP7-induced senescence is reversible.
4. We found that BMP7 induces MET in prostate cancer cells.
5. We found that BMP7 significantly suppresses the growth of CSCs on CaP scaffold model.
6. We found that BMP7 secreted from bone stromal cells inhibits the growth of CSCs through activation of p38 and NDRG1 in bone.

REPORTABLE OUTCOMES

Peer reviewed publications

Aya Kobayashi, Hiroshi Okuda, Fei Xing, Puspa R. Pandey, Misako Watabe, Shigeru Hirota, Sudha K. Pai, Wen Liu, Koji Fukuda, Christopher Chambers, Andrew Wilber and Kounosuke Watabe “Bone morphogenetic protein 7 in dormancy and metastasis of prostate cancer stem-like cells in bone” In preparation

Abstract/presentation

1. Aya Kobayashi, Hiroshi Okuda, Puspa R. Pandey, Misako Watabe, Sudha K. Pai, Shigeru Hirota, Fei Xing, Wen Liu, Bo Xia, Kounosuke Watabe
Poster Presentation: “BMP7 regulates dormancy and recurrence of prostate cancer stem cell in bone”
Joint MRS-AACR Conference: Metastasis and the Tumor Microenvironment, 2010, Philadelphia, PA
2. Aya Kobayashi, Hiroshi Okuda, Puspa Pandey, Misako Watabe, Sudha K. Pai, Fei Xing, Shigeru Hirota, Wen Liu, and Kounosuke Watabe
Poster Presentation: “Dormancy and recurrence of prostate cancer stem cell are regulated by bone morphogenetic protein 7 in bone”
Department of Defense (DOD) Prostate Cancer Research Program (PCRP) Innovative Minds in Prostate Cancer Today (IMPACT) Conference, 2011, Orlando, FL

Employment

Aya Kobayashi, Ph.D. (postdoctoral fellow) has been partly supported by the current grant..

CONCLUSIONS

We have shown that BMP7 secreted from bone stromal cells induces senescence through activation of p38, p21 and NDRG1. We have also shown that this senescence is reversible and the CSCs regain the ability to proliferate upon withdrawal of BMP7. Therefore, we are now in a strong position to pursue the rest of tasks. In the next fiscal year, we will address two important points. First, we will clarify whether BMP7 blocks the Wnt pathway and whether the Wnt pathway interferes with the BMP7-induced senescence and MET. Secondly, we will compare the expression level of BMP7 and Wnt ligand in sera of prostate cancer patients with or without metastatic disease. The results of this research will provide us with strong rationale for using BMP7 as an anti-recurrence agent to treat prostate cancer patients with metastatic disease.

So what?

Despite the significant improvement in recent therapeutic technologies, metastatic disease is still the ultimate cause of death in prostate cancer patients (3). Therefore, understanding the underlying mechanisms of dormancy and recurrence in bone metastasis is of paramount interest for developing an effective approach to treat and prevent recurrent prostate cancer. We believe that elucidation of such mechanisms will provide a paradigm shift in this research field and eventually lead to the development of a novel therapeutic approach to this devastating disease.

REFERENCES

1. Cancer Facts & Figures. *American Cancer Society*, 2009.
2. Polyak, K. and Weinberg, R. A. Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. *Nat Rev Cancer.*, 9: 265-273, 2009.
3. Suva LJ, Washam C, Nicholas RW and Griffin RJ. Bone metastasis: mechanisms and therapeutic opportunities. *Nat Rev Endocrinol.*, 7: 208-218, 2011