

AD _____

Award Number: W81XWH-09-1-0450

TITLE: Neuroendocrine Differentiation in Prostate Cancer: Role of Bone Morphogenetic Protein-6 and Macrophages

PRINCIPAL INVESTIGATOR: Dr. Isaac Kim

CONTRACTING ORGANIZATION: UMDNJ-Robert Wood Johnson Medical School
Piscataway, NJ 08854

REPORT DATE: July 2011

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-07-2011		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 15 JUN 2010 - 14 JUN 2011	
4. TITLE AND SUBTITLE Neuroendocrine Differentiation in Prostate Cancer: Role of Bone Morphogenetic Protein-6 and Macrophages				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-09-1-0450	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Dr. Isaac Kim E-Mail:kimiy@umdnj.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) UMDNJ-Robert Wood Johnson Medical School Piscataway, NJ 08854				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT In the grant proposal we have hypothesized that tumor-derived bone morphogenetic protein-6 (BMP-6) induces tumor associated macrophages (TAMs) to express interleukin-6 (IL-6) via a crosstalk between the Smad-dependent and the p38 pathway; IL-6 in turn drives neuroendocrine (NE) differentiation of prostate cancer cells. To test this proposal, three specific aims were proposed: 1) To investigate the mechanism of NE differentiation induced by BMP-6 in vivo. 2) To investigate the mechanism of IL-6 induction by BMP-6 in macrophages. 3)To study the efficacy of dorsomorphin, a small molecule inhibitor of BMP signaling, on NE differentiation of prostate cancer in vivo. To date, aims 1 and 2 have been completed while aim 3 is currently progressing. When BMP-6 overexpressing prostate cancer cell line Tramp-BMP6 was injected subcutaneously into IL-6 knockout (KO) and conditional macrophage-null mice, neuroendocrine differentiation was no longer observed. Mechanistically, series of studies including shRNA knockdowns and immunoprecipitation assays have confirmed that Smad5 and GAT4 interact to induce IL-6 expression in macrophages. Currently, these finding are being confirmed in vivo using a small molecule inhibitor of BMP-6, dorsomorphine.					
15. SUBJECT TERMS Prostate cancer, bone morphogenetic protein-6, macrophages, interleukin-6					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			USAMRMC
			UU	10	19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	2
Body.....	2
Key Research Accomplishments.....	3
Reportable Outcomes.....	3
Conclusion.....	4
References.....	5
Appendices.....	6

INTRODUCTION

Prostate cancer is the most commonly diagnosed and the second leading cause of cancer deaths in the US (Jemal et al., 2009). Although surgery and radiation are quite effective for an organ-confined disease, the outlook for patients with a metastatic prostate cancer is still bleak due to the inevitable emergence of hormone refractory prostate cancer (HRPC). Since neuroendocrine (NE) cells are androgen receptor negative and frequently found in HRPC (Abrahamsson et al., 1998), understanding the biology of NE cells in prostate cancer may uncover new targets of intervention in patients with HRPC. In this regard, we have observed that the NE cells in human prostate cancer tissues were frequently positive for BMP-6 and co-localized with tumor associated macrophages (TAMs). In tissue culture, co-culture of macrophages and prostate cancer cells led to the expression of NE markers by prostate cancer cells in the presence of BMP-6. A survey of potential macrophage-derived mediators of NE differentiation demonstrated a dramatic induction of interleukin-6 (IL-6) by BMP-6. In investigating the mechanism of IL-6 induction by BMP-6 in macrophages, we have unexpectedly observed that both the canonical BMP-signaling pathway mediated by Smads and the non-canonical Smad-independent pathway mediated by p38 are required. Based on these results, the present grant proposal was funded to further dissect the mechanism as well as the biology of the tumor-host interaction loop composed of BMP-6, TAMs, and IL-6 in the context of NE cells in prostate cancer. This is the progress report for year 2 of the grant.

BODY

Based on the preliminary data, we have hypothesized that **tumor-derived BMP-6 induces TAMs to express IL-6 via a crosstalk between the Smad-dependent and the p38 pathway; IL-6 in turn drives NE differentiation of prostate cancer cells.** To test this hypothesis, 3 specific aims were proposed: 1) To investigate the mechanism of NE differentiation induced by BMP-6 in vivo; 2) To investigate the mechanism of IL-6 induction by BMP-6 in macrophages; 3) To study the efficacy of dorsomorphin, a small molecule inhibitor of BMP signaling, on NE

differentiation of prostate cancer in vivo. Aim1 and parts of aim 2 were completed in year 1 and the past year was spent on completing aim 2 and part of aim 3.

Previously, we have demonstrated that both Smad-dependent and non-Smad pathways are required for the induction of IL-6 by BMP-6 in macrophages. Specifically, a physical interaction between Smad1 and GATA4 was required for BMP's activation of IL-6 promoter in macrophages. During the past year, we have observed that within the Smad-dependent pathway Smad1 alone is not sufficient for the activation of IL-6 promoter by BMP-6; in addition to Smad1, Smad4 was required (Fig 1). More importantly, when Smad4 was knocked down with shRNA, Smad1 translocated to the nucleus (Fig 2). These observations suggest that Smad4 is also required for the transcriptional activation of IL-6 promoter by BMP-6. Concerning the role of Smad-independent pathway and GATA4, we have confirmed that the expression of dominant negative GATA4 blocks the induction of IL-6 expression by BMP-6 (Fig 3).

As for the cross-talk between Smad1 and GATA4, immunoprecipitation previously confirmed a physical interaction. To further support this observation, we have attempted EMSA (electrophoretic mobility shift assay) as proposed in the task 2c of aim 2. But using the proposed antibodies against Smad1 and GATA4, we were not successful in demonstrating the supershifting of IL-6 promoter upon BMP-6 activation. Based on the results of the Smad4 overexpression study, we now propose that Smad4 is another critical player in BMP-6 induced activation of IL-6 promoter. Indeed, it is possible that Smad4 may be the critical link that connects Smad1/GATA4 to IL-6 promoter. We have recently initiated another EMSA experiment to test this hypothesis.

Simultaneously, we have begun investigating the effect of dorsomorphin, a small molecule inhibitor of BMP-6, on neuroendocrine differentiation of prostate cancer. To this end, a murine prostate cancer cell line with tetracycline-inducible expression of BMP-6 was used. This cell line, designated Tramp C2-BMP6, was injected into 40 C57BL/6 mice. The mice were divided into 4 groups as follows:

Cell Line	Tramp C2-BMP6 (n=40)			
	No (n=20)		Yes (n=20)	
Doxycycline				
Dorsomorphin	No (n=10)	Yes (n=10)	No (n=10)	Yes (n=10)

The tissues will be harvested and analyzed for NE differentiation in 2-4 weeks.

KEY RESEARCH ACCOMPLISHMENTS

1. Completion of task 2b and progress on task 2c.
2. Initiation of animal studies examining dorsomorphine in NE differentiation of prostate cancer cells.

REPORTABLE OUTCOMES

1. Published 1 manuscript.

Geun Taek Lee, Seok Joo Kwon, Jae-Ho Lee, Seong Soo Jeon, Kee Taek Jang, Han Yong Choi, Hyun Moo Lee, Wun Jae Kim, Seong Jin Kim, and Isaac Yi Kim. Induction of IL-6 expression by BMP-6 in macrophages requires both Smad and p38 signaling pathways. Journal of Biological Chemistry 285:39401-8, 2010.

CONCLUSION

During the second year of funding, we have demonstrated that BMP-6 induces IL-6 expression in macrophages via Smad1/Smad4/GATA4 interaction. The third year of funding will focus on completing the preclinical studies using a small molecule inhibitor of BMP signaling to determine whether neuroendocrine differentiation of prostate cancer cells is reversible.

REFERENCES

Abrahamsson, P.A., Cockett, A.T., and di Sant'Agnese, P.A. (1998). Prognostic significance of neuroendocrine differentiation in clinically localized prostatic carcinoma. Prostate Suppl 8, 37-42.

Jemal, A., Siegel, R., Ward, E., Hao, Y., Xu, J., and Thun, M.J. (2009). Cancer statistics, 2009. CA Cancer J Clin 59, 225-249.

Appendix




Figure 1. Smad1 and 4 were overexpressed in RAW 264.7 cells. When these cells were treated with BMP-6 100 ng/ml, only the simultaneous expression of both Smad1 and 4 increased BMP-6-induced IL-6 promoter activity as indicated by luciferase activity.




Figure 2. To determine the effect of Smad4 on nuclear translocation of Smad1, shRNA was used to knock-down Smad4. The results demonstrated that the knock-down of Smad4 had no significant effect on nuclear translocation of Smad1.




Figure 3. RAW 264.7 was transfected with dominant negative GATA4 (GATA4DN) and the effect on IL-6 expression was measured using RT-PCR. Following transfection with GATA4DN, baseline induction of IL-6 was again observed, suggesting that the transfection process itself can influence IL-6 expression. When treated with BMP-6, induction of IL-6 was no longer observed in cells expressing GATA4DN.