

Early Student Support for Shipboard LADCP/ χ pod Profiling of Internal Wave Structure and Dissipation in the Luzon Strait

Jonathan D. Nash
College of Oceanic & Atmospheric Sciences
Oregon State University
Corvallis OR 97331-5503
ph: (541) 737-4573 fx: (541) 737-2064
email: nash@coas.oregonstate.edu

Award #: N00014-11-1-0367
<http://mixing.coas.oregonstate.edu/>

LONG-TERM GOALS

We seek a more complete and fundamental understanding of the hierarchy of processes which transfer energy and momentum from large scales, feed the internal wavefield, and ultimately dissipate through turbulence. This cascade impacts the acoustic, optical, and biogeochemical properties of the water column, and feeds back to alter the larger scale circulation. Studies within the **Ocean Mixing Group** at OSU emphasize observations, innovative sensor / instrumentation development and integration, and process-oriented internal wave and turbulence modeling for interpretation.

OBJECTIVES

Luzon Strait represents a major source of internal tides and NLIWs in the SCS. However, unlike other regions of strong internal wave generation (i.e., Hawaii), Luzon Strait is believed to be highly dissipative. We seek to understand the character of this enhanced nonlinearity and turbulence, and how it affects internal wave generation and transmission. Specifically, we intend to:

- identify hotspots of generation and dissipation,
- quantify the structure and variability of wave energy, its flux and dissipation at the generation site.
- link the broader spatial structure, temporal content, and energetics of the internal wave field to the topography, forcing, and mesoscale influences (i.e., Kuroshio).

APPROACH

Much of the turbulent dissipation in Luzon Strait was anticipated to be deep, outside the range of tethered microstructure profilers, and evolving too rapidly for autonomous profilers. We have used a 2-fold approach to quantify this deep turbulence:

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Early Student Support for Shipboard LADCP/ pod Profiling of Internal Wave Structure and Dissipation in the Luzon Strait				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University, College of Oceanic & Atmospheric Sciences, Corvallis, OR, 97331-5503				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

1. rapid profiles at abyssal depths are obtained using standard shipboard CTD, augmented with ADCPs, turbulence sensors, and a motion package. These allow us to systematically obtain 36-h yo-yo timeseries, from which energy generation, energy fluxes, and energy dissipation can be directly measured.
2. Five moorings were successfully deployed and recovered in 2 of the most energetic regions of the strait. Recovery cruises were mid Aug and early Sept, 2011, so these data are just starting to be analyzed.




Figure 1: An example 1.5-day record of T (top) and velocity (bottom; with T contoured) from the stablemoor mooring. Time is in yeardays (2011); depth in m. Peak flows and displacements exceed 1 m/s and 300 m.

WORK COMPLETED

This award has so far provided 4 months of support for graduate student Byungho Lim. Byungho helped with instrument preparations for this summer's fieldwork and participated in two of the IWISE cruises this past summer. He is currently continuing coursework to fulfill requirements of the OSU Physical Oceanography program.

Byungho will be involved in the analysis of moored and shipboard data collected this summer. Examples of these data (from instruments recovered in Aug and Sept 2011) are shown in figures 1 and 2. These focus on 2 aspects of the flow: (1) the radiated energy and local dissipation at a site of strong wave generation (the Stablemoor mooring; figure 1), and (2) the structure of wave and dissipation events on the western ridge near station N2. At that site, 5 moorings were deployed at 1-km spacing to capture the evolution of billowing as the waves break at the ridge crest (figure 2).




Figure 2: The figure at right shows a 16 h segment of temperature data from the 4 T-chains deployed over the ridge near N2. These moorings were spaced 1 km apart and capture displacements and turbulence in x,z,t space. The depictions at right rival numerical simulations in resolution but capture the represent the true reality and breadth of scales of the real ocean. These figures show the evolution and spatial structure of a 300-m tall wave which grows to 500 m and breaks. Short-wavelength features are coherent across the mooring array and can be tracked (grey arrow). 100-500 m tall turbulent overturns are also visible, particularly at mooring T3 at 0900.

This example represents one 16-h time window within the 3-month deployment. Our objectives are to analyze these records in detail in order to quantify both internal wave and turbulent components. We will combine data from LADCP/ χ pod, MP- χ Pods, the moorings shown here, the stablemoor (at A1) and other ancillary moored and model data in these analyses. Ultimately, we will

1. characterize the spatial and temporal variability of high-dissipation events
2. determine the physics of high-wavenumber generation from the surface and internal tides, and the subsequent breakdown into turbulence
3. assess how these dynamics are related to those which have been numerically modeled, with a goal of understanding whether these can be parameterized.

RELATED PROJECTS

Profiling and moored operations are being coordinated with M Alford (UW); analysis of turbulence data are being conducted in conjunction with J MacKinnon (UCSD), H Simmons (UAF) and L. St. Laurent (WHOI). Data/model integration and comparisons will be made with Simmons, Klymak, and Buijsman.

REFERENCES

- Klymak, J.M., J.N. Moum, J.D. Nash, E. Kunze, J.B. Girton, G.S. Carter, C.M. Lee, T.B. Sanford, and M.C. Gregg, 2006: An estimate of tidal energy lost to turbulence at the Hawaiian Ridge. *J. Phys. Oceanogr.*, 36, 1148-1164.
- Moum, J.N. and J.D. Nash, 2009: Mixing measurements on an equatorial ocean mooring, *J. Atmos. and Oceanic Tech.*, 26, 317-336.
- Nash, J.D., E. Kunze, C.M. Lee and T.B. Sanford, 2006: Structure of the baroclinic tide generated at Kaena Ridge, Hawaii, *J. Phys. Oceanogr.*, 36(6), 1123-1135.