

Analysis of the in-Water and Sky Radiance Distribution Data Acquired During the RaDyO Project

Kenneth J. Voss

Physics Department, University of Miami
Coral Gables, Fl. 33124

phone: (305) 284-2323 ext 2 fax: (305) 284-4222 email: voss@physics.miami.edu

Award Number: N000141110153

<http://optics.physics.miami.edu>

LONG-TERM GOALS

My work involves experimentally investigating the interrelationships and variability of optical properties in the ocean and atmosphere. My goal is to define the variability of the optical properties, particularly those dealing with light scattering, and to improve the prediction capabilities of image and radiative transfer models used in the ocean. My near term ocean optics objectives have been: 1) to improve the measurement capability of measuring the in-water and above-water spectral radiance distribution and extending this capability to polarization, 2) to investigate the variability of the Point Spread Function (PSF) as it relates to the imaging properties of the ocean, and 3) to improve the characterization of the Bi-directional Reflectance Distribution Function (BRDF) of benthic surfaces in the ocean, and 4) to understand the capabilities and limitations of using radiative transfer to model the BRDF of particulate surfaces.

OBJECTIVES

The major objective of this research is to understand the downwelling spectral polarized radiance distribution, in the near surface of the ocean.

APPROACH

We have built, with ONR support (through the DURIP program) a camera system capable of measuring the polarization state of the downwelling radiance distribution. This instrument follows in the footsteps of other instruments we have developed (Voss and Liu, 1997) and uses a combination of 3-4 images of the radiance distribution to form this polarized radiance distribution. Because the downwelling radiance distribution is very dynamic, we need to have a system that will quickly make these images as matched as possible, so this required a completely new design.

The system we have designed uses 4 fisheye camera lenses with coherent fiber bundles behind each image. Each fisheye will have a polarizer in a different orientation. After the image is in the coherent fiber bundle, these bundles will be brought together and imaged on a CCD array camera, through a filter changer (for spectral information). Thus in a single image we will have 4 separate fisheye images of the scene, each with different polarization information. This instrument was used in the

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2011	2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011		
4. TITLE AND SUBTITLE Analysis of the in-Water and Sky Radiance Distribution Data Acquired During the RaDyO Project			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Miami, Physics Department, Coral Gables, FL, 33124			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RaDyO program. The work of this proposal was to analyze the data acquired during the RaDyO program, and prepare articles for publication.

WORK COMPLETED

During the last year we have submitted 3 papers from our own work, and collaborated on an additional 4 papers with RaDyO co-investigators. 3 of these papers have been published, while the remaining papers are accepted and will be published in a JGR special issue including other RaDyO publications. Along with this, the graduate student working on the project finished his Ph.D. thesis and has graduated.

RESULTS

Our measurements found that the polarization behavior of the light field in water near the surface is dominated by the refracted sky light, and depends strongly on the wave-induced curvatures of the water surface. Therefore, near the surface the polarization is predominately inside the Snell's cone. At 1.5 m depth the maximum Degree of Linear Polarization, *DoLP*, for 520 nm, was found to be about 65%, similar to the skylight. Near the surface in clear water this *DoLP* had no clear spectral dependence; however, for more turbid waters, the path radiance from skylight decreases the *DoLP* in proportion to the available skylight. As one progresses in the water column the polarization due to light scattering by the water increases, thereby reducing the effect of refracted sky light. Thus, with the increase of the water depth, the maximum in the polarization moves from the refracted direction of 90° scattering in air, to 90-100 degree scattering angle in water, as expected from the oceanic Mueller matrix [Voss and Fry, 1984]. The peak of the radiance also shifts towards the center (zenith) away from the refracted solar position. This shift is more prominent when the water attenuation coefficient increases, as in coastal waters (SBC). The maximum *DoLP* also decreases with increasing depth. Our data showed that, at a comparable depth, the maximum *DoLP* in clear water is larger than in turbid water. We also found that the pattern of χ remains very constant with depth and location, which is interesting, as this is thought to be used by animals for navigation in some cases. This work is detailed in Bhandari et al. [2011].

Our data, for both the sky and in the water, was also used by the two major modeling groups in RaDyO, to successfully validate their radiative transfer models. This work is detailed in the papers by Yu et al. [2011] and Xu et al. [2011]

One very interesting thing that came out of our research over the past year was the observation of off-axis neutral points in the upwelling light field.[Voss et al., 2011] Neutral points in the atmosphere have been studied since the 1800's. They are very sensitive to changes in the atmospheric conditions and are only rarely found off axis. No one had experimentally observed these points off-axis neutral points in the upwelling in-water light field before. We found these in our measurements in the case of very clear water. Our early modeling efforts have indicated that the off-axis position of the neutral points is caused by the addition of the diffused skylight radiance adding to the polarized field of the direct solar radiance. This was a fun, and interesting, offshoot of our polarization measurements.

IMPACT/APPLICATIONS

The goal of the overall RadYO program is to understand how the radiance distribution is modified in the near surface, and what factors are important to this modification. Our work is showing how the

near surface polarized radiance distribution is modified as it is transmitted through the air-sea interface and then into the water column.

RELATED PROJECTS

This project was part of the overall ONR RadYO program. We also have had DURIP support to build the instrument, fundamental to this work. Our work on the polarized radiance distribution is also related to our efforts with NASA funding to look at both the upwelling radiance distribution and the polarized upwelling radiance distribution.

REFERENCES

P. Bhandari, K. J. Voss, L. Logan, and M. Twardowski, “The variation of the polarized downwelling radiance distribution with depth in the coastal and clear ocean, *J. Geophys. Res.*, May 2011. [Accepted, refereed]

K. J. Voss, A. C. R. Gleason, H. R. Gordon, G. W. Kattawar, and Y. You, “Observation of non-principal plane neutral points in the in-water upwelling polarized light field”, *Opt. Expr.*, **7**, 5942-5952 (2011). [published, refereed]

Z. Xu, D. K. P. Yue, L. Shen, K. J. Voss, “Patterns and Statistics of In-water Polarization under conditions of linear and nonlinear ocean surface waves,” *J. Geophys. Res.*, May 2011. [Accepted, refereed]

Y. You, G. W. Kattawar, K. J. Voss, P. Bhandari, J. Wei, M. Lewis, C. Zappa, and H. Schultz, “Polarized light field under dynamic ocean surfaces: numerical modeling compared with measurements”, *J. Geophys. Res.*, May 2011. [Accepted, refereed]

PUBLICATIONS

P. Bhandari, K. J. Voss, and L. Logan, “An instrument to measure the downwelling polarized radiance distribution in the ocean”, *Optics Express*, **19**, 17609-17620 (2011). [published, refereed]

T. D. Dickey, G. W. Kattawar, and K. J. Voss, “Shedding new light on light in the ocean”, *Physics Today*, **64**, April, 44-49 (2011). [published, refereed]

K. J. Voss, A. C. R. Gleason, H. R. Gordon, G. W. Kattawar, and Y. You, “Observation of non-principal plane neutral points in the in-water upwelling polarized light field”, *Opt. Expr.*, **7**, 5942-5952 (2011). [published, refereed]

P. Bhandari, K. J. Voss, L. Logan, “The variation of the polarized downwelling radiance distribution with depth in coastal water, *Ocean Optics XX*, Anchorage, Al. October, 2010.

M. Lewis, J. Wei, R. Van Dommelen, and K. J. Voss, “Quantitative estimation of the underwater radiance distribution”, *J. Geophys. Res.*, May, 2011. [Accepted, refereed]

Y. You, G. W. Kattawar, K. J. Voss, P. Bhandari, J. Wei, M. Lewis, C. Zappa, and H. Schultz, “Polarized light field under dynamic ocean surfaces: numerical modeling compared with measurements”, J. Geophys. Res., May 2011. [Accepted, refereed]

Z. Xu, D. K. P. Yue, L. Shen, K. J. Voss, “Patterns and Statistics of In-water Polarization under conditions of linear and nonlinear ocean surface waves,” J. Geophys. Res., May 2011. [Accepted, refereed]

P. Bhandari, K. J. Voss, L. Logan, and M. Twardowski, “The variation of the polarized downwelling radiance distribution with depth in the coastal and clear ocean, J. Geophys. Res., May 2011. [Accepted, refereed]