

Optical Observations with Milliarcsecond Resolution of Stars, Their Environments and Companions

Jason J. Sanborn^{1,2†} and Robert T. Zavala³

¹Lowell Observatory, Flagstaff, AZ 86001, USA

²Northern Arizona University, Flagstaff, AZ 86001, USA

³United States Naval Observatory Flagstaff Station, Flagstaff, AZ 86001, USA

Observations with milliarcsecond resolution using the Navy Optical Interferometer have been obtained for a number of stellar systems which include high-mass binaries, eclipsing binaries, and radio stars. These observations also reveal the previously unseen companions in single-lined spectroscopic binaries via directly measured flux ratios. We will present examples of published and ongoing research efforts of these systems to illustrate how an optical interferometer contributes to our knowledge of stars, their environment, and companions. These studies include a conclusive revealing of the previously unseen companion in the single-lined binary Φ Herculis, the direct determination of orbital parameters in the wide and close orbits of Algol, and revealing the orbit of β Lyrae with spatially resolved images of the H α emission.

Keywords: optical interferometry, binaries, astrometry, Navy Optical Interferometer

1. INTRODUCTION

The Navy Optical Interferometer (NOI), formerly the Navy Prototype Optical Interferometer (NPOI), is a joint project of the Naval Research Laboratory (NRL), and the United States Naval Observatory (USNO) in cooperation with Lowell Observatory, located at Anderson Mesa near Flagstaff, Arizona. For further instrument details see Armstrong et al. (1998) and for details regarding NOI's observations and data reductions see Hummel et al. (2003), and the references therein.

Optical interferometry offers a method of directly determining stellar diameters, the morphology of H α emission regions, and astrometric orbital parameters that govern the interactions of binary systems. The fundamental observed quantity for the NOI is the squared visibility (V^2 , Armstrong et al. 1998), which is the Fourier transform of the source brightness distribution (see Thompson et al. 2001 for an introduction to interferometry). After detecting interference fringes from a binary star an orbit may be

fit (Pan et al. 1993). With the orbit, the visibility data from all observations can be used to fit for the magnitude differences of the binary system in question (Hummel et al. 2003). Results of this technique as applied to the binary stars Phi Herculis, Beta Lyrae and Algol are reviewed in this work.

2. RESOLUTION OF SINGLE-LINED SPECTROSCOPIC BINARIES: Φ HERCULIS

Φ Herculis (HD 145389, HR 6023) is a mercury-manganese (HgMn) star, showing a wide variety of spectral abundance anomalies with both depletions and enhancements (Adelman et al. 2004). Aikman (1976) identified Φ Her as a single-lined spectroscopic binary. The secondary component avoided detection until NOI observations became available beginning in 1998. These interferometric observations lead to the identification of the spectral type of the secondary. Identification of the secondary's spec-

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Jan 11, 2012 Revised Jan 30, 2012 Accepted Jan 31, 2012

[†]Corresponding Author

E-mail: jaisen1@lowell.edu

Tel: +1-928-310-6840 Fax: +1-928-774-6296

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2012		2. REPORT TYPE		3. DATES COVERED 00-00-2012 to 00-00-2012	
4. TITLE AND SUBTITLE Optical Observations With Milliarsecond Resolution Of Stars, Their Environments And Companions			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Naval Observatory Flagstaff Station,Flagstaff,AZ,86001			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES J. Astron. Space Sci. 29(1), 63-67 (2012)					
14. ABSTRACT Observations with milliarsecond resolution using the Navy Optical Interferometer have been obtained for a number of stellar systems which include high-mass binaries, eclipsing binaries, and radio stars. These observations also reveal the previously unseen companions in single-lined spectroscopic binaries via directly measured flux ratios. We will present examples of published and ongoing research efforts of these systems to illustrate how an optical interferometer contributes to our knowledge of stars, their environment, and companions. These studies include a conclusive revealing of the previously unseen companion in the single-lined binary &#934; Herculis, the direct determination of orbital parameters in the wide and close orbits of Algol, and revealing the orbit of &#946; Lyrae with spatially resolved images of the H&#945; emission.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

tral type enables a better understanding of the chemical abundances of the primary.

In Fig. 1 the calibrated V^2 data shows the cosine wave signature of a binary source (Michelson 1920) as seen with an interferometer. As described in Pan et al. (1993) and Hummel et al. (2003), a fit for the binary parameters was applied and a measure of the magnitude difference of the secondary relative to the primary was found. With the assumption that both stars are on the main sequence and with the determination of the magnitude differences, 2.57 in the V band and 2.39 in the R band, a prediction of the secondary's spectral type as A8 V was found. This spectral type was verified spectroscopically with follow up observations utilizing Dominion Astrophysical Observatory (DAO) spectra (Zavala et al. 2007). An independent confirmation of the secondary spectral type was also made by Dworetzky & Willatt (2006).




Fig. 1. The calibrated squared visibilities of Φ Her for Navy Optical Interferometer observations made on 1998 May 16. This data shows the characteristic cosine wave signature of a binary.




Fig. 2. The left panel shows the observed spectra overlaid on normalized synthetic spectra for Φ Her A (solid blue line) and Φ Her B (red dashed line). In these images we can see the primary and secondary components of Mg I (2) 5183.6042. In the right panel is the same data but now is overlaid with a line representing the summed synthetic spectra for Φ Her A and B.

Fig. 2 illustrates how the synthetic spectra of the components, constrained by the brightness ratio from the NOI, accurately represent the observed spectra. In the right panel of Fig. 2 the DAO spectra are shown with an overlay of the summed synthetic spectra for both Φ Her A and B. In the left panel of Fig. 2 the DAO observed spectra is overlaid on individual normalized synthetic spectra for Φ Her A and B.

3. HA EMISSION IMAGING AND ORBIT DETERMINATION: B LYRAE

β Lyrae is an interacting eclipsing binary system consisting of a ~ 3 solar mass B6-8II star which has filled its Roche lobe and a ~ 13 solar mass early B star, which is completely obscured and hidden within a thick accretion disk (Harmanec 2002). In 2002 the NOI observed β Lyrae as part of an experiment to detect the binarity of this system and the $H\alpha$ emission associated with the accretion stream. NOI data from this experiment is presented along with images indicating the position of the $H\alpha$ emitting regions relative to the continuum photo-center as a function of orbital phase, providing a means for analyzing the $H\alpha$ morphology (Schmitt et al. 2009).

In Fig. 3 (left panel) the position of the $H\alpha$ emission is plotted relative to the continuum photo-center for the five nights of data. The orbit of the $H\alpha$ emission is also plotted by fitting a line to the data points with an orbital solution found to be oriented along a position angle of $248.8 \pm 1.7^\circ$. Using parameters derived by Harmanec et al. (1996), Linnell (2000), and Harmanec (2002), the orbits of the $H\alpha$ and continuum photo-centers were fit and converted to positions in the sky using the inclination de-




Fig. 3. Separation of the two binary components (left panel, blue data points with larger errors), with each point showing the uncertainty in the position of the separation of the two sources. Besides each point an indication of the orbital phase and the date is given along with a dotted line representing a least squares fit to the data. The dotted line gives the orbit in the plane of the sky. The position of the peak of H α emission relative to the continuum photo-center is also displayed within the left panel (red square data points). The series of panels on the right show the H α images of β Lyrae. Each panel corresponds to a different night and is organized, in a clockwise fashion, of increasing orbital phase. The white star points to the photo-center of the system. The lowest contour in each figure corresponds to the 3 σ level. The reconstructing beam size and orientation is shown in the bottom left corner of each panel.

rived by Linnell et al. (1998). One important observational result is the H α semi-major axis was observed to be less than the continuum semi-major axis, providing evidence that the NOI was effectively imaging the orbit of this system. This is due to the H α emission originating within the disk of the more massive star, and the continuum photo-center being located nearer the less massive star (Schmitt et al. 2009). These H α images each correspond to a different night and are organized in order of increasing orbital phase and are also shown with the correct orbital orientation.

4. DIRECT DETERMINATION OF ORBITAL PARAMETERS: ALGOL

Algol is the prototype for a well-known class of eclipsing binaries. The Algol system is a hierarchical triple system (close binary Algol A-B and triple Algol AB-C) and is well summarized by Söderhjelm (1980). Speckle (Labeyrie

et al. 1974) and optical (Pan et al. 1993) interferometry resolved the orbit of the AB-C component, but the ascending node of these orbits differed by 180° from that used in the High Precision PARallaxCollectingSatellite (HIPPARCOS) double star solution (Lindgren et al. 1997). As Algol is 1 of 12 radio stars used to link the HIPPARCOS optical reference frame to the International Celestial Reference System (ICRS, Kovalevsky et al. 1997), a resolution of this 180° ambiguity was needed to reconcile the time-variable systematic offset of the photo-center between the HIPPARCOS orbital orientation and the interferometric orientation.

Interferometric observations using the NOI have resulted in the first resolved images of the triple system, as seen in Figs. 4 and 5. These observations have led to an unambiguous determination that the close pair orbit (A-B) is retrograde and nearly orthogonal to the plane of the wide (AB-C) orbit (Zavala et al. 2010). The improvements made to the AB-C orbital plane orientation have removed the time-variable systematic offset expected




Fig. 4. Images of the Algol triple system made from Navy Optical Interferometer data on 2006 Oct 29 are seen in the left panel. In the right panel observations from 2006 Oct 30 are shown to emphasize the motion of Algol B between the two epochs. The Algol C component can be found in the upper right hand of each panel. To help guide the eye, the approximate positions of Algol B are indicated at each epoch by a filled in red circle.




Fig. 5. The left panel is an illustration showing the Navy Optical Interferometer (NOI) astrometric results for the AB-C orbit superposed on top of previously determined orbits from Mark III and CHARA observations. In the right panel an illustration of NOI astrometric results are plotted with the astrometry of Pan et al. (1993) and rotated by 180° . A vector from the origin indicates the periastron point.

between the orbit used in the HIPPARCOS solution and the results of Pan et al. (1993) and Csizmadia et al. (2009). NOI observations have also led to the first directly measured magnitude differences for the three stars in Algol. Estimates made in previous studies were performed by modeling photometric light curve and spectroscopic data and spanned more than one magnitude, from 3.92 ± 0.88 (Richards et al. 1988) to 2.71 ± 0.15 (Kim 1989). The NOI directly measured V band magnitude difference favored a magnitude less than 3 (Zavala et al. 2010).

NOI magnitude differences help to constrain results obtained by modeling photometric light curves and spectroscopic data by adding directly measured Δ -magnitude results. Due to the NOIs capability of directly measuring position angles and magnitude differences in close binaries, further observations of double and triple star

systems will also benefit from the directly measured fundamental parameters that govern these dynamic stellar systems.

REFERENCES

- Adelman SJ, Proffitt CR, Wahlgren GM, Leckrone DS, Dolk L, Heavy element abundances in late-B and early-A stars. I. Co-added IUE spectra of HgMn stars, *ApJS*, 155, 179-189 (2004). <http://dx.doi.org/10.1086/424802>
- Aikman GCL, The spectroscopic binary characteristics of the mercury-manganese stars, *PDAO*, 14, 379-410 (1976).
- Armstrong JT, Mozurkewich D, Rickard LJ, Hutter DJ, Benson JA, et al., The Navy Prototype Optical Interferometer, *ApJ*, 496, 550-571 (1998). <http://dx.doi.org/10.1086/305365>

- Csizmadia Sz, Borkovits T, Paragi Zs, Ábrahám P, Szabados L, et al., Interferometric observations of the hierarchical triple system Algol, *ApJ*, 705, 436-445 (2009). <http://dx.doi.org/10.1088/0004-637X/705/1/436>
- Dworetzky MM, Willatt R, The secondary star of the HgMn binary Phi Her (2006). [arXiv:astro-ph/0612432](http://arxiv.org/abs/astro-ph/0612432)
- Harmanec P, The ever challenging emission-line binary beta Lyrae, *AN*, 323, 87-98 (2002). [http://dx.doi.org/10.1002/1521-3994\(200207\)323:2<87::AID-ASNA87>3.0.CO;2-P](http://dx.doi.org/10.1002/1521-3994(200207)323:2<87::AID-ASNA87>3.0.CO;2-P)
- Harmanec P, Morand F, Bonneau D, Jiang Y, Yang S, et al., Jet-like structures in β Lyrae. Results of optical interferometry, spectroscopy and photometry, *A&A*, 312, 879-896 (1996).
- Hummel CA, Benson JA, Hutter DJ, Johnston KJ, Mozurkewich D, et al., First observations with a co-phased six-station optical long-baseline array: application to the triple star η Virginis, *AJ*, 125, 2630-2644 (2003). <http://dx.doi.org/10.1086/374572>
- Kim H-I, BV light curve analysis of Algol, *ApJ*, 342, 1061-1067 (1989). <http://dx.doi.org/10.1086/167662>
- Kovalevsky J, Lindegren L, Perryman MAC, Hemenway PD, Johnston KJ, et al., The HIPPARCOS catalogue as a realisation of the extragalactic reference system, *A&A*, 323, 620-633 (1997).
- Labeyrie A, Bonneau D, Stachnik RV, Gezari DY, Speckle interferometry. III. High-resolution measurements of twelve close binary systems, *ApJ*, 194, L147-L151 (1974). <http://dx.doi.org/10.1086/181689>
- Lindegren L, Mignard F, Söderhjelm S, Badioli M, Bernstein, H-H, et al., Double star data in the HIPPARCOS catalogue, *A&A*, 323, L53-L56 (1997).
- Linnell AP, Progress on a model for β Lyrae, *MNRAS*, 319, 255-266 (2000). <http://dx.doi.org/10.1046/j.1365-8711.2000.03898.x>
- Linnell AP, Hubeny I, Harmanec P, New light synthesis and spectrum synthesis constraints on a model for beta Lyrae, *ApJ*, 509, 379-391 (1998). <http://dx.doi.org/10.1086/306484>
- Michelson AA, On the application of interference methods to astronomical measurements, *ApJ*, 51, 257-262 (1920). <http://dx.doi.org/10.1086/142550>
- Pan X, Shao M, Colavita MM, High angular resolution measurements of Algol, *ApJ*, 413, L129-L131 (1993). <http://dx.doi.org/10.1086/186976>
- Richards MT, Mochnaki SW, Bolton CT, Nonsimultaneous multicolor light-curve analysis of Algol (beta Persei), *AJ*, 96, 326-336 (1988). <http://dx.doi.org/10.1086/114810>
- Schmitt HR, Pauls TA, Tycner C, Armstrong JT, Zavala RT, et al., Navy Prototype Optical Interferometer imaging of line emission regions of β Lyrae using differential phase referencing, *ApJ*, 691, 984-996 (2009). <http://dx.doi.org/10.1088/0004-637X/691/2/984>
- Söderhjelm S, Geometry and dynamics of the Algol system, *A&A*, 89, 100-112 (1980).
- Thompson AR, Moran JM, Swenson GW, Interferometry and synthesis in radio astronomy, 2nd ed. (Wiley, New York, 2001).
- Zavala RT, Adelman SJ, Hummel CA, Gulliver AF, Caliskan H, et al., The mercury-manganese binary star ϕ Herculis: detection and properties of the secondary and revision of the elemental abundances of the primary, *ApJ*, 655, 1046-1057 (2007). <http://dx.doi.org/10.1086/510108>
- Zavala RT, Hummel CA, Boboltz DA, Ojha R, Shaffer DB, et al., The Algol triple system spatially resolved at optical wavelengths, *ApJ*, 715, L44-L48 (2010). <http://dx.doi.org/10.1088/2041-8205/715/1/L44>