

Native Plant Species Suitable for Ecological Restoration

Thomas Smith

U.S. Army, Engineer Research and Development Center,
Champaign, IL

David Wester

Texas Tech University, Lubbock, TX

Sandra Rideout-Hanzak

Texas A & M University, Kingsville, TX

May 10, 2011




US Army Corps of Engineers
BUILDING STRONG



Photo credit: David Wester

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 10 MAY 2011		2. REPORT TYPE		3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE Native Plant Species Suitable for Ecological Restoration				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S . Army Engineer Research and Development Center,Construction Engineering Research Laboratory,PO Box 9005,Champaign,IL,61826-9005				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the NDIA Environment, Energy Security & Sustainability (E2S2) Symposium & Exhibition held 9-12 May 2011 in New Orleans, LA.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 25	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Engineer Research and Development Center



Background

- DoD manages approximately 25 million acres in US
- Large training mission component
- Variety of ecosystems



Military Range Landscape



TEXAS TECH UNIVERSITY
Department of Natural
Resources Management



BUILDING STRONG®

Western Ragweed



Cheatgrass and Crested Wheatgrass



Terminology

- Restoration - *The process of repairing man caused damage to the diversity and dynamics of indigenous ecosystems*
- Revegetation - *Reestablishment of vegetation*
- Reclamation - *Returning the community to the conditions are comparable to that prior to disturbance*
- Rehabilitation - *A partial movement along the trajectory from degraded to the original state*
- Remediation, mitigation - *Activities that lessen the degree of damage*


Ecoregions

Collapsed Bailey's 19 divisions into the following ecoregions

- Grasslands
 - ▶ Temperate Steppe (Central and Northern Great Plains)
 - ▶ Tropical/Subtropical (Southern Great Plains, Rio Grande Plains, Coastal Prairies)
 - ▶ Prairie (Tallgrass prairie)
- Southwestern U.S.
 - ▶ Temperate Desert (Great Basin)
 - ▶ Tropical/Subtropical Desert (Chihuahuan, Sonoran deserts)
- Mediterranean
 - ▶ California Grasslands
 - ▶ California Chaparral
- Western Mountains (Rocky Mountains)
- Northwestern Mountains (Pacific Northwest)
- Humid Temperate Domain: Warm Continental Division
- Humid Temperate Domain: Hot Continental Division
- Humid Temperate Domain: Hot Continental Division (Mountains)
- Humid Temperate Domain: Subtropical Division
- Savannah Division (southern Florida)

Ecoregions

(Modified from Bailey (1995) –Ecoregions of the United States)



Approach

- For each major division, we surveyed relevant professional, technical, and other literature to document existing resources from which to develop species lists. Additionally, scientific (peer-reviewed) literature and “gray literature” (published technical reports and documents produced by government agencies and non-governmental research groups)
- Native plant species suitable for ecological restoration purposes were identified for each ecological unit. Several non-native but widely naturalized species also were included.



BUILDING STRONG®

Approach

- Species distribution data were compiled from Stubbendieck et al. 2003, the PLANTS database (<http://plants.usda.gov/>), the Fire Effects Information system (<http://www.fs.fed.us/database/feis>), and Vallentine (1989).
- Species were further described with respect to specific management considerations
 - ▶ growth form
 - ▶ season of growth
 - ▶ ease of establishment,
 - ▶ longevity,
 - ▶ soil adaptation,
 - ▶ drought and salinity tolerance,
 - ▶ cold-hardiness, and
 - ▶ erosion control


Approach

- Growth form -- graminoid, forb, shrub, tree
- Season of growth -- warm season, cool season, evergreen
- Ease of establishment -- synthesis of characteristics – e.g., seed viability, germination, seedling vigor
- Longevity -- stand survival
- Drought tolerance -- relative value judgment
- Salinity tolerance -- $< 10\%$ reduction in plant growth
- Cold hardiness -- minimal tolerable temperature = lowest temperature recorded in historical range

Approach

Soil adaptation

Erosion control





ECOLOGY OF PLANTS, Second Edition, Figure 4.1 (Part 2) © 2006 Sinauer Associates, Inc.



Approach

Cold hardiness



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Results

Summary of results of data study, organized by common name.

Common and scientific name	Growth		Soil			Plant characteristics						Region												Comments	
	Growth form	Season of growth	Sandy	Silty	Clayey	Ease of establishment	Longevity	Drought tolerance	Cold hardiness	Salinity tolerance	Erosion control	Temperate Steppe	Tropical/Subtropical	Prairie	Temperate Desert	Tropical/subtropical desert California Grasslands	Humid Temperate: Subtropical	Humid Temperate: Hot continental	Humid Temperate: Mountain segment	Humid Temperate: Warm continental	Western mountains: Rocky Mountains	Western mountains: Northwestern Mtns	Western Mountains: California chaparral		Southern Florida: Savanna
Alfalfa (<i>Medicago sativa</i>)	F	W	1	1	2	1	1	2	2	2	1	•	•	•	•	•									Most widely planted legume; over 400 varieties available; an introduced species but widely naturalized
Arizona cottontop (<i>Digitaria californica</i>)	F	W	1	1	2	2	1	1	2	2	1		•			•									Variety "Loetta" suitable for southwestern US
Arrowleaf balsamroot (<i>Balsamorhiza sagittata</i>)	F	C	2	1	1	2	1	1	1	3	1	•			•	•									
Bitterbrush (<i>Purshia tridentata</i>)	S	E	1	2	3	2	1	1	1	2	1	•			•	•					•	•	•		Varieties "Lassen," "Fountain Green," and "Maybell"
Black cherry (<i>Prunus serotina</i>)	T	W	1	2	2	1	2	2	1	2	1			•		•		•	•	•	•				



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Results

[illegible]Department of Natural
Resources Management

Results

Common and scientific name	Growth		Soil			Plant characteristics						Region												Comments		
	Growth form	Season of growth	Sandy	Silty	Clayey	Ease of establishment	Longevity	Drought tolerance	Cold hardness	Salinity tolerance	Erosion control	Temperate Steppes	Tropical/Subtropical	Prairie	Temperate Desert	Tropical/subtropical desert	California Grasslands	Humid Temperate: Subtropical	Humid Temperate: Hot continental	Humid Temperate: Mountain segment	Humid Temperate: Warm continental	Western mountains: Rocky Mountains	Western mountains: Northwestern Mtns		Western Mountains: California chaparral	Southern Florida: Savanna
Bluestem, little (<i>Schizachyrium scoparium</i>)	G	W	1	1	1	2	1	1	1	3	1	•	•	•		•		•	•	•	•				•	Varieties "Aldous," "Camper," "Cimmaron," "Pastura" and "Blaze" suitable for central Great Plains.
Bluestem, sand (<i>Andropogon gerardii</i> var. <i>paucipilus</i> ; <i>A. hallii</i>)	G	W	1	2	3	2	1	2	1	3	1	•	•	•		•										Varieties "Elida," "Goldstrike," and "Garden" suitable for northern to southern Great Plains; variety "Woodward" suitable for southern Great Plains.
Boneset, common (<i>Eupatorium perfoliatum</i>)	F	W	1	1	1	1	1	2	1	1	1	•		•				•	•	•	•					
Brome, meadow (<i>Bromus biebersteinii</i>)	G	C	1	1	1	2	1	2	1	3	1	•		•												"Regar" – not native



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Bluestems



BUILDING STRONG®

Results

Common and scientific name	Growth		Soil			Plant characteristics						Region												Comments	
	Growth form	Season of growth	Sandy	Silty	Clayey	Ease of establishment	Longevity	Drought tolerance	Cold hardness	Salinity tolerance	Erosion control	Temperate Steppe	Tropical/Subtropical	Prairie	Temperate Desert	Tropical/subtropical desert	California Grasslands	Humid Temperate: Subtropical	Humid Temperate: Hot continental	Humid Temperate: Mountain segment	Humid Temperate: Warm continental	Western mountains: Rocky Mountains	Western mountains: Northwestern Mtns		Western Mountains: California chaparral
Honey locust (<i>Gleditsia triacanthos</i>)	T	W	3	1	1	1	1	1	1	3	1							•	•	•					
Huckleberry, grouse (<i>Vaccinium scoparium</i>)	S	?	1	1	2	1	1	2	2	3	1	•									•	•			
Indiangrass (<i>Sorghastrum nutans</i>)	G	W	1	1	1	2	1	2	1	2	1	•	•	•		•		•	•	•	•				Varieties "Holt" (NE), "Llano" (NM), "Lometa" (TX), "Osage" (KS/OK), "Oto" (NE/KS) and "Tomahawk" (ND/SD)
Indiangrass, lopsided (<i>Sorghastrum secundum</i>)	G	W	1	2	3	2	2	2	1	2	1						•							•	
Indian ricegrass (<i>Stipa hymenoides</i>)	G	C	1	1	2	2	1	1	1	2	1	•		•	•	•									Varieties "Nezpar," "Paloma" and "Rimrok"
Leadplant (<i>Amorpha canescens</i>)	S	W	1	1	2	1	1	1	1	3	1									•					



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Results

Common and scientific name	Growth		Soil		Plant characteristics		Region																Comments			
	Growth form	Season of growth	Sandy	Silty	Clayey	Ease of establishment	Longevity	Drought tolerance	Cold hardiness	Salinity tolerance	Erosion control	Temperate Steppe	Tropical/Subtropical	Prairie	Temperate Desert	Tropical/subtropical desert	California Grasslands	Humid Temperate: Subtropical	Humid Temperate: Hot continental	Humid Temperate: Mountain segment	Humid Temperate: Warm continental	Western mountains: Rocky Mountains		Western mountains: Northwestern Mtns	Western Mountains: California chaparral	Southern Florida: Savanna
Pine, ponderosa (<i>Pinus ponderosa</i>)	T	E	1	1	3	1	1	1	1	2	1				•	•						•	•	•		
Prairie blazing star (<i>Liatris pycnostachya</i>)	F	W	1	1	1	1	1	2	1	3	1			•												Variety "Eureka"
Needle and thread (<i>Hesperostipa comata</i>)	G	C	1	1	2	1	1	1	1	1	1	•			•	•	•									
Needlegrass, Columbia (<i>Achnatherum nelsonii</i>)	G	C	2	1	1	2	1	1	1	2	1	•			•											
Needlegrass, foothill (<i>Nassella lepida</i>)	G	C	1	2	3	1	1	2	2	3	1						•									
Needlegrass, green (<i>Nassella viridula</i>)	G	C	3	1	1	2	2	1	1	2	1	•			•								•			
Needlegrass, nodding (<i>Nassella cernua</i>)	G	C	2	1	1	1	1	1	2	3	1						•							•		
Needlegrass, purple (<i>Nassella pulchra</i>)	G	C	1	1	1	1	1	1	2	3	1						•							•		



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Needle and Thread



Results

Common and scientific name	Growth		Soil			Plant characteristics				Region												Comments				
	Growth form	Season of growth	Sandy	Silty	Clayey	Ease of establishment	Longevity	Drought tolerance	Cold hardiness	Salinity tolerance	Erosion control	Temperate Steppe	Tropical/Subtropical	Prairie	Temperate Desert	Tropical/subtropical desert	California Grasslands	Humid Temperate: Subtropical	Humid Temperate: Hot continental	Humid Temperate: Mountain segment	Humid Temperate: Warm continental		Western mountains: Rocky Mountains	Western mountains: Northwestern Mtns	Western Mountains: California chaparral	Southern Florida: Savanna
Penstemon, Rocky Mountain (<i>Penstemon strictus</i>)	F		1	2	3	1		2	1	3	1				*											Variety "Bandera" suited for clayey soils
Prairie junegrass (<i>Koeleria macrantha</i>)	G	C	1	2	2	1	1	1	1	3	1	*	*	*	*	*	*			*	*	*	*	*		
Prairie willow (<i>Salix humilis</i>)	S		1	2	3	2	1	3	1	3	1			*				*	*	*						
Red fescue (<i>Festuca rubra</i>)	G	C	3	1	1	1	1	2	1	3	1	*			*	*	*			*	*	*	*	*		
Rose, wild (<i>Rosa woodsii</i>)	S	C	1	1	2	1	1	2	1	3	1	*			*	*					*		*			
Russet buffaloberry (<i>Shepherdia canadensis</i>)	S	C	1	1	2	3	1	1	1	2	1	*	*													
Sacaton, alkali (<i>Sporobolus airoides</i>)	G	W	1	1	1	2	1	2	1	1	1	*	*		*	*	*									Suitable for alkaline or saline soils
Sagebrush, big (<i>Artemisia tridentata</i>)	S	E	3	1		2	1	1	1	2	1	*			*	*					*		*			Variety "Hobblecreek"



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Results

Common and scientific name	Growth		Soil			Plant characteristics						Region													Comments	
	Growth form	Season of growth	Sandy	Silty	Clayey	Ease of establishment	Longevity	Drought tolerance	Cold hardiness	Salinity tolerance	Erosion control	Temperate Steppe	Tropical/Subtropical	Prairie	Temperate Desert	Tropical/subtropical desert	California Grasslands	Humid Temperate: Subtropical	Humid Temperate: Hot continental	Humid Temperate: Mountain segment	Humid Temperate: Warm continental	Western mountains: Rocky Mountains	Western mountains: Northwestern Mtns	Western Mountains: California chaparral		Southern Florida: Savanna
Switchgrass (<i>Panicum virgatum</i>)	G	W	1	1	1	2	1	2	1	2	1	•	•	•				•	•	•	•				•	Many varieties available (e.g., "Alamo" (TX), "Blackwell" (OK), "Dacotah" (ND), "Forestburg" (SD), "Kanlow" (OK), "Nebraska 28" (NE), "Grenville" (NM))
Sycamore (<i>Platanus occidentalis</i>)	T	W	1	1	2	1	1	3	1	3	1							•	•	•	•					
Tall tickseed (<i>Coreopsis tripteris</i>)	F	W	1	1	2	1	1	1	1	1	1							•	•	•	•					
Tufted hairgrass (<i>Deschampsia caespitosa</i>)	G	C	1	1	1	2	1	3	1	3	1			•		•					•	•	•	•		
Tulip poplar (<i>Liriodendron tulipifera</i>)	T	W	1	1	2	1	1	3	2	3	1							•	•	•	•					



TEXAS TECH UNIVERSITY

Department of Natural
Resources Management



BUILDING STRONG®

Other Resources

- Bailey, R.G. 1995. Ecoregions. U.S. Forest Service. <http://www.fs.fed.us/land/ecosystem/>
- Stubbenbeck, J., S.L. Hatch, and L.M Landholt. 2003. North American wildland plants. In 6th ed. *North American Range Plants*. U. of Nebraska Press, Lincoln, NE.
- U.S. Department of Agriculture. PLANTS data base. <http://plants.usda.gov/>
- U.S. Department of Agriculture, Forest Service. Fire Effects Information System. <http://www.fs.fed.us/database/feis>
- U.S. Department of the Army. 2011. Guidance on native plant species
- Valentine, J.F. 1989. Range development and improvements. 3rd ed. Academic Press, New York, NY.



BUILDING STRONG®

Questions?

Thomas.Smith@usace.army.mil
U.S. Army Corps of Engineers, ERDC-CERL
Champaign, IL
217-373-5898



BUILDING STRONG®