

REPORT DOCUMENTATION PAGE					<i>Form Approved</i> OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.						
1. REPORT DATE (DD-MM-YYYY) 14-10-2011		2. REPORT TYPE Final Report			3. DATES COVERED (From - To) March 01,2008-May 31, 2011	
4. TITLE AND SUBTITLE Numerical Solution of Optimal Control Problem under SPDE Constraints				5a. CONTRACT NUMBER FA9550-08-1-0119		
				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Hongmei Chi (PI) Yanzhao Cao (Co-PI)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Florida A& M University 400 Foot-Hilyer Admin Cntr Tallahassee, FL 32307-3200					8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Office Of Scientific Research Research 875 North Randolph Street Arlington,VA 22203-1768					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-OSR-VA-TR-2012-0861	
12. DISTRIBUTION/AVAILABILITY STATEMENT Distribution A						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT The overall goal of this AFOSR sponsored research program is to construct fast and efficient numerical algorithms for solving stochastic partial differential equations and apply them to solve optimal control problems under uncertainty which is described by stochastic partial differential equations. It is well understood that effective numerical methods for stochastic partial differential equations (SPDES) are essential for uncertainty quantification. In the last decade much progress has been made in the construction of numerical algorithms to efficiently solve SPDES with random coefficients and white noise perturbations. However, high dimensionality and low regularity are still the bottleneck in solving real world applicable SPDES with efficient numerical methods. This project is intended to address the mathematical aspects of numerical approximations of SPDES, including error analysis and complexity analysis and development of new efficient numerical algorithms.						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF: a. REPORT b. ABSTRACT c. THIS PAGE			17. LIMITATION OF ABSTRACT		18. NUMBER OF PAGES	
					19a. NAME OF RESPONSIBLE PERSON	
					19b. TELEPHONE NUMBER (Include area code)	

AFOSR FINAL REPORT

Principal Investigator:

Hongmei Chi
Department of Computer and Information Sciences
Florida A&M University
Tallahassee, Florida, 32307

Co-Principal Investigator:

Yanzhao Cao
Department of Mathematics & Statistics
Auburn University
Auburn, AL 36849

Research Title:

Numerical Solutions of Optimal Control Problems
Under SPDES Constraints

Grant No:

FA9550-08-1-0119

Period of Performance:

April 15, 2008 –May 31, 2011

Numerical Solutions for Optimal Control Problems under SPDE Constraints

FA9550-08-1-0119

Hongmei Chi¹ and Yanzhao Cao²

*¹Department of Computer and Information Sciences
Florida A&M University, Tallahassee, FL 32307*

*²Department of mathematics
Auburn University, Auburn, AL 36849*

1 Introduction

The overall goal of our AFOSR sponsored research program is to construct fast and efficient numerical algorithms for solving stochastic partial differential equations and apply them to solve optimal control problems under uncertainty which is described by stochastic partial differential equations.

Uncertainty quantification has been an active research area in the past 10 years with many significant applications ranging from signal processing to aircraft wing designs. It is well understood that effective numerical methods for stochastic partial differential equations (SPDES) are essential for uncertainty quantification. In the last decade much progress has been made in the construction of numerical algorithms to efficiently solve SPDES with random coefficients and white noise perturbations. However, high dimensionality and low regularity are still the bottleneck in solving real world applicable SPDES with efficient numerical methods. This project is intended to address the mathematical aspects of numerical approximations of SPDES, including error analysis and complexity analysis and development of new efficient numerical algorithms. Our contributions toward this objective include

(i). *Optimal Kronecker sequence for the quasi Monte Carlo method and the Monte Carlo method for stochastic Stokes equations.* The generation of appropriate high-

quality quasirandom sequences (low-discrepancy sequences) is crucial to the success of quasi-Monte Carlo methods. We present a new algorithm for finding an optimal Kronecker sequence within choices of irrationals. we illustrated with numerical experiments why our algorithm is efficient for breaking these correlations.

(ii). *Efficient algorithms for stochastic partial differential equations.* We are interested in the the orthogonal polynomial expansion of **d-variable** functions. Fast and efficient algorithms of evaluating such expansions are essential for efficiently solving stochastic partial differential equations with spectral methods where d may be fairly large.

(iii). *High order numerical method for nonlinear filter problems.* We first convert the nonlinear filter problem into a problem of solving a stochastic partial differential equations (SPDES). We then construct a high order method to solve the SPDES and apply our algorithm to solve a ship tracking problem.

2 Brief Overview of Accomplishments

During the prior three-year grant period we have published 12 papers that have been supported by AFOSR funding. Selections of our current accomplishments are summarized here to provide an introduction to the concepts we have developed in the past and upon which future planned studies are based.

2.1 The optimal Kronecker Sequence for quasi Monte Carlo simulations

Kronecker sequences are easy and fast to implement. However, problems with Kronecker sequences arise from correlations between the choice of parameters which are used for different dimensions. These correlations cause Kronecker sequences to have poor two-dimensional projections for some pairs of dimensions.




Figure 1: Left: 1000 points of dimension 1 and 2 projection for original Kronecker sequence; right: 1000 points of dimension 26 and 27 projection for original Kronecker sequence.

In Figure 1, we can see high correlations and similarities in the poor two-dimensional projections for original Kronecker sequences.

There are at least two possible ways to break the correlations we have seen in the Kronecker sequence. One is by increasing the difference between the bases for any pair of dimensions; the other is to select better basis for the Kronecker sequence. It became clear that the smaller the partial quotients in the continued fractions of the irrational number, the more uniformly distributed the s -dimensional Kronecker sequence is [4]. We blend those two methods and propose a novel approach by using generalized golden ratio, which is widely applied in many applications in computer science [6]. The 2-D projections derived from our new algorithm are shown in Figure 2. The patterns appeared in Figure 1 disappear in Figure 2.

We have demonstrated that generalization of golden ratio is the simplest and most effective method to break this correlation in original Kronecker sequences. It is also very easy to implement the original Kronecker sequences and generate high-quality of the optimal Kronecker sequences.

In Bratley et al's paper [1], the Faure and Sobol sequences are used to evaluate high dimensional integrals, and the errors in the numerical results for over 30 dimensions become quite large. When we compared their results with our optimal Kronecker sequences, we found that our method has much smaller errors comparing to the




Figure 2: Left: 1000 points of dimension 1 and 2 projection for optimal Kronecker sequence; right: 1000 points of dimension 26 and 27 projection for optimal Kronecker sequence.

benchmark Faure and Sobol methods.

2.2 High order numerical method for nonlinear filter problems

Assume that $X = X_t$ is a signal process and $Y = Y_t$ is the observation. Because of noises, X and Y are solutions of the following stochastic differential equations.

$$\begin{cases} dX_t &= b(X_t)dt + \sigma(X_t)dU_t \\ dY_t &= h(X_t)dt + dV_t \end{cases}$$

Here U_t, V_t are independent Brownian motions which reflect the noises of the signal and the observation. The filter problem is to seek the best estimation of $\phi(X)$ given the observation Y . Mathematically this amounts to seek a conditional expectation $E(\phi(X_t) | (Y_s, 0 \leq s \leq t))$, i.e.,

$$E(\phi(X_t) | (Y_s, 0 \leq s \leq t)) = \underset{\{Z_t \text{ is } Y_{(0,t)} \text{ measurable}\}}{\text{Argmin}} \quad (E|\phi(X_t) - Z_t|^2)$$

According to Zaikai filter theory, the best approximation is given by

$$E(\phi(X_t) | (Y_s, 0 \leq s \leq t)) = \frac{\int \phi(x)p(t, x)dx}{\int p(t, x)dx}.$$




Figure 3: Sample path of the ship location




Figure 4: Un-normalized probability distribution of the ship location

Also according to the Zakai's theory, p satisfies the following stochastic partial differential equation.

$$dp_t = \left(\sum_{i,j=1}^d \sigma \sigma^* \frac{\partial^2 p_t}{\partial x_i \partial x_j} + \sum_{i=1}^p b_i \frac{\partial p_t}{\partial x_i} \right) dt + p_t h^* dY_t.$$

So the nonlinear filter problem is solved as long as the above stochastic partial differential equation is solved. We have derived a higher order method to solve the nonlinear stochastic differential equations. We achieve the high order method through solving a backward forward doubly stochastic differential equations. Detail of the method can be found in [3].

We then applied the method to solve a problem of tracking a ship in open water through the angle observation of the ship. Figure 3 shows a sample path of the ship and Figure 4 shows the probably distributions of the ship location at four different times.

2.3 Error analysis of finite element approximations for stochastic Stokes equations

In this subproject, we consider the steady-state Stokes equation with the forcing term perturbed by white noise:

$$\left\{ \begin{array}{l} -\nu \Delta u(x, \xi) + \nabla p(x, \xi) = f + \sigma(x) \dot{W}(x, \xi) \quad x \in D \subset R^d (d = 2, 3), \quad \xi \in \Omega, \\ \operatorname{div} u = 0 \quad \text{in } D, \\ u = 0, \quad \text{on } \partial D. \end{array} \right. \quad (1)$$

Here $(\Omega, \mathcal{F}, P)$ is a probability space and $\dot{W} = (\dot{W}^1, \dots, \dot{W}^d)$ is the white noise such that

$$E(\dot{W}^j(x) \dot{W}^j(x')) = \delta(x - x'), \quad x, x' \in \Omega, \quad j = 1, \dots, d.$$

Our strategy of solving this problem consists of two steps. In the first step, we construct an approximate white noise $\dot{W}_h$ based the finite element partition of the physical domain D . In the second step, we construct a finite element approximation (u_h, p_h) for the solution (u, p) of (1). Under certain assumptions on the regularity of the domain and using the Taylor-Hood finite element, we proved the following error estimates [2].

$$\left\{ \begin{array}{l} E(\|u - u_h\|^2 + \|p - p_h\|_{-1}^2) \leq C |\ln h| h^2 \quad d = 2, \\ E(\|u - u_h\|^2 + \|p - p_h\|_{-1}^2) \leq Ch \quad d = 3. \end{array} \right. \quad (2)$$

The significance of the error estimate is twofold:

- It indicates that the due to the lack of regularity, the convergence order of the finite element solution for stochastic Stokes equations is much lower than that for deterministic Stokes equations.
- It provides a guidance for the number of samples used in Monte Carlo simulations to evaluate the statistics of the solutions.

We also constructed practical numerical algorithms to find the finite element approximations using the Monte Carlo method and verified our theoretical results through

numerical experiments ([2]).

2.4 Fast and efficient numerical algorithms for stochastic partial differential equations

We first developed a fast algorithm of high dimension orthogonal polynomial expansions on sparse grids. Then we apply this algorithm to solve stochastic partial differential equations (SPDEs). This is achieved through the following four steps.

- **Step 1.** Use only sparse indices in the orthogonal expansions. For a d dimensional problem, this reduces the number of terms in the orthogonal expansion from $O(n^d)$ to $O(n \log^{d-1} n)$, a tremendous saving of computing cost.
- **Step 2.** Use fast Fourier transform as well as sparse grid to compute the coefficients of the orthogonal expansion. This reduces the computing cost from $O(n^d)$ to $O(n \log^{(d+1)n})$, yet another significant reduction of computing complexity.
- **Step 3.** Apply the fast orthogonal expansion algorithm to solve stochastic partial differential equations with random coefficients and forcing term using the spectral method. The remarkable feature of our algorithm is that the coefficient matrix is sparse and each element can be calculated analytically.

Our numerical experiments show that the overall computing complexity of solving the stochastic partial differential equation is in the same magnitude as computing the orthogonal expansion of the forcing term, which is proved to be $O(n \log^{d+1} n)$. In comparison, the complexity of conventional spectral method is at least $O(n^d)$, which is prohibitively expensive when the dimensional d is large as in the case of solving stochastic partial differential equations. The numerical verification of the computational complexity is shown in Figure 2.4.




Figure 5: The CPU times in numerical experiment v.s. theoretical estimates Solid lines: $O(n \log^{d+1} n)$, $d = 4, 6, 8$ Dashed lines: CPU times in numerical experiment

References

- [1] P. Bratley and B. Fox, Algorithm 659: Implementing Sobol's quasirandom sequence generator, *ACM Trans. Math. Softw.*, Vol. 14, 88–100, (1988).
- [2] Yanzhao Cao, Z. Chen and Max Gunzburger, Finite element method approximation and error estimates for stochastic Stokes equations, *Advances in Computational Mathematics*. Vol. 33, 215–230, (2010).
- [3] Feng Bao, Yanzhao Cao and Weidong Zhao, Numerical solutions for forward backward doubly stochastic differential equations and Zakai equations, *International Journal of Uncertainty Quantification*, DOI: 10.1615, (2011).
- [4] J. Goodman and J. O'Rourke, Handbook of discrete and computational geometry, *CRC Press, Inc.*, (2004).
- [5] S. Joe and F. Kuo, Constructing Sobol Sequences with Better Two-Dimensional Projections, *SIAM J. Sci. Comput.*, Vol. 30, 2635–2654, (2008).

- [6] S. Sen and R. Agarwal, Golden ratio in science, as random sequence source, its computation and beyond, , Vol. 56, Comput. Math. Appl. 469–498, (2008).
- [7] X. Wang and K.T. Fang, The Effective dimension and quasi-Monte Carlo, ***Journal of Complexity***, Vol. 19, 101-124, (2003).
- [8] B. Vandewoestyne, R. Cools and T. Warnock, On obtaining quadratic and cubic error convergence using weighted Kronecker-sequences, ***Computing***, Vol. 80, 75–94, (2007).

3 Personnel Supported During Duration of Grant

Hongmei Chi, PI

Yanhzao Cao, Co-PI

Kevin Lawrence, Graduate Student, Florida A&M University, Tallahassee, Florida

Trenton Johnson, Undergraduate Student, Florida A&M University, Tallahassee, Florida

Reginald McGee, Undergraduate student, Florida A&M University, currently Ph. D. student in Mathematics at Purdue University

Haiyan Tian, Visiting Scholar, University of Southern Mississippi

Bin Wu, Visiting Scholar, Zhongshan University, China

Ying Jiang, Visiting Scholar, Zhongshan University, China

4 Publications

- 1 Feng Bao, Yanzhao Cao and Weidong Zhao, Numerical solutions for forward backward doubly stochastic differential equations and and Zakai equations, ***International Journal of Uncertainty Quantification***, DOI: 10.1615, (2011).

- 2 Yanzhao Cao and Li Yin, Spectral method for nonlinear stochastic partial differential equations of elliptic type, accepted by *Numer. Math. Theo. Meth. App.*, Vol. 4 38–52, (2011).
- 3 Yanzhao Cao, Z. Chen and Max Gunzburger, Finite element method approximation and error estimates for stochastic Stokes equations, *Advances in Computational Mathematics*. Vol. 33, 215–230, (2010).
- 4 B. Vandewoestyne and H. Chi and R. Cools, An empirical Investigation of Different Scrambling Methods for Faure Sequences, *Mathematics and Computers in Simulation* , Vol. 81, 522–535 (2010).
- 5 H. Chi and Y. Cao, Generating Parallel Random Sequences via Parameterizing EICGs For Heterogeneous Computing Environments, *ICCSA 2010, March 23- 26, Fukuoka, Japan*.
- 6 Yanzhao Cao, Bin Wu and Yuesheng Xu, A fast collocation method for solving stochastic integral equations, *SIAM J. Numer. Anal.* Vol. 47(5), 3744 – 3767, (2009).
- 7 Yanzhao Cao, Z. Chen and Max Gunzburger, An ANOVA analysis for a class of partial differential equations with uncertain boundary conditions, , *Int. J. Numer. Anal. & Model.*, Vol. 6, 256-273, (2009).
- 8 H. Chi, Parallel Quasirandom Number Generations for Heterogeneous Computing Environments, *International Journal of Parallel, Emergent and Distributed Systems*, Vol. 24, 21–29, (2009).
- 9 Yanzhao Cao, H. Chi, C. Milton and W. Zhao, Exploitation of sensitivity derivatives on quasi Monte Carlo simulations, *Monte Carlo Methods and Applications*, Vol. 14, 269–279, (2008).
- 10 Yanzhao Cao, Ran Zhang and Kai Zhang, Finite Element and Discontinuous Galerkin Method for Stochastic Helmholtz Equation in Two- and Three-Dimensions. *J. Comp. Math.*, Vol. 26, 702–715, (2008).

- 11 Yanzhao Cao, K. Zhang and R. Zhang, Finite Element Method and Discontinuous Galerkin Method for Stochastic Scattering Problem of Helmholtz Type, *Potential Analysis*, Vol. 28, 301–319, (2008).