

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.					
1. REPORT DATE (DD-MM-YYYY) 30-09-2011		2. REPORT TYPE Final Report		3. DATES COVERED (From - To) Dec 2008-June 2011	
4. TITLE AND SUBTITLE Solving Differential Equations with Random Ultra-Sparse Numerical Discretizations				5a. CONTRACT NUMBER FA-9550-09-0403	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) David M. Bortz Andrew J. Christlieb				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Colorado - Boulder				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Office of Scientific Research Computational Mathematics Program 875 N. Randolph Ste. 325, Rm 3112 Arlington, VA 22203				10. SPONSOR/MONITOR'S ACRONYM(S) AFOSR	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL - OSR - VA - TR - 2012 - 0547	
12. DISTRIBUTION/AVAILABILITY STATEMENT A					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We proposed a novel approach which employs random sampling to generate an accurate non-uniform mesh for numerically solving Partial Differential Equation Boundary Value Problems (PDE-BVPs). From a uniform probability distribution U over a 1D domain, we considered a M discretization of size N where $M \gg N$. The statistical moments of the solutions to a given BVP on each of the M ultra-sparse meshes provide insight into identifying highly accurate non-uniform meshes. We used the pointwise mean and variance of the coarse-grid solutions to construct a mapping $Q(x)$ from uniformly to non-uniformly spaced mesh-points. The error convergence properties of the approximate solution to the PDE-BVP on the non-uniform mesh are superior to a uniform mesh for a certain class of BVPs. In particular, the method works well for BVPs with locally non-smooth solutions. We fully developed a framework for studying the sampled sparse-mesh solutions and provided numerical evidence for the utility of this approach as applied to a set of example BVPs.					
15. SUBJECT TERMS Numerical Partial Differential Equations - Non-uniform Mesh - Random Sampling					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code)

Solving Differential Equations with Random Ultra-Sparse Numerical Discretizations

D. M. Bortz¹ and A. J. Christlieb²

1) Department of Applied Mathematics, University of Colorado

2) Department of Mathematics, Michigan State University

Email: dmbortz@colorado.edu, christlieb@math.msu.edu

Final Report

Grant #: FA9550-09-1-0403

David Bortz

Department of Applied Mathematics
University of Colorado, Boulder

Abstract We proposed a novel approach which employs random sampling to generate an accurate non-uniform mesh for numerically solving Partial Differential Equation Boundary Value Problems (PDE-BVP's). From a uniform probability distribution $\mathcal{U}$ over a 1D domain, we considered M discretizations of size N where $M \gg N$. The statistical moments of the solutions to a given BVP on each of the M ultra-sparse meshes provide insight into identifying highly accurate non-uniform meshes. We used the pointwise mean and variance of the coarse-grid solutions to construct a mapping $Q(x)$ from uniformly to non-uniformly spaced mesh-points. The error convergence properties of the approximate solution to the PDE-BVP on the non-uniform mesh are superior to a uniform mesh for a certain class of BVP's. In particular, the method works well for BVP's with locally non-smooth solutions. We fully developed a framework for studying the sampled sparse-mesh solutions and provided numerical evidence for the utility of this approach as applied to a set of example BVP's.

Summary Over the duration of this grant, while developing our SMRT methodology for solving BVP-PDEs, the core of our research efforts have include the following: substantial refinement to our algorithm, extension of the algorithm to higher dimensions, and establishing the theoretical well-posedness of our approach [3,4]. All of these topics are linked by a desire to efficiently exploit the high parallellizability of our approach and future implementation on massively parallel multi-core technologies. Lastly, we have also been invited to contribute a review article on computing on GPU's to SIAM Review [5]. The focus of this effort is one type of computation which is substantially accelerated on GPU's.

We now give a brief summary of our progress.

Scandalously Parallellizable Mesh Generation The PI and his collaborator are developing an SMRT framework to generate non-uniform meshes for solving PDE's [3,4,5]. These discretizations can offer superior solution accuracy and convergence properties to that of uniform spacing. We offer a brief overview of our proposed algorithm as well as the establishment of a preliminary theoretical framework [3]. Also, in [4] we extended results in [2] to the identification of Q using an optimization technique using results from probability theory. However, we discovered that the approximation technique described below was substantially more efficient.

We consider a monotonically non-decreasing function $Q : \bar{\mathbf{I}} \rightarrow \bar{\mathbf{I}}$ which is absolutely continuous on a finite number of compact subsets of $\bar{\mathbf{I}}$ and restricted at the endpoints to $Q(0) = 0$, $Q(1) = 1$. The purpose of the function Q is to map the uniformly spaced mesh to a non-uniformly spaced one. The goal is to develop a strategy for identifying a Q such that, e.g., the approximate solution to the Poisson problem

$$u''(Q(x)) = f(Q(x)) \text{ s.t. } u(Q(0)) = A; \quad u(Q(1)) = B,$$

has convergence properties (in n) superior to a uniform spacing. The core of our approach is to identify Q via a sparse stochastic approximation. We repeatedly sample from a distribution P and then use pointwise statistical moments of the coarse solutions to generate the desired non-uniform mesh function Q . Naturally, different classes of problems call for different strategies for generating Q . Our results, however, suggest that a more generalizable strategy may exist. Before presenting our conclusions, we briefly establish some notation.

Let p be a function taking a point $\xi \in \bar{\mathbf{I}}$ and a random vector of length n , and mapping them to a single random variable

$$p(\xi, \mathbb{X}_{(n)}(P)) \equiv \mathbb{E}_K \left[\{U(\mathbb{X}_{(n)}(P))\}_{K=k} | X_{(k)} = \xi \right]. \quad (1)$$

The function U takes a discretization of the domain and solves the BVP. The operator $\mathbb{E}_K$ denotes expectation with respect to a uniform distribution on $\{1, \dots, n\}$ where the distribution of the index random variable K and $\{\cdot\}_K$ denotes the K th element of a vector. We note that this function returns a random variable for each ξ . Let the pointwise mean of p be defined for $\xi \in \bar{\mathbf{I}}$ as

$$\mu(\xi) \equiv \mathbb{E}_P \left[\mathbb{E}_K \left[\{U(\mathbb{X}_{(n)}(P))\}_{K=k} | X_{(k)} = \xi \right] \right]. \quad (2)$$

The pointwise variance of p is defined for $\xi \in \bar{\mathbf{I}}$ as

$$v(\xi) \equiv \mathbb{V}_P \left[\mathbb{E}_K \left[\{U(\mathbb{X}_{(n)}(P))\}_{K=k} | X_{(k)} = \xi \right] \right], \quad (3)$$

where $\mathbb{V}_P$ denotes variance with respect to P , $\mathbb{E}_K$ denotes expectation with respect to $\mathcal{U}\{1, \dots, n\}$, the distribution of the index random variable K , and $\{\cdot\}_K$ denotes the K th element of a vector.

Answers to the critical questions for this approach are depicted below

For each candidate Q , how many sample sparse grids need to be generated? The relationship between the mesh size n and the number of samples m is non-trivial. and Figure 1 illustrates this by depicting the error in $\bar{v}$ (relative to $\bar{v}$ computed with $m = 3000$ sampled from a uniform distribution on $\bar{\mathbf{I}}$) for a range of n and m values. For a given n , though, we do note that the error in the $\bar{v}$ computation is decreasing. In Figure 2 we depict the number of samples of vector size n which are needed to ensure three digits of accuracy in estimating the variance. Since the number was consistently below 1000 over a range of n , we let $m = 15000$ in all subsequent simulations (unless otherwise specified).

In what way do the random solutions converge to the actual solution? For a conventional finite difference discretization, we would consider the error E in the solution

$$\begin{aligned} \|E(Q, \mathbf{x}_n^0)\| &= \|u(Q(\mathbf{x}_n^0)) - U(Q(\mathbf{x}_n^0))\| \\ &= \left\| A_{Q(\mathbf{x}_n^0)}^{-1} \left(A_{Q(\mathbf{x}_n^0)} u(Q(\mathbf{x}_n^0)) - f_{Q(\mathbf{x}_n^0)} \right) \right\| \\ &\leq \left\| A_{Q(\mathbf{x}_n^0)}^{-1} \right\| \left\| \tau_{Q(\mathbf{x}_n^0)} \right\|, \end{aligned}$$




Figure 1: $\log_{10}$ of the error in the computation of $\bar{v}$ (sampling from a uniform distribution on $\bar{\mathbf{I}}$) as a function of m and n . Note the general downward trend along both the m and n axes.




Figure 2: For each n , the vertical axis reflects the number of samples needed to compute the variance with 3 digits of accuracy relative to $\bar{v}$ (sampling from uniform distribution on $\bar{\mathbf{I}}$) with $m = 3000$.

which is bounded above by the spectral radius of the inverse of the finite difference operator $A_{Q(\mathbf{x}_n^0)}^{-1}$ and a truncation error $\tau_{Q(\mathbf{x}_n^0)}$. For the non-uniform three-point-stencil approximating the second derivative, the truncation error is $O(\max_k |h_k|)$. For our development, we consider a probabilistic version of this error, with the following conditions.

CONDITION C1. *For a given P , the spectrum of $A_{\mathbb{X}_{(n)}(P)}^{-1}$ is bounded in $[0, 1]$.*

CONDITION C2. *For a given P , the truncation error induced by a finite difference approximation to the second derivative is first order in the largest step-size h .*

THEOREM 1. Under Condition C1 and C2, the expected error converges pointwise to zero.

See [3] for support of these conditions as well as a proof of the theorem.

How should Q be constructed? The function Q is created using the statistical moments of the sampled sparse-mesh solutions and based on results in [1]. For the problems with second derivatives we define Q as

$$Q(x) = \left[\frac{q_1(\cdot)}{q_1(1)} \right]^{-1}(x),$$

where

$$q_1(x) = \int_0^x \sqrt{|\mu'(\xi; U(\mathbb{X}_{(n)}(P)))|} d\xi,$$

and the superscript -1 is an inverse function operator. Essentially, this definition will pile up points in regions with a steep solution in an effort to provide higher order accuracy for the nonuniform second derivative discretization.

For the problem with a second power of the first derivative, we define Q as

$$Q(x) = \left[\frac{q_2(\cdot)}{q_2(1)} \right]^{-1}(x),$$

where

$$q_2(x) = \int_0^x \mu''(\xi; U(\mathbb{X}_{(n)}(P)))^2 v(\xi; U(\mathbb{X}_{(n)}(P)))^3 d\xi,$$

and v is defined above. Evidence for improvement in error convergence is depicted in Figures 3-4.

We hypothesize that the reason $q_1(x)$ works well is that the μ' may converge faster than μ . We also hypothesize that the function $q_2(x)$ works well because the second derivative (when cast as the local curvature) is inversely proportional to the local variance of a random variable (a result which is well known in the semi-parametric nonlinear regression literature). Essentially, while the μ'' may not converge quickly, the product $\mu''v$ does. We also found that multiplication by an extra v dramatically improves the computed Q , though an explanation is not immediately clear. A deeper understanding of the spectrum of $A_{\mathbb{X}_{(n)}(P)}$ and how it depends upon the choice of P will be essential to explaining the efficiency of $q_2(x)$. We plan to explore both of these issues in a future paper [4].

Acknowledgment/Disclaimer This work was sponsored (in part) by the Air Force Office of Scientific Research, USAF, under grant/contract number FA9550-09-1-0403. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily




Figure 3: Error convergence for uniformly and non-uniformly spaced points for the steady-state Hamilton-Jacobi BVP.




Figure 4: Error convergence of the different mesh mappings for the singular BVP.

representing the official policies or endorsements, either expressed or implied, of the Air Force Office of Scientific Research or the U.S. Government.

References

1. D.M. Bortz and P.W. Nelson, "Model Selection and Mixed-Effects Modeling of HIV Infection Dynamics," Bull. Math. Biol., (2006) 68:2005-2025.
2. H.T. Banks and D.M. Bortz, "Inverse problems for a class of measure dependent dynamical systems," J. Inv. Ill-Posed Problems, (2005) 13:103-121.

Publications

3. D.M. Bortz and A.J. Christlieb, "Scandalously Parallelizable Mesh Generation," in revision SIAM J. Scientific Computation.
4. D.M. Bortz and E.C. Byrne, "Identification of Conditional Probability Measures," in revision Inverse Problems.
5. D.M. Bortz and A.J. Christlieb, "Analysis of Random Mesh Generation Methods," in preparation.
6. D.M. Bortz, A.J. Christlieb, J. Cohen, and F. Fahroo, "Computations on GPU's," in preparation.

Personnel Supported by Grant During 10-11

D.M. Bortz, Assistant Professor, University of Colorado, Boulder

A. J. Christlieb, Associate Professor, Michigan State University

Honors & Awards Received

University of Graz, Invited speaker for summer school on mathematical methods in mathematical biology 2009

University of Colorado, Junior Faculty Award 2008

University of Michigan, Horace H. Rackham Faculty Fellowship 2003

Changes in research objectives, if any: None.

Change in AFOSR program manager, if any: None

Extensions granted: No Cost Extension was granted till 6/30/11 due to initial nine month delay in receiving funds.

New Discoveries None.