

ASSESSING THE TOXICITY AND BIOAVAILABILITY OF 2,4-DINITROANISOLE IN ACUTE AND SUB-CHRONIC EXPOSURES USING THE EARTHWORM, *EISENIA FETIDA*


Jessica G. Coleman

Research Biologist

US Army Engineer Research and
Development Center
Vicksburg, MS



US Army Corps of Engineers
BUILDING STRONG



2,4 DNAN



Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE JUN 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Assessing the Toxicity and Bioavailability of 2,4-Dinitroanisole in Acute and Sub-Chronic Exposures Using the Earthworm, Eisenia Fetida				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Army Engineer Research and Development Center,Environmental Laboratory,3909 Halls Ferry Road Waterways Experiment Station,Vicksburg,MS,39180-6199				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the NDIA Environment, Energy Security & Sustainability (E2S2) Symposium & Exhibition held 14-17 June 2010 in Denver, CO.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 24	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			


Project Team

- Sandra M. Brasfield- Research Biologist
- Frances C. Hill- Research Chemist
- Choo Y. Ang- Cell Biologist
- Jeffery A. Steevens- Research Toxicologist
- Ganna Gry'nova- Student, Jackson State University
- Robert Boyd- Research Assistant


This work was supported by the Army's Environmental
Quality Technology Basic Research Program



Overview



DNAN




TNT



- The Army currently produces and employs 2,4,6-trinitrotoluene (TNT) in traditional munitions, but is facing a mandate to replace TNT with less sensitive and less toxic compounds
- 2,4-dinitroanisole (DNAN) is currently being tested by the DOD
- Hypothesis: DNAN will exhibit a similar mechanism of action, but reduced toxicity relative to the closely related compounds, DNT and DNP
- Preliminary results obtained through acute and sub-chronic terrestrial studies suggest that DNAN is **less toxic** relative to TNT




2,4-Dinitroanisole



- Not a new entergetic → use in Amatol 40 (50% DNAN, 35% ammonium nitrate, 15% RDX) in the warhead of some V-1 flying bombs during World War II
- Gained popularity because of scarcity of TNT
- 10% reduction in explosive impact
- Little information exists on the environmental risk of DNAN, driving a need for further research in this area



Assessing the Environmental Toxicity of DNAN



Dermal
Exposure

Ingestion



Whole
Organism


Tissue

Cell

- Terrestrial ecosystems likely impacted due to munitions deployment over land
- Chose the earthworm *Eisenia fetida* as the terrestrial receptor
- Investigated multiple acute and sub-chronic exposure pathways:
 - ▶ Acute 72 hr dermal toxicity studies
 - ▶ Acute 7 day range finder soil studies
 - ▶ Sub-chronic 28 day soil studies
- Cellular toxicity endpoints coupled with exposures through *in vitro* Neutral Red Based Toxicology Assay (NRRT)



Dermal Filter Paper Exposure




- OECD Guideline 207 (1984)
- Duration: 72 Hours, n=10
- Treatments: DNAN and TNT dissolved in 100% methanol, 1ml spiked onto filter paper: 0, 0.0001, 0.001, 0.01, 0.1 mg/cm²
- 1 earthworm per vial
- NRRT Analysis conducted from n=5 per treatment

**Toxicity/
Bioaccumulation**


**Cellular Stress
NRRT**



Dermal Toxicity of DNAN/TNT




DNAN/TNT Bioaccumulation




Cellular Stress/NRRT

Two Methods of Measurement

1)



Extract 50 ul from clitellum



2)



Expose worm to GGE t= 2mins
collect coelomocyte solution




1 row per worm/treatment,
obtain measurements through
spectrophotometer


- NRRT is a biomarker of cellular stress; viable cells stain red with dye
- Coelomic fluid extracted and analyzed in two different methods, 2nd method chosen for final study:
 - ▶ Coelomocytes collected in 2 ml Guaiacol Glyceryl Ether (GGE) solution, centrifuged, decanted, washed with 1X Phosphate Buffered Saline (PBS) and re-suspended in 1,500 ul PBS
 - ▶ Kinetic readings obtained from spectrophotometer every 5 minutes for 1 hour to determine neutral red retention time



Comparison of Total Cells and NRRT



DNAN & TNT Toxicity in Human Liver HEPG2 Cells



Soil Exposures




Toxicity & Body Burdens



Reproductive Toxicity



NRRT




- Range Finder-(n=3, 10 worms per rep) determine lethal and non-lethal range of DNAN
- Sub-chronic 28 day studies - (n=3, 10 worms per rep) comparatively assessed DNAN and TNT toxicity, bioaccumulation, and cellular stress through NRRT analysis
- Soil studies provide multiple routes of exposure through dermal contact and ingestion



Materials and Methods: Soil Prep

- Field soil spiked with increasing concentrations of DNAN and TNT dissolved in methanol
 - ▶ Range Finder Concentration: 0-300 mg/kg
 - ▶ 28 Day Exposure Concentrations: 0-100 mg/kg
- Tumbled overnight, placed in test containers, hydrated to 85% WHC




Materials and Methods: Test Conditions



**Earthworms
depurated 24-
hours
Adults 0.3-0.6 g**



**10 added
per
treatment
n=3**





**Test conducted at 22°C, 80% humidity,
continuous light**

Endpoints : bioaccumulation, toxicity, growth, reproduction, NRRT



Results: DNAN Acute Range Finder




n=3, 10 per rep
T=7 days



BUILDING STRONG®


Sub Chronic 28-Day Exposure



TNT and DNAN tested concentrations: 0, 12.5, 25, 50, 100 mg/kg n=3, 10 per rep




Sub-Chronic Preliminary Results




Results: 28 Day Exposure NRRT

DNAN




TNT



BUILDING STRONG®

Computational Chemistry and Toxicity of DNAN

First stage of alkaline hydrolysis of 2,4-Dinitroanisole





- Initial exposures conducted with DNAN parent compound

- Additional exposures to be conducted with DNAN degradation product predicted by computational chemistry models


- TS (Transition State)
- MC (anionic Meisenheimer complexes)
- DS (direct substitution of nitro group w/phenol)




Kinetics simulations



First stage of alkaline hydrolysis of 2,4-Dinitroanisole




Electrostatic Potential (ESP) maps for selected transition states



Second stage of alkaline hydrolysis of 2,4-Dinitroanisole

We have considered attack of second hydroxyl anion on nitro phenol DS₂, formed on the first stage as a result of direct substitution.




Transition states geometries				Products geometries			
TS	Bond C-O, Å	Bond C-R (R=H,CH ₃), Å	□OCR	TS	Bond C-O, Å	Bond C-R (R=H,CH ₃), Å	□OCR
TS_21	2.096	1.380	95.584	MC_21	1.485	1.465	103.631
TS_23	2.012	1.085	88.177	MC_23	1.514	1.099	105.804
TS_24	1.748	1.605	93.154	DS_24	1.395	-	-

** For bonds: first column is bond length in Å, second column is Wiberg index of bond order*


Reaction thermo chemistry									
Activation parameters					Products parameters				
Reaction	ΔE, kcal/mol	ΔH, kcal/mol	ΔG, kcal/mol	ΔS, cal/mol*K	Product	ΔE, kcal/mol	ΔH, kcal/mol	ΔG, kcal/mol	ΔS, cal/mol*K
TS_21	13.74	13.53	20.79	-24.38	MC_21	6.08	5.84	13.06	-24.24
TS_23	15.65	15.49	22.47	-23.42	MC_23	11.46	11.18	18.47	-24.43
TS_24	29.48	29.37	36.12	-22.66	DS_24	-30.31	-30.78	-35.25	16.97

Kinetics simulations



2,4-dihydroxytoluene is highly dominating product because of low stability of competitive pathways adducts. However, its formation requires enormous amount of time.

Future Direction of Project




Determine toxicity of predicted breakdown product in terrestrial systems



Overview: Results/Findings

- DNAN resulted in lower toxicity to relative to TNT compounds in *E. fetida* exposures
- Two methods tested for NRRT analysis; kinetic readings obtained from spectrophotometer found to be less variable
- Better definition of link between computational chemistry and toxicology as it relates to munitions
- Information previously unavailable on the bioavailability and toxicity of DNAN to terrestrial organisms
- Data obtained from our exposures will benefit the Army by reducing uncertainty about the environmental effects of DNAN; therefore, reducing cost associated with unnecessary site remediation.





US Army Engineer Environmental
Laboratory, Vicksburg, MS,

Contact Information:

Jessica.g.coleman@us.army.mil

601-634-3976



BUILDING STRONG®