

Multi-Scale Model-Driven Sampling with Autonomous Systems at a National Littoral Laboratory: Turbulence Characterization from an AUV

Edward R. Levine

Naval Undersea Warfare Center, Division Newport

Code 8211

Newport, RI 02841-1708

phone: (401) 832-4772 fax: (401) 832-2146

email: levineer@tech.npt.nuwc.navy.mil

Document #: N0001498WX30325, N0001498WX30334, N0001498WX30417

<http://marine.rutgers.edu/MAS>

LONG-TERM GOAL

My long term goal is to utilize turbulence measurements obtained from small Autonomous Underwater Vehicle (AUV) based sensors as the subgrid characterization tool in combined coastal ocean observation/prediction networks.

OBJECTIVES

I wish to use AUV-based turbulence measurements to quantify mixing in shallow water physical process studies (upwelling regions, fronts, boundary layers) within the context of the LEO-15 based National Ocean Partnership Program (NOPP) coupled ocean observation/modeling system. This includes estimating mixing levels, identifying regions of enhanced mixing, determining the horizontal spatial scale of mixing events, defining the role of boundary layers, and parameterizing results for coastal predictive model testing studies of subgrid scale processes.

APPROACH

My approach is to integrate an optimum turbulence sensor suite into a small, logistically simple, AUV, with input from the ocean turbulence and modeling communities. Then, I seek to establish this small AUV as a viable platform for coastal turbulence research. Towards this end, I obtain horizontal profiles of dissipation rate, temperature microstructure, 3-dimensional small scale velocity, larger scale vertical shear of horizontal current, and stratification in the coastal environment.

Subsequently, I studied mixing in the context of the multi-scale measurements surrounding the NOPP LEO-15 node site in the Mid-Atlantic Bight. I sampled adaptively using the Rutgers University continental shelf model SCRUM (Song and Haidvogel, 1994).

The sensors provided data for estimates of eddy diffusivity profile (Gargett and Moum (1995), eddy viscosity profile (using the truncated TKE equation), bulk and gradient Richardson numbers, fluxes [using the correlation technique]. These data enable us to evaluate turbulence closure schemes associated with subgrid mixing processes as parameterized by 3 different surface boundary layer submodels in SCRUM, the coastal circulation model.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Multi-Scale Model-Driven Sampling with Autonomous Systems at a National Littoral Laboratory: Turbulence Characterization from an AUV				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Undersea Warfare Center,Division Newport ,Code 8211,Newport,RI,02841				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

WORK COMPLETED

A turbulence sensor package was electrically and mechanically integrated into the REMUS AUV (Levine and Lueck, 1998). Sensors include two shear probes, an ultra-fast thermistor, an upward and downward looking ADCP, two CTDs, and an ADV-O. Preliminary field trials were conducted in Narragansett Bay to test sensor/platform synergy and software viability.

Utilizing these techniques, scientific studies of the Ocean Boundary Layer (OBL) during an upwelling event were conducted near the Rutgers University LEO-15 site on the inner continental shelf off New Jersey during July 1998. In the field experiment, high quality data were obtained from all sensors, and data analysis is proceeding well.

RESULTS

Using model-based adaptive sampling, the AUV was deployed along a trajectory through components of the Leo-15 upwelling region, including the upwelling gyre and offshore jet. These measurements were made synoptically with those from other platforms which characterize larger scale structures on the nearby continental shelf. Model predictions which include assimilated data from the wide variety of sampling platforms is also available for intercomparison .

Results indicate that the modified REMUS AUV (Fig 1) was a viable platform for turbulence data acquisition in the coastal ocean. For example, a data sample obtained from the vertical and transverse shear probes, for a short transect in the upwelling center during peak upwelling on July 21 1998, has been analyzed. These data have been processed to remove noise associated with vehicle vibrations. This process is done using data from accelerometers located in the probe pressure case directly behind the probe mounts, utilizing the techniques of Levine and Lueck (1998). Also, the data have been low pass filtered. The autospectra for this segment are shown in Figure 2, and correspond to a dissipation rate of approximately 5×10^{-8} W/kg based on the computed spectral variance over the wavenumber band of the filtered data. The Nasmyth “universal spectrum” (Oakey, 1982) for this dissipation rate and appropriate viscosity agrees well in level and shape with the observations out to wavenumbers close to the physical size of the sensing tip of the shear probes.

IMPACT/APPLICATION

The AUV-based turbulence measurements provide a unique horizontal profiling view of the variability of the mixing environment that cannot be obtained by more conventionally sampling measurements, and this approach can be further exploited in yo-yoed horizontal sections. These techniques will be invaluable in upwelling process studies in which competing turbulence closure model alternatives are testing in SCRUM to parameterize subgrid processes. Competing OBL alternatives include those of Mellor-Yamada (1974) Large et al. (1994), and Price et al. (1986). Features such as the evolution of the upwelling front can be tested.

TRANSITIONS

Our AUV sensor technologies, hardware and software, are being considered for inclusion as tactical oceanography payloads for the Manta UUV Initiative, proposed for an upcoming Accelerated Research Initiative (ARI).

RELATED PROJECTS

1 My AUV-based turbulence measurement system is also being utilized in NOPP studies with the Harvard led LOOPS project. Measurements were also made in Cape Cod Bay in September 1998.

REFERENCES

A. E. Gargett and J. N. Moum. 1995. Mixing effects in tidal fronts: results from direct and indirect measurements of density flux. *J. Phys. Ocean.*, 25, 2583-2608.

W. G., Large, J. C. McWilliams, and S. C. Doney. 1994: Oceanic vertical mixing: a review and model with a non-local boundary parameterization. *Rev. Geophys.*, 32, 363-403.

Levine, E. R., R. G. Lueck, 1998. Turbulence measurements with an autonomous underwater vehicle. *Journal of Atmospheric and Oceanic Technology*, Special Issue on Ocean Turbulence Measurement, in press.

G. L. Mellor and T. Yamada. 1974. A hierarchy of turbulence closure models for planetary boundary layers. *J. Atmos. Sci.*, 31, 1791-1806.

N. S. Oakey. 1982. Determination of the rate of dissipation of turbulent energy from simultaneous temperature and velocity shear measurements. *J. Phys. Ocean.*, 12, 256-271.

J. F. Price., R. A. Weller, and R. Pinkel. 1986. Diurnal cycling: Observations and models of the upper ocean response to diurnal heating, cooling, and wind mixing. *J. Geophys. Res.*, 91, C7, 8411-8427.

Y. Song and D. Haidvogel. 1994. A semi-implicit ocean circulation model using a generalized topography following coordinate system. *J. Comput. Phys.*, 115, 228-244.

PUBLICATIONS

Levine, E. R., D. Connors, R. Shell, and R. Hanson, 1997: Autonomous Underwater Vehicle-Based Sampling of Estuarine Hydrography, *Journal of Atmospheric and Oceanic Technology*, 14, 6, 1444-1454.

Levine, E. R., R. G. Lueck, 1998: Turbulence measurements with an autonomous underwater vehicle. *Journal of Atmospheric and Oceanic Technology*, Special Issue on Ocean Turbulence Measurement, in press.

Glenn, S. M., D. B. Haidvogel, O. M. E. Scofield, C J. von Alt, and E. R. Levine, 1998: Coastal Predictive Skill Experiments. *Sea Tech.*, April 1998, 66-72.

Levine, E. R., R. G. Lueck, 1998b: A small AUV-based turbulence measurement system for NOPP combined coastal observation/prediction networks. *EOS, Trans. Am. Geophys. Un.*, 1998 Ocean Sciences meeting, San Diego, Ca, Feb, 1998.

Levine, E. R., 1998: Tactical oceanographic sampling from an UUV, National Defense Industrial Association Conference, San Diego, Feb 1998.

PATENTS

D. French, E. Levine, and R. Lueck. 1998. A rigid sting extension for ocean turbulence measurement from an unmanned underwater vehicle. Navy case #78733.




Fig. 1 The REMUS AUV instrumented with turbulence sensors




Fig. 2 A comparison of shear probe autospectra with the “Universal” spectrum