

# **Transfer of Contaminants Between the Water Column and Bottom Sediments: The Role of Deposit- and Suspension-Feeding Benthic Invertebrates**

Cheryl Ann Butman  
Applied Ocean Physics and Engineering Department  
Mailstop 12  
Woods Hole Oceanographic Institution  
Woods Hole, MA 02543  
phone: (508) 289-2442 fax: (508) 457-2194 e-mail: [cbutman@whoi.edu](mailto:cbutman@whoi.edu)

Robert A. Wheatcroft  
Oceanic and Atmospheric Sciences Department  
Oregon State University  
104 Ocean Administration Building  
Corvallis, OR 97331-5506  
phone: (541) 737-3891 fax: (541) 737-2064 e-mail: [raw@oce.orst.edu](mailto:raw@oce.orst.edu)  
Award #: N00014-93-1042

## **LONG-TERM GOAL**

The ultimate goals of this research are to determine the dominant mechanisms whereby and the rates at which fine particles (<0.05 mm) are incorporated into marine sediments. Particular focus is on developing a mechanistic understanding of the role that deposit- and suspension-feeding benthic invertebrates play in particle deposition and burial relative to the abiotic case.

## **OBJECTIVES**

The objectives of this project are to manipulate and test various combinations of hydrodynamic, sedimentological and biological parameters to determine the magnitude of: (1) net accumulation of fine particles in sandy and “reworked” (mixed grain size) sediments relative to muddy sediments, (2) biodeposition by suspension feeders relative to deposit feeders, and (3) vertical transport modes and rates of sediment transport within the bed by deposit feeders.

## **APPROACH**

Carefully tailored laboratory flume and still-water experiments are used to quantify effects of suspension- and deposit-feeding benthic invertebrates on the transport of fine-grained sediment to the seabed, and its subsequent burial and retention within the bed. Using organisms from several functional and taxonomic groups, the flume and still-water studies are designed to determine the magnitude of biological effects (e.g., biodeposition, vertical transport in the bed, net accumulation) on fine-particle movement relative to the abiotic case. A variety of techniques are used to visualize and quantify flow and sediment transport, including digital particle image velocimetry (DPIV), optical backscatter sensors (OBSs), x-radiography and deliberate particle tracers. During FY98, we also sampled intertidal sediments to determine the vertical distribution of fine particles relative to the density of a surface deposit feeding worm (*Enoplobranchus sanguineus*) that was shown to subduct fine sediments in the lab.

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>1998</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-1998 to 00-00-1998</b>	
4. TITLE AND SUBTITLE <b>Transfer of Contaminants Between the Water Column and Bottom Sediments: The Role of Deposit- and Suspension-Feeding Benthic Invertebrates</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Woods Hole Oceanographic Institution, Department of Applied Ocean Physics and Engineering, Woods Hole, MA, 02543</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM002252.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

Aspects of this research are being conducted by J. Steven Fries as part of his dissertation research in the MIT/WHOI Joint Program. The goal of Fries's research is to determine the influence of the viscous sublayer on the deposition of particles to sand boundaries. The work is being conducted in a large, racetrack flume, recently constructed in WHOI's Coastal Research Center, and the techniques involve the use of DPIV to visualize and quantify the flow field, an array of OBSs to quantify fine particle transport, and direct sampling to measure fine-particle deposition.

## **WORK COMPLETED**

During FY98, one manuscript (Fries et al., in press) was revised and accepted for publication. Steve Fries wrote and defended his dissertation research proposal, and began his thesis research. He also continued with the analysis of near-bed velocity field and bottom-stress measurements collected over and within animal-tube mimic arrays in flume flows.

New research conducted during FY98 focussed on improving understanding of the effects of a surface deposit-feeding polychaete on fine-particle subduction in intertidal sands. Our previous laboratory experiments with the terebellid polychaete, *Enoplobranchus sanguineus*, demonstrated that it can non-locally transport fine-grained sediment to depths of several centimeters in the seabed. A logical consequence of this activity is that areas supporting terebellid populations should have a higher inventory of fine-grained sediments than areas without terebellids. This species lives in sandy intertidal sediments where fine particles typically do not accumulate. Thus, significant subduction of fine particles by this species would have important sedimentological, ecological and applied implications.

## **RESULTS**

During FY98, we sampled the vertical distribution of fine sediments in an intertidal sand flat with patchily distributed populations of *E. sanguineus*. One hundred cores were taken at low tide in Fairhaven Harbor, MA. There were 50 cores, one meter apart, along two perpendicular transect lines. The lines were oriented parallel and perpendicular to shore. Two types of cores were taken at each sampling location, a large-diameter core for enumeration of *E. sanguineus* individuals, and a small-diameter for quantification of the percentage of fine (<63  $\mu\text{m}$ ) sediments. The sediment cores were vertically sectioned into three depth horizons: 0-2, 2-6, and 6-10 cm. There was no significant trend in the percentage of fine sediments as a function of worm density for the upper two depth horizons, but for the 6-10 cm fraction the percentage of fine material increased with increasing worm density. Thus, these results support the *a priori* prediction that *E. sanguineus* subducts fine particles in sandy sediments.

As part of Fries' dissertation research, the fallout of fine suspended material was observed over boundaries of fixed sand roughness in a series of flume experiments. During these runs, particles collected in particle streaks usually associated with viscous sublayer flow structures over hydraulically smooth boundaries. The loss of material from suspension violated the anticipated balance between resuspension induced by bed shear stress and gravitational settling. This effective trapping of particles within the viscous sublayer will be studied in considerable detail.

## **IMPACT/IMPLICATIONS**

In recent times, increasingly stricter regulations have been imposed on the discharge of toxic chemicals to harbors in order to improve water quality. Many of these chemicals adhere to and thus transport as fine particles. In determining sites and rates of toxic chemical accumulation, depositional (muddy) regions have had the greatest focus of attention because fine particles are not expected to be deposited or retained in erosion (sandy) locales. This research suggests, however, that bioturbating activities of sand-dwelling organisms can affect significantly the retention of fine particles in sands. Moreover, because horizontal fluid flux is higher in erosional as compared to depositional sites, sandy areas potentially are exposed to more (toxic-laden) mass per unit time than muddy areas.

## **TRANSITIONS**

This research suggests that fine-particle accumulation should be studied in erosional, as well as depositional areas in order to understand the processes responsible for contaminant transport and retention in harbors associated with highly urbanized and industrialized cities.

## **RELATED PROJECTS**

This project is part of the Harbor Processes research program and is related to other projects within this program that address biological effects on sediments and sediment transport (e.g., Linda Shafner, et al.).

## **REFERENCES**

Fries, J.S., Butman, C.A. and Wheatcroft, R.A. in press: Subcritical ripple formation induced by biogenic mounds, *Marine Geology*.

## **PUBLICATIONS**

Fries, J.S., Butman, C.A. and Wheatcroft, R.A. in press: Subcritical ripple formation induced by biogenic mounds, *Marine Geology*.

Wheatcroft, R.A., Starczak, V.R. and Butman, C.A. in press: The impact of population abundance on the deposit-feeding rate of a cosmopolitan worm, *Limnology and Oceanography*.