

Extension of Sediment Geochronology to Coarse-Grained Sediments

Steven A. Kuehl
Virginia Institute of Marine Science
College of William and Mary
Gloucester Pt., VA 23062
phone: (804) 684-7118 fax: (804) 684-7118 email: kuehl@vims.edu

Award Number: N00014-05-1-0088
<http://www.vims.edu/>

LONG-TERM GOALS

Knowledge of seabed dynamics in nearshore environments is needed to address problems ranging from to the evolution of coastal geomorphology to the changing nature of seabed acoustical properties. One successful approach, used primarily in fine-grained sedimentary environments, has been the application short-lived particle-reactive radioisotopes to quantify sediment deposition and accumulation rates, and to measure the depth and intensity of physical and biological mixing. Unfortunately, the utility of this approach in coarser sediments typical of coastal environments has been hindered by the low inherent concentration of these isotopes in sandy sediments, and the relatively high lower limit of detection using traditional decay counting techniques. The proposed study would attempt to extend this approach into sandy sedimentary environments by taking advantage of a relatively new measurement technique, high-resolution inductively coupled plasma-mass spectroscopy (ICP-MS).

OBJECTIVES

Recent studies by Kenna (2002) have successfully measured Pu concentrations as low as 0.5 fg g^{-1} , a value roughly two orders of magnitude lower than possible by traditional decay counting techniques. $^{239/240}\text{Pu}$ is one of the major byproducts of thermonuclear explosions, and hence is a key time marker for the period since 1954, the onset of intensive atmospheric weapons testing. This study would attempt to measure $^{239/240}\text{Pu}$ concentrations in sediment cores collected from Duck, NC. A transect of core samples with a range of textural characteristics will be obtained from other ongoing studies of this area and analyzed for $^{239/240}\text{Pu}$. The major goal will be to determine the limits of this new technology in its application to sediment geochronology of sandy sediments.

APPROACH

Recent advances in high-resolution inductively coupled plasma-mass spectroscopy (HR ICPMS) present an unparalleled opportunity to advance the measurement of environmental levels of bomb Pu isotopes in sediment geochronology. A combination of advances in instrumentation and separation techniques has increased the sensitivity of bomb Pu measurement by two orders of magnitude over that typically achieved by traditional decay counting techniques. In fact, the present technology allows the measurement of Pu levels as low as 0.5 fg g^{-1} (Kenna, 2002). For perspective, a rough equivalent in terms of radioactivity is only a single disintegration per day, a quantity that could never be measured using decay counting. HR ICPMS measurement of bomb Pu, therefore, is likely to replace traditional

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2005		2. REPORT TYPE		3. DATES COVERED 00-00-2005 to 00-00-2005	
4. TITLE AND SUBTITLE Extension of Sediment Geochronology to Coarse-Grained Sediments				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Virginia Institute of Marine Science, College of William and Mary, Gloucester Pt, VA, 23062				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT Knowledge of seabed dynamics in nearshore environments is needed to address problems ranging from to the evolution of coastal geomorphology to the changing nature of seabed acoustical properties. One successful approach, used primarily in fine-grained sedimentary environments, has been the application short-lived particle-reactive radioisotopes to quantify sediment deposition and accumulation rates, and to measure the depth and intensity of physical and biological mixing. Unfortunately, the utility of this approach in coarser sediments typical of coastal environments has been hindered by the low inherent concentration of these isotopes in sandy sediments, and the relatively high lower limit of detection using traditional decay counting techniques. The proposed study would attempt to extend this approach into sandy sedimentary environments by taking advantage of a relatively new measurement technique, high-resolution inductively coupled plasma-mass spectroscopy (ICP-MS).					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

decay counting techniques in the near future for application to problems in marine sediment geochronology. The dramatically lower limit of detectability achieved using this technique should allow us to extend studies heretofore limited to fine-grained sedimentary environments to the sandy environments of the nearshore and inner continental shelf.

Measurement of Pu isotopes will be made using an approach adapted from that of Kim et al. (2000) and Kenna (2002). Sample material (5-10 g) will be spiked with a ^{242}Pu reference solution to monitor chemical yield. The sample will then be combusted at 550° C to remove organic matter before partial acid leaching with HNO_3 . Following filtration, the Pu will be oxidized from Pu(III) to Pu(IV) using a sodium nitrate solution. Separation and purification of the Pu isotopes in solution will be made using TEVA-SpecTM ion exchange resin columns as an actinide-nitrate complex. Eluted samples with Pu will be dried and brought to a final volume of 1 ml in 10% HNO_3 for measurement on the ICP-MS. Using a similar approach, Kenna (2002) was able to achieve chemical yields between 70% and 95%.

WORK COMPLETED

All ordering of supplies and laboratory reagents to set up the new Pu extraction facility at VIMS has been completed. The VIMS radiation safety office has expended considerable effort over the past year in order to add Pu to the institute's radioisotope license. This is needed before a Pu standard can be purchased for this project. A major change in the radioisotope licensing authority for VIMS, from region II to region I, has resulted in an unusual delay in NRC approval. VIMS is required to renew its license on a 5 year renewal period and up until last year had dealt with NRC Region II, Atlanta, GA. Because of their familiarity with the VIMS RSO, adding new isotopes had previously been a relatively straightforward and timely process. The change to Region I, King of Prussia, PA, required an entirely new program/license application for the institute. In addition, for the first time, VIMS has had to provide a financial assurance instrument in accord with the type, quantity and risk involved with the particular type of RAM on hand. This process is now complete and we now expect approval by the NRC in the next few weeks.

RESULTS

As indicated above, all of the preliminary preparation and set up for the Pu extraction line has been completed. The unforeseen complexity in renewal of the VIMS radioisotope license, including the addition of Pu to the license, has delayed the purchase of the necessary Pu radioisotope standard. However, the approval process is nearly complete and we anticipate being able to begin extractions later this Fall.

REFERENCES

- Dellapenna, T.M., S.A. Kuehl, and L.C. Schaffner, 1998, Seabed mixing and particle residence times in biologically and physically dominated estuarine systems: A comparison of lower Chesapeake Bay and the York River subestuary, *Estuarine, Coastal and Shelf Science*, v. 46, pp. 777-795.
- Kenna, T.C., 2002, Determination of plutonium isotopes and neptunium-237 in environmental samples by inductively coupled plasma mass spectrometry with total sample dissolution, *Journal of Analytical Atomic Spectrometry*, v, 17, pp. 1471-1479.

Kim, C.S., C.K Kim, J.I. Lee, and K.J. Lee, 2000, Rapid determination of Pu isotopes and atom ratios in small amounts of environmental samples by an on-line sample pre-treatment system and isotope dilution high resolution inductively coupled plasma mass spectrometry, *Journal of Analytical Atomic Spectrometry*, v. 15, pp. 247-255.

Kuehl, S.A., C.A. Nittrouer, M.A. Allison, L.E.C. Faria, D.A. Dukat, J.M. Jaeger, T.D. Pacioni, A.G. Figueiredo, and E.C. Underkoffler, 1996, Sediment deposition, accumulation, and seabed dynamics in an energetic fine-grained coastal environment, *Continental Shelf Research*, v. 16, pp. 787-815.

Santschi, P.H., Y.H Li, D.M. Adler, M. Amdurer, J. Bell, and U.P Nyffeler, 1983, The relative mobility of natural (Th, Pb and Po) and fallout (Pu, Am, Cs) radionuclides in the coastal marine environment: Results from model ecosystems (MERL) and Narragansett Bay, *Geochimica et Cosmochimica Acta*, v. 47, pp 201-210.