

Biogenic Ripple Destruction: Rates, Modes and Subsurface Consequences

Robert A. Wheatcroft
College of Oceanic & Atmospheric Sciences
Oregon State University
Corvallis, OR 97331
phone: (541) 737-3891 fax: (541) 737-2064 e-mail: raw@coas.oregonstate.edu

Award Number: N00014-04-1-0537
<http://www.coas.oregonstate.edu/faculty/wheatcroft.html>

LONG-TERM GOALS

The ultimate objective of this research program is to obtain a predictive understanding of the physical and biological processes that control the microtopography of the sediment-covered seafloor. Our primary means to address this topic is to make focused field measurements that extend and test conceptual frameworks.

OBJECTIVES

This project, part of the “Ripple DRI”, had as its goal the quantitative measurement of ripple destruction rates and modes, as well as their subsurface consequences. It is important to develop a predictive understanding of ripple destruction for at least two reasons. First, in many environments where ripples occur they are ephemeral features. That is, the near-bed flows are of insufficient magnitude to move sediment continuously as bed load (a prerequisite of ripple formation and maintenance). During the periods between sediment transport events, ripples in most settings will be subject to biogenic destruction, and all of the physical, chemical and biological consequences of the ripples will be altered or cease to occur altogether. Second, under weak transport conditions ripple geometry likely depends on the preceding bed morphology (i.e., the seabed has a memory, Fries et al. 1999). Thus, without better knowledge of the evolution of ripple geometry during non-transport periods (i.e., during the destructive phase) it will be more difficult to predict morphology when the flow increases above the critical shear stress (the constructive phase). The second general but related area of inquiry in this project involves ripples as templates for subsurface structure. By “templates” we mean that ripples, because they perturb near-bed and subsurface fluid and particle fluxes (e.g., Huettel et al. 1996), may cause repeatable subsurface variations in diverse properties, especially bulk density.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 2005		2. REPORT TYPE		3. DATES COVERED 00-00-2005 to 00-00-2005	
4. TITLE AND SUBTITLE Biogenic Ripple Destruction: Rates, Modes and Subsurface Consequences				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University, College of Oceanic & Atmospheric Sciences, Corvallis, OR, 97331				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT The ultimate objective of this research program is to obtain a predictive understanding of the physical and biological processes that control the microtopography of the sediment-covered seafloor. Our primary means to address this topic is to make focused field measurements that extend and test conceptual frameworks					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

APPROACH

Our focused set of field measurements at the SAX04 (Ft. Walton Beach, FL) and nearby sites was to include:

- (1) deliberate, noble-metal tracer experiments to quantify horizontal and vertical biodiffusivities in the presence/absence of transient bioturbators (fish),
- (2) time-series measurements of ripple profiles using a diver-operated laser system, and
- (3) high-resolution vertical profiles of porosity using an *in situ* resistivity probe (Wheatcroft, 2002). The first two measurements were designed to test a simple model relating ripple destruction rate to the intensity of horizontal bioturbation, whereas the latter measurement extends results obtained during SAX99 that suggest ripples may yield deterministic porosity patterns in the subsurface.

WORK COMPLETED

Our field measurements were wholly dependent on divers. Therefore, the nearby passage of hurricane Ivan on 16 September had major negative effects on our research plan. We were the first divers in the water following the hurricane (on 19 September) and found visibility at the bottom to be essentially zero. Over the next week conditions improved to <0.5 m visibility, but this level still precluded most of our measurements. We returned in late October, when conditions were marginally better (2+ m visibility), but by then could not conduct our tracer or profile measurements. It is important to recognize that there was a very strong vertical suspended-sediment concentration gradient such that water even 2 m above bottom had much higher visibility. Unfortunately, we were working in the lower 0.5 m. All was not lost, however, as we were able to make a large number of resistivity measurements.

RESULTS

Two results are highlighted. First, mud delivered to the study area following hurricane Ivan preferentially accumulated in ripple troughs compared to crests. Therefore, the subsurface porosity (or bulk density) field underlying ripple troughs showed much greater vertical fluctuations (Fig. 1), which has potentially important implications for sound scattering. Second, we detected a clear relationship between the porosity of buried mud layers and the thickness of the overlying sand layer (Fig. 2), whereby thicker sand layers resulted in a more fully consolidated mud.




Figure 1. A subset of porosity profiles obtained in situ from the upper 6 cm of the seabed at the SAX04 study site in late October 2004. Profiles collected from ripple troughs had many subsurface fluctuations, whereas profiles from ripple crests did not.




Figure 2. A plot of sand layer thickness (mm) versus mud layer porosity showing a linear relationship ($r^2 = 0.79$), whereby sand-layer thicknesses of <5 mm resulted in mud porosities of >0.8, whereas a sand layer thickness of 40 mm resulted in a porosity <0.5.

An additional, albeit qualitative, result was our observation that fish abundance was higher adjacent to seafloor structures such as tripods, and the resultant microtopography was more heavily impacted by their foraging activities.

RELATED PROJECTS

Our rippled degradation research is related to that of Richardson (NRL), Boudreau and Hay (Dalhousie). Their acoustical measurements and automaton models of ripple degradation would have benefited from the tracer experiments and our proposed caging experiments that were intended to exclude transient bioturbators. We continue to interact with these colleagues, with the hope of making improved field measurements of ripple degradation in the future. In addition, our resistivity measurements complement those made by Tang (APL-UW).

REFERENCES

Fries, J.S., C.A. Butman & R.A. Wheatcroft. 1999. Ripple formation induced by biogenic mounds. *Marine Geology*, 159: 289-303

Huettel, M., W. Ziebis and S. Forster. 1996. Flow-induced uptake of particulate matter in permeable sediments. *Limnology and Oceanography*, **41**: 309-322.

Wheatcroft, R.A. 2002. In situ measurements of near-surface porosity in shallow-water marine sands. *IEEE Journal of Oceanic Engineering*, 27: 561-570.

PUBLICATIONS

Wheatcroft, R.A., P.L. Wiberg & 6 others. 2006a. Post-depositional alteration and preservation of sedimentary strata. In: *Continental Margin Sedimentation: From Sediment Transport to Sequence Stratigraphy*, C. A. Nittrouer et al. (eds), Blackwell Press, (in press).

Wheatcroft, R.A. Time-series measurements of macrobenthos and bioturbation intensity on a flood-dominated continental shelf, northern California. *Progress in Oceanography* (in review).