

Coupled Ocean-Atmosphere Modeling of the Coastal Zone

Eric D. Skyllingstad

College of Oceanic and Atmospheric Sciences, Oregon State University

104 Ocean Admin. Bldg., Corvallis, OR 97331

Phone: (541) 737-5697 Fax: (541) 737-2540 Email: skylling@coas.oregonstate.edu

Roger M. Samelson

College of Oceanic and Atmospheric Sciences, Oregon State University

104 Ocean Admin. Bldg., Corvallis, OR 97331

Phone: (541) 737-4752 Fax: (541) 737-2064 Email: rsamelson@coas.oregonstate.edu

Award Number: N00014-06-1-0328

LONG-TERM GOAL

The long-range goal of this project is to improve our ability to understand and predict environmental conditions in the coastal zone.

OBJECTIVES

The primary scientific objectives of the proposed research are to use a coupled atmosphere--ocean model to investigate and quantify the interaction between the oceanic and atmospheric boundary layers and its effect on environmental conditions in the coastal zone. The main focus will be on boundary layer interactions under coastal upwelling conditions, in which cold, upwelled ocean surface water induces the development of stable internal boundary layers in the atmosphere and thereby reduces low-level winds and surface stress.

APPROACH

The approach used in this project is to combine numerical model results with in-situ and remote-sensing observations to understand and quantify physical processes in the coastal, coupled atmosphere-ocean and test their representation in mesoscale atmospheric models.

WORK COMPLETED

Work has continued on testing a coupled version of the COAMPS and ROMS models. Results from an idealized coastal upwelling scenario have been reported in a paper published in the Journal of Physical Oceanography (Perlin et al., 2007). Further model development has continued on generalizing the coupled model for domains with differing grid sizes. Extension of the model to more realistic coastal domains has begun with an initial case representing a coastal point or cape feature.

RESULTS

Work performed under this research grant has shown that the effect of SST gradient is significant for near-shore wind forcing. Perlin et al. (2007) performed idealized experiments using a coupled

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2007		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Coupled Ocean-Atmosphere Modeling Of The Coastal Zone				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University, College of Oceanic and Atmospheric Sciences, 104 Ocean Admin. Bldg, Corvallis, OR, 97331				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

atmosphere/ocean model based on COAMPS and ROMS developed for the purpose of studying mesoscale air-sea interaction. Result from these experiments (Figure 1) suggest that air-sea interaction during coastal upwelling can reduce surface wind stress by as much as a factor of 2 over distances of about 20 km, as cold water outcrops near the coast and causes the development of an internal boundary layer in the atmosphere. The effects on surface stress are sufficiently strong that they can begin to affect the ocean upwelling circulation on relatively short timescales. Between 3 and 15 km from the coast, offshore Ekman transport in these simulations increased more rapidly with offshore distance, and vertical velocities at the base of the surface layer were stronger for the coupled case than for the uncoupled (ocean only) case. The coupled case also generated a thinner coastal surface layer with gradual seaward deepening. Overall, adjustment of surface Ekman divergence to the coastal boundary occurred over a slightly greater offshore distance in the coupled case than in the uncoupled case. Because of the wind reduction in the coupled case, vertical mixing in the upwelling zone was weaker, resulting in a shallower surface mixed layer. This delayed the merging of the surface and bottom boundary layers over the inner shelf and allowed further inshore propagation of the bottom upwelling front. Within 2 km off the coast, local cross-shore transport estimates were slightly greater and SST slightly cooler for the coupled case. Weaker near shore wind stresses in the coupled case also resulted in consistently smaller alongshore transport in the oceanic surface layer.




Figure 1. Surface meridional wind stress at the end of a 72-h run for two simulations. Note the factor of 2 reduction in wind stress at the model coastline. (from Perlin et al. 2007).

Results from the idealized channel flow simulations support the general hypothesis that SST variability can have significant control over the surface wind stress. However, observations (e.g., Perlin et al., 2004) show that variations in coastal topography can also cause large changes in the coastal wind forcing. Therefore, including coastal variations into the coupled simulation model was a principal goal in our most recent proposed effort.

As in the channel flow case, we first examined an idealized scenario for topographic variability for two coastal bends as shown in Figure 3a. Periodic boundaries were again used in the north-south direction with open boundaries along the east and west. The advantage in this approach over a full, realistic simulation of the coast line is the ability to distinguish the role of the terrain and SST variations from other potential influences, such as land surface variability, diurnal heating, and unresolved topographic features.




Figure 3. Sea-surface temperature ($^{\circ}\text{C}$ shaded) and wind stress (N m^{-2} , vectors and contours) after 192 hours of simulated time. Note the sharp decrease in wind stress from north to south along the coast.

Development and documentation of the coupled model will continue with adaptation of the Earth System Modeling Framework (www.esmf.ucar.edu). N. Perlin is currently involved with a NOPP project to couple a sediment transport model with ROMS using ESMF as the overall model framework. This work complements our coupled modeling efforts and will leverage further development of the model interface.

TRANSITIONS

The coupled model code has been made available to the general community. For example M. Spall (WHOI, personal communication) is using the model to study open-ocean air-sea coupling in the North Atlantic.

REFERENCES

Perlin, N., R. M. Samelson, and D. B. Chelton, 2004. Scatterometer and model wind and wind stress in the Oregon - northern California coastal zone. *Monthly Weather Review*, 132, 2110-2129.

PUBLICATIONS

- Perlin, N., E. D. Skillingstad, R. M. Samelson, and P. L. Barbour, 2007. Numerical simulation of air-sea coupling during coastal upwelling, *J. Phys. Oceanogr.*, 37, 2081-2093.
- Skillingstad, E. D., D. Vickers, L. Mahrt, and R. Samelson, 2007. Effects of mesoscale sea-surface temperature fronts on the marine boundary layer. *Boundary-Layer Meteorol.*, 123, 219-237.