

Correspondence

Infection of the Gulf Coast tick, *Amblyomma maculatum* (Acari: Ixodidae), with *Rickettsia parkeri*: first report from the State of Delaware

DAVID A. FLORIN¹, JU JIANG², RICHARD G. ROBBINS³ & ALLEN L. RICHARDS^{1,2}

¹ Uniformed Services University of the Health Sciences, Department of Preventive Medicine and Biometrics, 4301 Jones Bridge Road, Bethesda, MD 20814-4799, U.S.A. E-mail: david.florin@usuhs.edu

² Naval Medical Research Center, Viral and Rickettsial Diseases Department, 503 Grant Avenue, Silver Spring, MD 20910-7500, U.S.A.

³ Armed Forces Pest Management Board, Office of the Deputy Under Secretary of Defense for Installations and Environment, Building 172, U.S. Army Garrison Forest Glen, Silver Spring, MD 20910-1230, U.S.A.

Abstract

The molecular detection of *Rickettsia parkeri* in a Gulf Coast tick, *Amblyomma maculatum*, collected in Delaware represents the first evidence of the human pathogen *R. parkeri* associated with *A. maculatum* in the state. A total of five adult (2 male and 3 female) Gulf Coast ticks were collected from tick drags conducted during a two-day sampling event (21–22 May 2012) at Bombay Hook National Wildlife Refuge, near Smyrna, Delaware. All specimens were tested for the presence of *Rickettsia* with a genus-specific quantitative real-time polymerase chain reaction (qPCR) assay; one of the female specimens tested positive. This specimen was then assessed for the presence of *Rickettsia parkeri* and *Candidatus Rickettsia andeanae* by two species-specific qPCR assays. The presence of *R. parkeri* DNA was detected, whereas *Candidatus R. andeanae* DNA was not.

Key words: *Amblyomma maculatum*, *Rickettsia parkeri*, Delaware

Introduction

The Gulf Coast tick, *Amblyomma maculatum* Koch, has a range that extends from Peru to the southeastern United States, with a documented presence in many northeastern states, including New York, Connecticut and Maine (Teel *et al.* 2010). The occurrence of the species in these far northern states has generally been viewed as incidental dispersion; larval and/or nymphal stages are displaced from the Southeast when attached to migrating avian hosts. While the displaced immature stages may eventually molt into adults, the adult ticks do not complete a life cycle in these areas, probably because no stage is able to overwinter. Yet, the species does appear to be undergoing a range expansion from the past range boundary of South Carolina to the mid-Atlantic East Coast, as evidenced by large numbers of collected specimens, multiple collection sites, and multi-year data from North Carolina (Varela-Stokes *et al.* 2011) and Virginia (Wright *et al.* 2011, Formadel *et al.* 2011) that indicate established, viable populations in those states. In Delaware, the occurrence of *A. maculatum* has been documented only once (Lancaster 1973) in a listing that is suspected of being based upon incidental collection(s) – there is no evidence in the available literature that the species has been consistently found in the state during field collections or as an ectoparasite of animals/humans.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2013		2. REPORT TYPE		3. DATES COVERED	
4. TITLE AND SUBTITLE Infection of the Gulf Coast tick, Amblyomma maculatum (Acari: Ixodidae), with Rickettsia parkeri: first report from the State of Delaware			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Uniformed Services University of the Health Sciences, Department of Preventive Medicine and Biometrics, 4301 Jones Bridge Road, Bethesda, MD, 20814			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Materials and methods

This report describes a single 2-day (21–22 May 2012) collecting trip to the Bombay Hook National Wildlife Refuge (NWR), near Smyrna, Delaware, where field drags were used to collect ticks. The single collection methodology and the extremely short collection period prevented determination of whether or not an established population of *A. maculatum* exists at the site. Instead, our goal was to collect any and all *A. maculatum* and test the collected specimens for the presence of *Rickettsia parkeri*, the causative agent of Tidewater spotted fever in humans (Jiang *et al.* 2012). The collection event was part of a larger effort to determine the geographical range of *R. parkeri* in the mid-Atlantic and New England regions. The time frame of mid May was selected to optimize the collection of adult *A. maculatum*. All collections were made within approximately 150 m of the GPS coordinates 39° 15' 47.8"N, 75° 28' 07.2"W, in a field of secondary growth near the entrance to the hiking trail leading to the Shearneck Observation Tower. All specimens were immediately preserved in 100% ethanol and later transported to the Naval Medical Research Center, Silver Spring, MD, where processing of the ticks and molecular analysis were conducted. Laboratory materials and methods used have been previously published in Jiang *et al.* (2012).

Results and discussion

A total of five adult (2 male and 3 female) Gulf Coast ticks were collected at Bombay Hook NWR. Additionally, 55 specimens (21 adults, 34 nymphs) of *Amblyomma americanum* (Linnaeus) were collected, as were 13 adults of *Dermacentor variabilis* (Say). Molecular analyses for *R. parkeri* were conducted on *A. americanum* and *A. maculatum*, with all of the *A. americanum* testing negative.

Of the five *A. maculatum* specimens, a single female was positive for the *Rickettsia* genus-specific 17-kDa protein gene (*htrA*) (Ct value: 20.99) and subsequently was positive for the *Rickettsia parkeri*-specific qPCR assay (Ct value: 20.81; calculated *R. parkeri* genome copies in this tick: 1.69×10^8) but negative for the *Candidatus R. andeanae*-specific qPCR assay. We believe that this is the first time *R. parkeri* has been detected in Delaware within a collected tick specimen, thus adding to the U.S. geographic distribution of *R. parkeri* described in Sumner *et al.* (2007). It is noteworthy that a confirmed human case of *R. parkeri* infection was recently reported from Harford County, in northern Maryland, about 80 km from Bombay Hook NWR (Paddock *et al.* 2008). The next step is to determine whether there is an established population of *A. maculatum* at the Bombay Hook NWR and what percentage of the population is infected with *R. parkeri*, so that a robust epidemiological risk assessment can be developed.

Acknowledgements

This research was funded by USUHS intramural grant R087M6, Surveillance for *Rickettsia parkeri* in ticks collected from Maryland and the northeast region of the United States, and DoD AFHSC/GEIS work unit #0000188M.0931.001.A0074. The views expressed herein are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, or the U.S. Government. This work was prepared as part of official military duties of the lead author. Title 17 U.S.C. §105 provides that "Copyright protection under this title is not available for any work of the United States Government." Title 17 U.S.C. §101 defines a U.S. Government work as a work prepared by a military service member or employee of the U.S. Government as part of that person's official duties.

References

- Formadel, C.M., Zhang, X., Smith, J.D., Paddock, C.D., Arias, J.R. & Norris, D.E. (2011) High rates of *Rickettsia parkeri* infection in Gulf Coast ticks (*Amblyomma maculatum*) and identification of “*Candidatus Rickettsia andeanae*” from Fairfax County, Virginia. *Vector-Borne and Zoonotic Diseases*, 11, 1535–1539.
<http://dx.doi.org/10.1089/vbz.2011.0654>
- Jiang, J., Stromdahl, E.Y. & Richards, A.L. (2012) Detection of *Rickettsia parkeri* and *Candidatus Rickettsia andeanae* in *Amblyomma maculatum* Gulf Coast ticks collected from humans in the United States. *Vector-Borne and Zoonotic Diseases*, 12, 175–182.
<http://dx.doi.org/10.1089/vbz.2011.0614>
- Lancaster, J.L., Jr. (1973) *A Guide to the Ticks of Arkansas*. Bulletin 779. Agricultural Experiment Station, Division of Agriculture, University of Arkansas, Fayetteville. 39 pp.
- Paddock, C.D., Finley, R.W., Wright, C.S., Robinson, H.N., Schrodt, B.J., Lane, C.C., Ekenna, O., Blass, M.A., Tamminga, C.L., Ohl, C.A., McLellan, S.L.F., Goddard, J., Holman, R.C., Openshaw, J.J., Sumner, J.W., Zaki, S.R. & Ereemeeva, M.E. (2008) *Rickettsia parkeri* rickettsiosis and its clinical distinction from Rocky Mountain spotted fever. *Clinical Infectious Diseases*, 47, 1188–1196.
<http://dx.doi.org/10.1086/592254>
- Sumner, J.W., Durden, L.A., Goddard, J., Stromdahl, E.Y., Clark, K.L., Reeves, W.K. & Paddock, C.D. (2007) Gulf Coast ticks (*Amblyomma maculatum*) and *Rickettsia parkeri*, United States. *Emerging Infectious Diseases*, 13, 751–753.
<http://dx.doi.org/10.3201/eid1305.061468>
- Teel, P.D., Ketchum, H. R., Mock, D.E., Wright, R.E. & Strey, O.F. (2010) The Gulf Coast tick: a review of the life history, ecology, distribution, and emergence as an arthropod of medical and veterinary importance. *Journal of Medical Entomology*, 47, 707–722.
<http://dx.doi.org/10.1603/ME10029>
- Varela-Stokes, A.S., Paddock, C.D., Engber, B. & Toliver, M. (2011) *Rickettsia parkeri* in *Amblyomma maculatum* ticks, North Carolina, USA, 2009–2010. *Emerging Infectious Diseases*, 17, 2350–2353.
<http://dx.doi.org/10.3201/eid1712.110789>
- Wright, C.L., Nadolny, R.M., Jiang, J., Richards, A.L., Sonenshine, D.E., Gaff, H.D. & Hynes, W.L. (2011) *Rickettsia parkeri* in Gulf Coast ticks, southeastern Virginia, USA. *Emerging Infectious Diseases*, 17, 896–898.
<http://dx.doi.org/10.3201/eid1705.101836>

Accepted by Owen Seeman: 30 Nov. 2012; published 31 Mar. 2013