

Analysis of Data From TCS-08

David J. Raymond
Physics Department
New Mexico Tech
801 Leroy Place
Socorro, NM 87801
phone: (575) 835-5610 fax: (575) 835-5707 email: raymond@kestrel.nmt.edu

Award Number: N000140810241
<http://physics.nmt.edu/~raymond>

LONG-TERM GOALS

The long-term goals of this project are to understand better the mechanisms operative in tropical cyclogenesis and to transfer this knowledge to large-scale models in order to improve forecasts of tropical storm formation. Since convection constitutes the biggest uncertainty in this process, our focus is to understand how convection affects and is affected by cyclone-scale flows.

OBJECTIVES

Our objective in this segment of the program was to use our vorticity and thermodynamic analyses of developing and non-developing tropical cyclones to understand better the process of tropical cyclogenesis. In addition to results from TCS-08, we have added results from ITOP and the NSF-funded PREDICT project.

APPROACH

The TCS-08, ITOP, and PREDICT field programs provided a rich set of data including Eldora radar data (TCS-08) and dropsondes (all projects) from a variety of platforms. We have analyzed the data from all three of these projects, focusing on the production of the three-dimensional variational analyses (3D-VAR) of winds and thermodynamic data for all potential cyclogenesis cases. From these analyses, we have developed a coherent picture of the mechanisms underlying tropical cyclogenesis.

WORK COMPLETED

Graduate student Saška Gjorgjievska has finished and written up the diagnostic analysis of about two dozen tropical cyclogenesis missions from the TCS-08 and PREDICT field programs. Our vorticity analysis scheme of dropsonde and airborne Doppler radar data allowed her to extract the strength of the low-level and mid-level circulations in these cases, as well the circulation tendencies at each level. The balanced response to such a circulation is the development of a warm anomaly

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2012		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Analysis of Data From TCS-08				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Physics Department New Mexico Tech 801 Leroy Place Socorro, NM 87801				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




Figure 1: **Instability index vs. mean mid-level absolute vorticity for tropical cyclogenesis missions flown during TCS-08 and PREDICT. H2 and N2 indicate respectively a non-developing (Hagupit2) and a rapidly developing tropical disturbance (Nuri2).**

above the circulation maximum and a cold anomaly below. We characterized the development of these virtual temperature anomalies by an “instability index”

$$I = s^*(low) - s^*(mid)$$

where s^* is the saturated moist entropy, “low” indicates a disturbance-wide average over 1 – 3 km, and “mid” indicates an average over 5 – 7 km. Smaller instability index corresponds to a stronger warm-cold dipole of this type. Figure 1 shows that low instability index is correlated with strong mid-level vorticity, a result that would be expected if the temperature dipole is indeed a balanced response to convection and not, for instance, an unbalanced response to the convection.

The significance of instability index is its influence on convective mass fluxes. High instability index results in top-heavy vertical mass fluxes, as indicated in the example of Hagupit2 (H2 in figure1) in figure 2. Hagupit eventually intensified into a typhoon, but at the time of observation was a non-developing tropical wave. In contrast, Nuri2 (N2 in figure 1) was intensifying rapidly when observed and as figure 2 shows, its mass flux profile is bottom-heavy. The other cases in our sample behaved similarly. Levels at which the mass flux increases with height exhibit mass and vorticity convergence, resulting in increasing circulation with time, whereas decreasing mass flux with height shows the reverse. Thus an environment with high instability index results in the spinup of a mid-level vortex, whereas low instability index causes a low-level vortex to form and intensify. Figure 3 verifies this result for our data set.

One loose end is the plausible but unproven assertion that the temperature dipole observed in conjunction with a mid-level vortex is a balanced response to the vortex. To address this question, Raymond applied a non-linear balance potential vorticity inversion code (Raymond 1992) to an axisymmetric idealization of the observed potential vorticity distributions in two TCS-08 cases, Nuri1 and Nuri2. The upper panels of figure 4 show the observed and assumed potential vorticity profiles averaged over the two cases, while the lower panel shows the actual virtual potential temperature profiles and the profiles reconstructed from the potential vorticity inversion. The




Figure 2: Vertical mass flux profiles for the Nuri2 and Hagupit2 tropical cyclogenesis cases. Nuri2 was intensifying, Hagupit2 was not.




Figure 3: Low-level vorticity tendency vs. instability index. Large green symbols indicate spinup into a tropical storm within 48 hr of the observation.




Figure 4: **Horizontally averaged potential vorticity (upper panel) and virtual potential temperature perturbation (lower panel) for Nuri1 (left) and Nuri2 (right).** The thick dashed red line in the upper panel represents the ambient potential vorticity, while the thinner dashed red lines represent areally averaged observations in both panels. The solid blue lines show the averages over a corresponding region in the balanced model result.

agreement between the actual and reconstructed profiles is reasonably good, providing evidence that the balance assumption is reasonably valid in these cases.

Graduate student Ana Juračić is computing the moist entropy budget of intensifying and mature tropical cyclones using our 3D-VAR analyses. She has added the ITOP observations to our catalog of tropical cyclogenesis cases. Ordinary tropical convection tends to export moist entropy laterally. However, preliminary work indicates that intensifying systems import entropy while the net lateral balance for mature systems is close to zero.

Raymond and graduate student Michael Herman have published an initial paper on the question of causality in frictional convergence. Further work is underway to extend this work, which may be important in understanding the boundary layers of developing tropical cyclones.

RESULTS

In last year's annual report we outlined the following sequence of events for tropical cyclogenesis:

- Normal tropical conditions favor top-heavy convective mass flux profiles.
- Top-heavy mass flux profiles cause spinup of a mid-level vortex.

- A mid-level vortex results in a smaller instability index.
- Smaller instability index promotes convection with a bottom-heavy mass flux profile.
- Bottom-heavy convection promotes low-level spinup and tropical cyclogenesis.
- Cloud-resolving model results support the relationship between instability index and mass flux profile.

The analysis of a much larger sample of potential cyclogenesis cases supports this picture.

Our theoretical analysis of causality in frictional convergence raises questions about the dynamics of tropical cyclones that need to be addressed.

IMPACT/APPLICATIONS

If the above sequence of events for tropical cyclogenesis withstands scrutiny, then it will provide a useful conceptual framework for forecasting cyclogenesis.

Many of the lessons learned in this work have broader applicability to tropical convection in general. The knowledge generated should help in the effort to constrain the behavior of convective parameterizations in global models, thus enhancing the ability of these models to forecast tropical weather.

RELATED PROJECTS

The current project is related to work being pursued under a grant from the National Science Foundation entitled “Cumulus Convection and Large-Scale Tropical Flows”. The purpose of that NSF grant is to refine and extend our understanding of the thermodynamic control of convection and to begin to apply these results to a broader range of tropical disturbances. In addition to the Madden-Julian oscillation, we are particularly interested in equatorial Kelvin waves, tropical easterly waves, and the developmental stages of tropical cyclones. TCS-08 results will advance the purposes of this grant and expertise developed under this and previous NSF support will benefit our Office of Naval Research project as well. Graduate student Michael Herman is supported by this grant.

We also have NSF support for our role in the PREDICT project with a grant entitled “Vorticity and Thermodynamic Budgets in Easterly Waves – PREDICT Participation”. The field program for PREDICT concluded two years ago after obtaining extensive dropsonde data on a number of pre-tropical-depression disturbances as well as a number of non-developing systems. Results from this project complement and strengthen our TCS-08 results. Saška Gjorgjievska is supported by this project.

REFERENCES

Raymond, D. J., 1992: Nonlinear balance and potential-vorticity thinking at large Rossby number. *Quart. J. Roy. Meteor. Soc.*, **118**, 987-1015.

PUBLICATIONS

Gjorgjievska, S. and D. J. Raymond, 2012: Interaction between dynamics and thermodynamics in tropical cyclogenesis. Submitted to *Atmos. Chem. Phys.*

Raymond, D. J., 2012: The balanced thermodynamics of tropical cyclogenesis. Submitted to *Tellus-A*.

Raymond, D. J. and M. J. Herman, 2012: Frictional convergence, atmospheric convection, and causality. *Atmósfera*, **25**, 253-267.

Raymond, D. J., S. L. Sessions, and C. López Carrillo, 2011: Thermodynamics of tropical cyclogenesis in the northwest Pacific. *J. Geophys. Res.*, 116, D18101, doi:10.1029/2011JD015624.

Raymond, D. J. and C. López Carrillo, 2011: The vorticity budget of developing typhoon Nuri (2008). *Atmos. Chem. Phys.*, **11**, 147-163, doi:10.5194/acp-11-147-2011.

López Carrillo, C., and D. J. Raymond, 2011: Retrieval of three-dimensional wind fields from Doppler radar data using an efficient two-step approach. *Atmos. Meas. Tech.*, **4**, 2717-2733, doi:10.5194/amt-4-2717-2011.