

Technical Report ARMET-TR-12013

**SYNTHESIS OF BIS(1-METHYL-3-PROPYL-4-NITRO)IMIDAZOLIUM
DODECAHYDRODODECABORATE AND BIS(1-METHYL-3-PROPYL-5-
NITRO)IMIDAZOLIUM DODECAHYDRODODECABORATE SALTS: A NEW CLASS
OF ENERGETIC FUELS**

Peggy Sanchez
Paul Anderson

February 2013



U.S. ARMY ARMAMENT RESEARCH, DEVELOPMENT AND
ENGINEERING CENTER

Munitions Engineering Technology Center

Picatinny Arsenal, New Jersey

Approved for public release; distribution is unlimited.

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

The citation in this report of the names of commercial firms or commercially available products or services does not constitute official endorsement by or approval of the U.S. Government.

Destroy this report when no longer needed by any method that will prevent disclosure of its contents or reconstruction of the document. Do not return to the originator.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-01-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to Department of Defense, Washington Headquarters Services Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) February 2013		2. REPORT TYPE		3. DATES COVERED (From - To) 2009 to 2011	
4. TITLE AND SUBTITLE SYNTHESIS OF BIS(1-METHYL-3-PROPYL-4-NITRO)IMIDAZOLIUM DODECAHYDRODODECABORATE AND BIS(1-METHYL-3-PROPYL-5-NITRO)IMIDAZOLIUM DODECAHYDRODODECABORATE SALTS: A NEW CLASS OF ENERGETIC FUELS				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHORS Peggy Sanchez and Paul Anderson				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army ARDEC, METC Energetics, Warheads & Manufacturing Technology Directorate (RDAR-MEE-W) Picatinny Arsenal, NJ 07806-5000				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army ARDEC, ESIC Knowledge & Process Management (RDAR-EIK) Picatinny Arsenal, NJ 07806-5000				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) Technical Report ARMET-TR-12013	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Two new salts that ionically bind nitroimidazolium cations with the dodecahydro- <i>closa</i> -dodecaborane anion are reported from the synthesis of the corresponding nitrate, iodide, and potassium salts. These salts represent a category of tailorable energetic oxidizer-fuel compounds that can be synthesized in the aqueous environment under normal atmospheric conditions. Spectroscopic data - including single crystal x-ray diffraction, Fourier transform infrared (FTIR), melting points, and densities - are provided along with thermal gravimetric analysis of each compound.					
15. SUBJECT TERMS Boron Metathesis Energetic cation Energetic fuel Boron oxidation Dodecaborate Decaborate					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 23	19a. NAME OF RESPONSIBLE PERSON P. Sanchez
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (Include area code) (973) 724-2770

CONTENTS

	Page
Introduction	1
Background	2
Test Methods	3
Materials	3
Synthesis of Compound I, Ia, and II	3
Analysis	3
Results	4
Conclusions	4
Recommendations	4
References	5
Appendix	
A FTIR and Melting Point Data	7
B Single Crystal X-ray Analysis	11
Distribution List	17

ACKNOWLEDGMENTS

The authors would like to acknowledge Dr. Damon Parish of the Naval Research Laboratory and the University of Missouri, Columbia for the single crystal x-ray analysis and Dr. Reddy Damavarapu of the U.S. Army Armament Research, Development and Engineering Center, Picatinny Arsenal, New Jersey for providing the heterocyclic energetic salts.

INTRODUCTION

Dodecahydro-*closo*-dodecaborate [*closo*-B₁₂H₁₂]²⁻ and decahydro-*closo*-decaborate [*closo*-B₁₀H₁₀]²⁻ salts are caged boron analogues of aromatic hydrocarbons (fig.1). Due to both their thermal stability up to 800°C and high energy release upon combustion compared to their hydrocarbon analogues the *closo*-borate species are a promising class of compounds for energetic and burn rate enhancement research.




Figure 1

Metathesis reaction of dipotassium dodecahydro-*closo*-dodecaborate salt, [*closo*-B₁₂H₁₂]²⁻, with 1-methyl-5-nitro-propyl imidazolium iodide and with 1-methyl-4-nitro-propyl imidazolium iodide to form the corresponding methyl nitroimidazolium dodecaborate salts **I**, **Ia**, and **II** (all vertices are boron)

BACKGROUND

Despite having a high gravimetric and volumetric energy density, energetic boron research has been limited due to both kinetic and thermodynamic impediments: a kinetic bottleneck experienced during oxidation, and a solid phase oxide layer that encompasses the particle surface, requires high temperatures to remove and consequentially causes ignition difficulties (ref. 1). During combustion in an oxygenated environment, boron is oxidized first to boric acid, then to metaboric acid (HOBO), and finally to diborontrioxide with energy being released during each of the oxidative stages (fig. 2).




Figure 2
Equilibrium of (a) boric acid, (b) metaboric acid, and (c) diborontrioxide

Due to the thermal stability of metaboric acid (HOBO), the rate of reaction of k_1 is far greater than the rate of k_2 , thus, the energy release associated with metaboric acid oxidation during combustion is often not realized because of a kinetic bottleneck in the oxidation pathway. Thermodynamically, boron is inhibited by particle diffusion during combustion. In an oxygen-containing atmosphere, boron as a free metal, is surrounded by a fully oxidized diborontrioxide (B_2O_3) layer, which during ignition restricts further oxidation (ref. 2). At temperatures above the B_2O_3 melting point of 723 K, B_2O_3 liquefies and allows some oxygen to slowly diffuse and react with the boron particle surface. This reaction, though, produces more B_2O_3 that cools, condenses back to the boron surface, and further retards the boron particle combustion.

Closo-borate compounds such as dodecahydrododecaborate (DDB) and decahydro-decaborate (DB) are *closo*-polyhedral boron cages with 12 and 10 vertices, respectively. The DDB and DB possess the stability of benzene in a spherical orientation and are considered amongst the most stable compounds yet identified. The DDB and DB cages exhibit a negative two charge across the surface of the molecule and are susceptible to metathesis chemistry with singly and doubly charged cations. They are not metalloids like boron particulates, and thus effects of the oxide passivation layer typically present on boron particles is not a factor. Recently, Shackelford and coworkers used DDB and carborane DDB to co-precipitate in aqueous solution salts of various triazolium moieties (ref. 3). The reactions proceeded with high yields. The group also showed that open-air bench-top synthesis is possible even with perfluoronated versions of the DDB caged structure (ref. 4). In this follow-on work, they used *cis* and *trans* structures of triazolium-based derivatives of a butane structure. In concept, they effectively produced a molecularly engineered explosive, with an energetic molecule in direct contact with a fuel system.

It is in these regards that bis(1-methyl-3-propyl-4-nitro)imidazolium dodecahydro-dodecaborate salt [MP4NI (DDB)] and bis(1-methyl-3-propyl-5-nitro)imidazolium dodecahydro-dodecaborate salt [MP5NI (DDB)] were successfully synthesized. These new compounds represent a category of materials in which an energetic cation is ionically bound to a boron rich fuel. This is the first known synthesis of a molecule with an oxidative moiety (nitrate) directly bound to the fuel system (borate). The energetic properties of the compounds can be tuned by the use of different energetic cations to vary characteristics, such as oxygen or nitrogen balance and density. Use of DDB or DB eliminates the effect of the oxide layer on boron particulate combustion. This is due to

the simultaneous energy release from intimate contact energetic cation and DDB or DB fuel. The oxidation could proceed without inhibition by diffusion limitations associated with the sluggish metaboric and HOBO formation.

TEST METHODS

Materials

Potassium [*closo*-B₁₂H₁₂]²⁻ was obtained from the University of Missouri at Columbia. Pure MP5NI iodide salt, MP4NI iodide salt, and MP5NI nitrate salt were synthesized at the U.S. Army Armament Research, Development and Engineering Center, Picatinny Arsenal, New Jersey. They were used without further purification and analyzed by melting point and Fourier Transform Infrared (FTIR). This data can be seen in the appendix A.

Synthesis of Compound (I), (Ia), and (II)

An aqueous solution of the methylnitroimidazolium salt (2.27 mmol in 2 mL DI H₂O) is added drop wise with stirring to an aqueous solution of potassium [*closo*-B₁₂H₁₂]²⁻ salt (1.14 mmol in 1 mL DI H₂O) at ambient temperature (fig. 2). A white crystalline solid precipitates out of solution and is filtered and rinsed with excess DI H₂O to remove potassium iodide (KI) or potassium nitrate (KNO₃). The solid is then analyzed thermogravimetrically and spectrophotometrically.

Analysis

Single crystal x-ray diffraction spectroscopy of I and II synthesized from the iodide imidazolium salts was solved and refined using the Bruker SHELXTL software package at the University of Missouri in Columbia. X-ray crystallographic analysis of Ia prepared from the nitrate salt was obtained from the Naval Research Laboratory using SHELXTL-*plus* system of programs.

Infrared absorption data was recorded as nujol mulls suspensions on a Perkin Elmer Spectrum 100 FTIR in the range of 400 to 4000 cm⁻¹ at ambient temperature and analyzed using Perkin Elmer Spectrum software.

Thermogravimetric analysis (TGA) thermograms were recorded from a Perkin Elmer Diamond TGA using ceramic boats. The temperature was increased at a rate of 5°C/min from 50 to 400°C in argon. The argon was passed over a molecular sieve and oxygen traps to ensure it was highly pure. Decomposition and melting points were determined by subtracting the thermogram of the ceramic boat from that of the corresponding borate salt.

RESULTS

Three new crystal structures were reported. MP4NI(DDB) and MP5NI(DDB) were synthesized from MP4NI iodide and MP5NI iodide, respectively, in high yield. MP5NI(DDB) was also synthesized from the corresponding nitrate salt. Crystal spectra for each of the materials are in appendix B. The MP4NI(DDB) and MP5NI(DDB) are white crystalline solids that are insoluble in water.

Traditional use of FTIR or nuclear magnetic resonance proved futile as the results were unable to show the difference between a mixture of the corresponding starting materials and the desired product. To overcome this obstacle, the materials were observed under both FTIR and digital scanning calorimeter.

Dodecaborate is identified in FTIR spectra by boron-hydrogen (B-H) stretching vibrations in the range of 2400 cm^{-1} , and bending bands at 1000 and 700 cm^{-1} (ref. 5). To determine the exact wavelength of this particular dipotassium dodecaborate salt used in the reaction raw material was analyzed in a nujol mulls suspension and the corresponding FTIR spectrum indicated B-H stretching at 2485 cm^{-1} and 1075 cm^{-1} . The FTIR spectra of the reactant imidazolium salts were analyzed and compared to the metathesized DDB product. In all cases, the absorbance of the imidazolium moiety became less intense and the B-H stretch was apparent in the spectrum (app. A). A B-H stretch, in particular the strong, sharp peaks at 2485 cm^{-1} and 1075 cm^{-1} , shifts to the right indicating that there is a molecular attraction between the organic cation and *closo*-dodecaborate salt.

The use of FTIR along with a change in melting point was considered sufficient to determine that the metathesis reaction was successful. Both MP4NI(DDB) and MP5NI(DDB) are high melting point organic salts; TGA of the materials shows melting points at 203.0°C and 214.9°C after background subtraction (app. A). Melting of both compounds seems to occur almost simultaneously with gas evolution and decomposition shortly after. A shift to a single higher melting point from the starting materials, along with FTIR analysis, was used to indicate that a reaction proceeded and single crystal x-ray analysis was then obtained. The structures are shown in appendix B.

CONCLUSIONS

Bis(1-methyl-3-propyl-4-nitro)imidazoliuim dodecahydrododecaborate and bis(1-methyl-3-propyl-5-nitro)imidazolium dodecahydrododecaborate are two new organic *closo*-dodecaborate salts synthesized for potential energetic purposes. They represent a class of materials in which an energetic cation is in intimate contact with a borate fuel. They are a tailorable class of compounds through variation of the chemical properties of the energetic cations.

RECOMMENDATIONS

A thermogravimetric analysis study to explore the decomposition kinetics is recommended. The decomposed material can then be analyzed by x-ray diffraction to detect the presence or absence of metaboric acid, and determine if the kinetic bottleneck has been avoided. Calorimetry should also be obtained to determine heat of formation and combustion.

REFERENCES

1. Zhou, W; Yetter, A; Dryer, F.L.; Rabitz, H.; Brown, R.C.; and Kolb, C.E., "Effect of Flourine on the Combustion of "Clean" Surface Boron Particles," s.1: Elsevier Science Inc., Vol. 112, pp. 507-521, 1998.
2. Burton, R.L. and Krier, H., "High Pressure Boron Ignition and Combustion AASERT Grant Numer N00014-93-1-1206, Office of Naval Research, Arlington, VA, Vols. UILU ENG 96-0509, 1996.
3. Shackelford, Scott A.; Belletire, John L.; Boatz, Jerry A.; Schneider Stefan; Wheaton, Amanda K.; Wight, Brett, A.; Hudgens, Leslie M.; and Ammon, Herman L., "Pairing Heterocyclic Catons with closo-Icosahedral Borane and Carborane Anions. I. Benchtop Aqueous synthesis of Binary Triazolium and Imidazolium Salts with Limited Water Solubility, Organic Leters, Vol. 11, pp. 2623-2626, December 2009.
4. Shackelford, Scott A.; Belletire, John L.; Boatz, Jerry A.; Schneider, Stefan; Wheaton, Amanda K.; Wight, Brett A.; Ammon, Herman L.; Peryshkov, Dmitry V.; and Strauss, Steven H., "Bridged Heterocyclium Dicationic closo-Icosahedral Perfluoroborane, Borane, and Carborane Salts via Aqueous, Open-Air Benchtop Synthesis," Organic Letters, Vol. 12, pp. 2417-2717, December 2010.
5. Saldin, V.I.; Karpenko, M.A.; Ignat'eva, L.N.; Nikolenko, Yu M.; Buznik, V.M.; and Mikhailov, Yu M., "Hexamethylenetetrammonium Dodecahydro-closo-Dodecaborate: Synthesis and Study," Journal of Inorganic Chemistry, Vol. 54, pp. 13-16, January 2009.


APPENDIX A
FTIR AND MELTING POINT DATA

Table A-1
FTIR and melting point data of the dodecaborate and methylnitroimidazolium salts and their metathesized products

Molecular Formula	$K_2B_{12}H_{12}$	$C_7H_{12}N_4O_5$	$C_{14}H_{36}B_{12}N_6O_4$	$C_7H_{12}N_3O_2I$	$C_{14}H_{36}B_{12}N_6O_4$
Molecular Weight (g/mol)	220.02	232.19	482.21	297.09	482.21
TGA					
Melting point (°C)	>800	114±0.5	203.7±0.7	174.2±0.4	215.3±0.4
FTIR					
B-H	2487, 1075, 715	NA	2467, 1058	NA	2467, 1056, 713
NO₂	NA	1522, 1403	1530, 1399	1521, 1400	1520, 1394, 713

Reaction was successful with an increased melting point coupled with the appearance of B-H stretching in the product FTIR spectra.


APPENDIX B
SINGLE CRYSTAL X-RAY ANALYSIS



MP5NI(DDB)							
	Synthesized from MP5NI Iodide Salt				Synthesized from MP5NI Nitrate Salt		
Chemical formula	$C_{14}H_{36}B_{12}N_6O_4$				$C_{14}H_{36}B_{12}N_6O_4$		
Formula weight	482.21				482.21		
Space group	P 2 ₁ /c				P 1 2 ₁ /c 1		
Unit Cell dimensions							
a (Å)	8.8651(2)	α (°)	90		8.8649(19)	α (°)	90
b (Å)	18.1099(4) Å	β (°)	94.2820(10)		18.096(4)	β (°)	94.324(4)
c (Å)	16.2541(3) Å	γ (°)	90		16.288(4)	γ (°)	90
V (Å ³)	2602.25(10) Å ³				2605.5(10) Å ³		
ρ_{calc} /g cm ⁻³	1.231				1.229		
Wavelength	1.54178 Å				0.71073 Å		
Z	4				4		
Temperature (K)	100(2)				100(2)		
Crystal size	0.05 x 0.12 x 0.23 mm				0.48 x 0.17 x 0.03 mm ³		
Crystal habit	clear light colourless plate				clear light colourless plate		
Crystal system	monoclinic				monoclinic		
Theta range for data collection	3.66 to 72.03°				1.68 to 26.52°		
Index ranges	-10 ≤ h ≤ 9, -22 ≤ k ≤ 22, -19 ≤ l ≤ 19				-10 ≤ h ≤ 9, -22 ≤ k ≤ 22, -19 ≤ l ≤ 19		
Reflections collected	37939				23566		
Independent reflections	5007 [R(int) = 0.0261]				5352 [R _{int} = 0.0417]		
F(000)	1016				1016		
Goodness-of-fit on F ²	1.055				1.049		
Largest diff. peak and hole	0.575 and -0.307 e Å ⁻³				0.315 and -0.239 e Å ⁻³		
Absorption coefficient	0.620 mm ⁻¹				0.078 mm ⁻¹		
Max. and min. transmission	0.9697 and 0.8705				0.7454 and 0.6269		
Data / restraints / parameters	5007 / 0 / 324				5352 / 0 / 329		
Refinement method	Full-matrix least-squares on F ²				Full-matrix least-squares on F ²		

Figure B-1

Single crystal x-ray of I and Ia obtained from the University of Missouri and D. Parrish at NRL synthesized from the iodide and nitrate salt, respectively. Analysis shows two positively charged (1-methyl-3-propyl-5-nitro)imidazolium cations ionically bound to one dodecaborate anion.



MP4NI(DDB)			
Chemical formula	$C_{14}H_{36}B_{12}N_6O_4$		
Formula weight	482.21		
Space group	P 1 21/c 1		
Unit Cell dimensions			
a (Å)	8.50210(10)	α (°)	90
b (Å)	19.2046(3)	β (°)	114.3000(10)
c (Å)	8.55620(10)	γ (°)	90
V (Å ³)	1273.28(3)		
$\rho_{calc.}$ /Mg cm ⁻³	1.258		
Wavelength (Å)	1.54178 Å		
Z	2		
Temperature (K)	100(2)		
Crystal size	0.04 x 0.05 x 0.15 mm		
Crystal system	monoclinic		
Theta range for data collection	4.60 to 72.14°		
Index ranges	-10 ≤ h ≤ 10, -23 ≤ k ≤ 23, -10 ≤ l ≤ 10		
Reflections collected	18924		
Independent reflections	2505 [R(int) = 0.0225]		
F(000)	508		
Goodness-of-fit on F ²	1.045		
Largest diff. peak and hole	0.247 and -0.287 eÅ ⁻³		
Absorption coefficient	0.634 mm ⁻¹		
Max. and min. transmission	0.9751 and 0.9109		
Data / restraints / parameters	2505 / 0 / 165		
Refinement method	Full-matrix least-squares on F ²		

Figure B-2

Single crystal x-ray of II obtained from the University of Missouri. Analysis shows two positively charged (1-methyl-3-propyl-4-nitro)imidazolium cations ionically bound to one dodecaborate anion.




Figure B-3
1-x, 1-y, 1-z space grouping of MP5NI(DDb) showing the crystal packing of the material. a) stick diagram showing the view along the a-axis b) stick diagram showing view along the b-axis c) stick diagram showing view along the c-axis




Figure B-4
1-z, 1-y, 1-z space grouping of MP4NI(DDDB) showing the crystal packing of the material. a) stick diagram showing the view along the a-axis b) stick diagram showing view along the b-axis c) stick diagram showing view along the c-axis

DISTRIBUTION LIST

U.S. Army ARDEC
ATTN: RDAR-EIK
RDAR-GC
RDAR-MEE-W, P. Sanchez (2)
Picatinny Arsenal, NJ 07806-5000

Defense Technical Information Center (DTIC)
ATTN: Accessions Division
8725 John J. Kingman Road, Ste 0944
Fort Belvoir, VA 22060-6218

Commander
Soldier and Biological/Chemical Command
ATTN: AMSSB-CII, Library
Aberdeen Proving Ground, MD 21010-5423

Director
U.S. Army Research Laboratory
ATTN: AMSRL-CI-LP, Technical Library
Bldg. 4600
Aberdeen Proving Ground, MD 21005-5066

Chief
Benet Weapons Laboratory, WSEC
U.S. Army Research, Development and Engineering Command
Armament Research, Development and Engineering Center
ATTN: RDAR-WSB
Watervliet, NY 12189-5000

Director
U.S. Army TRADOC Analysis Center-WSMR
ATTN: ATRC-WSS-R
White Sands Missile Range, NM 88002

Chemical Propulsion Information Agency
ATTN: Accessions
10630 Little Patuxent Parkway, Suite 202
Columbia, MD 21044-3204

GIDEP Operations Center
P.O. Box 8000
Corona, CA 91718-8000

Synthesis of bis(1-methyl-3-propyl-4-nitro)imidazolium dodecahydrododecaborate and bis(1-methyl-3-propyl-5-nitro)imidazolium dodecahydrododecaborate salts : A New Class of Energetic Fuels

PART 1. Must be signed before the report can be edited.

- Mr. Steven Nicolich
ARDEC, EWETD (AMSRD-AAR-AEE-W)

PART 2. To be signed either when draft report is submitted or after review of reproduction copy.

This report is approved for publication.

Mr. Steven Nicolich
ARDEC, EWETD (AMSRD-AAR-AEE-W)

Division Chief (Date)

SMCAR Form 49, 1 Nov 94 supersedes SMCAR Form 49, 1 Nov 84 which may be used until supply is exhausted.

Stephen Leong

PDAR-CIS (DATE)