

Status of 1/25° Global HYCOM Simulations

Luis Zamudio¹ & Joe Metzger²

¹Center for Ocean-Atmospheric Prediction Studies,
Florida State University

²Naval Research Laboratory,
Stennis Space Center, Mississippi

Layer Ocean Model Workshop, Ann Arbor, Michigan, 21 - 23 May 2013

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAY 2013		2. REPORT TYPE		3. DATES COVERED 00-00-2013 to 00-00-2013	
4. TITLE AND SUBTITLE Status of 1/25degree Global HYCOM Simulations				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Florida State University,Center for Ocean-Atmospheric Prediction Studies,2000 Levy Avenue, Building A, Suite 292,Tallahassee,FL,32306-2741				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 15	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Background

- The 1/25° HYCOM/NCODA/CICE system (~3.5 km mid-latitude resolution) is scheduled to replace the 1/12° GOFS 3.1 at the Naval Oceanographic Office in FY16
- Preliminary hindcast of 1/25° HYCOM/NCODA ran on Navy DSRC Cray XT5 (Einstein), but it is not currently running on the new Navy DSRC IBMs
 - Hindcast length: May 2010 through January 2012
- Development path of 1/25° non-assimilative model is somewhat lagged due to cost and time
 - HPC Grand Challenge project supplies computer resources
- Miscellaneous results from various 1/25° non-assimilative simulations are presented here along with a process study that focuses on the Labrador Sea

1/25° global HYCOM: GLBa0.04-04.0


- Initialized from rest using T & S from GDEM 3.0
- Spun-up for 10 years with ERA40 climatology
- Topography from a modified version of NRL DBDB2
 - 5 m isobath as land-sea boundary
- Built-in energy loan (thermodynamic) sea ice model
- 32 layers, non-assimilative, no tidal forcing
- Relax SSS toward the Polar Science Center climatology

1/25° global HYCOM: GLBa0.04-04.1/04.2


- Initialized from GLBa0.04-04.0
- Interannual 0.5° NOGAPS forcing: 2003-2010
- Long-term NOGAPS wind stress mean was replaced by ERA40 mean for consistency on the large scale

GLBa0.04 simulations often produce a realistic and robust Gulf Stream

SSH mean over years 5-8



SSH variability over years 5-8



Figures 11 & 12 from Hurlburt et al. (2011, Operational Oceanography)


This was the best-to-date representation of the Gulf Stream from non-assimilative HYCOM; Kuroshio looked equally good

Impact of Increasing Horizontal Resolution on RMS SSH Variability in the Gulf of Mexico Region

1/12° Global HYCOM

1/25° Global HYCOM

From satellite altimeter data







- Core of high variability is farther north at 1/25° in western Gulf
- Higher variability extends farther to the west at 1/25°


Response to Tropical Storms

- ONR sponsor S. Harper asked us to investigate the upper ocean response to tropical storms as part of the Impact of Typhoons on the Ocean in the Pacific (ITOP) project – what is the response in existing global models?
- See an upper ocean response in $1/25^\circ$ HYCOM but it is muted because of the weaker than observed atmospheric signal in 0.5° NOGAPS
 - Typhoon Fanapi had Category 3 winds but only Tropical Storm force winds in NOGAPS
- Anticipate NAVGEM to perform better if distributed on 0.33° grid
 - OPTTEST showed improved tropical cyclone statistics

Upper Ocean Response to Typhoon Fanapi: Sept 2010




Impact of Irminger Ring Generation on the Labrador Sea Deep Convection




- Fan et al. (2013, JPO):
 - Irminger Rings transport buoyant fresh and warm water to Labrador Sea providing resistance to deep convection
 - Formed off west coast of Greenland where topographic slope changes
 - Diameters range from 15-70 km
 - Significant barotropic component to these eddies

Impact of Increasing Horizontal Resolution on EKE

2008
mean:
EKE
0 m




EKE
700 m




Better representation of Irminger Rings in 1/25° HYCOM

Strong Barotropic Component of Simulated Irminger Rings

Rel. Vort. ($1 \times 10^{-5} \text{ s}^{-1}$) at 500 m depth for 2007_001




Rel. Vort. ($1 \times 10^{-5} \text{ s}^{-1}$) at 1500 m depth for 2007_001




Significant relative vorticity signature of Irminger Rings even at 1500 m

Significant Difference in Labrador Sea Deep Convection in 2007 and 2008



The observations of Yashayaev and Loder (2009, GRL) show shallow (deep) Labrador Sea mixed layer during the 2007 (2008) winter season. That is primarily due to weak (strong) wind events (heat loss from the ocean to the atmosphere) during the 2007 (2008) winter season and a **secondary hypothesis is an increase (decrease) in the generation of Irminger Rings during the 2007 (2008) winter season.**

The EKE Time Series Supports the Secondary Hypothesis of Irminger Rings Impacting Winter Deep Convection



EKE at
500 m
in eddy
generation
region

1/25° global HYCOM: GLBb0.04-01.2

- Initialized from GLBa0.04-04.0
- Spun-up for 9 years with ERA40 climatology
- Topography from a modified version of GEBCO
 - 10 cm isobath as land-sea boundary

1/25° global HYCOM: GLBb0.04-01.5


- Initialized from GLBb0.04-01.2
- Interannual 0.5° NOGAPS forcing: Jan 2007 – Dec 2011
- No tidal forcing

1/25° global HYCOM: GLBb0.04-01.7

- Initialized from GLBb0.04-01.5
- Interannual 0.5° NOGAPS forcing: Jul 2007 – Jan 2011
- Tidal forcing


1-hourly Steric SSH

GLBb0.04-01.7: 2008 169 00 steric SSH




Internal Tidal Signature Seen in Steric Sea Surface Height

GLBb0.04-01.5: 2008 168 01 steric SSH



No Tides

GLBb0.04-01.7: 2008 168 01 steric SSH



Tides