

**Measurement of the Absolute Raman Cross
Sections of Diethyl Phthalate, Dimethyl
Phthalate, Ethyl Cinnamate, Propylene
Carbonate, Tripropyl Phosphate,
1,3-Cyclohexanedione, 3'-Aminoacetophenone,
3'-Hydroxyacetophenone, Diethyl
Acetamidomalonate, Isovanillin, Lactide,
Meldrum's Acid, p-Tolyl Sulfoxide, and Vanillin**

R.L. Aggarwal
L.W. Farrar
B.G. Saar
T.H. Jeys
R.B. Goodman

13 September 2013

Lincoln Laboratory
MASSACHUSETTS INSTITUTE OF TECHNOLOGY
LEXINGTON, MASSACHUSETTS



Prepared for the Joint Program Executive Office for Chemical and Biological Defense,
Joint Program Manager for Nuclear, Biological and Chemical Contamination Avoidance
under Air Force Contract FA8721-05-C-0002.

Approved for public release; distribution is unlimited.

This report is based on studies performed at Lincoln Laboratory, a federally funded research and development center operated by Massachusetts Institute of Technology. This work was sponsored by the Joint Program Executive Office for Chemical and Biological Defense (JPEO-CBD), Joint Program Manager for Nuclear, Biological and Chemical Contamination Avoidance (JPM NBC CA), under Air Force Contract FA8721-05-C-0002.

This report may be reproduced to satisfy needs of U.S. Government agencies.

JPEO-CBD security has reviewed this report, and it is releasable to the National Technical Information Service, where it will be available to the general public, including foreign nationals.

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER


Gary Tutungian
Administrative Contracting Officer
Enterprise Acquisition Division

Non-Lincoln Recipients

PLEASE DO NOT RETURN

Permission has been given to destroy this document when it is no longer needed.

Massachusetts Institute of Technology
Lincoln Laboratory

Measurement of the Absolute Raman Cross Sections of Diethyl Phthalate,
Dimethyl Phthalate, Ethyl Cinnamate, Propylene Carbonate, Tripropyl
Phosphate, 1,3-Cyclohexanedione, 3'-Aminoacetophenone,
3'-Hydroxyacetophenone, Diethyl Acetamidomalonate, Isovanillin, Lactide,
Meldrum's Acid, p-Tolyl Sulfoxide, and Vanillin

R.L. Aggarwal

L.W. Farrar

B.G. Saar

T.H. Jeys

Group 82

R.B. Goodman

Group 81

Technical Report 1175

13 September 2013

Approved for public release; distribution is unlimited.

Lexington

Massachusetts

ABSTRACT

We have measured the Raman spectra of several chemicals for 785-nm excitation using a Raman spectrometer with focal length of 500 mm, f-number of 6.5, and resolution of $2.0 \pm 0.1 \text{ cm}^{-1}$. Values of $1.92 \pm 0.12 \times 10^{-29}$, $2.50 \pm 0.15 \times 10^{-29}$, $3.10 \pm 0.19 \times 10^{-28}$, $1.17 \pm 0.07 \times 10^{-29}$, and $1.53 \pm 0.09 \times 10^{-29} \text{ cm}^2$ have been determined for the Raman cross section σ_R of neat Diethyl phthalate (DEP), dimethyl phthalate (DMP), ethyl cinnamate (EC), propylene carbonate (PC), and tripropylene phosphate (TPP) for the Raman modes at 1041, 1040, 1638, 849, and 726 cm^{-1} , respectively. The values of the linewidth (FWHM) Γ for these modes have been determined to be 6.6 ± 0.1 , 7.0 ± 0.1 , 11.3 ± 0.2 , 15.8 ± 0.2 , and $29.0 \pm 0.3 \text{ cm}^{-1}$, respectively. Values of $1.47 \pm 0.12 \times 10^{-28}$, $1.67 \pm 0.13 \times 10^{-29}$, $5.15 \pm 0.40 \times 10^{-29}$, $3.25 \pm 0.26 \times 10^{-29}$, $1.57 \pm 0.13 \times 10^{-28}$, $2.07 \pm 0.17 \times 10^{-29}$, $1.00 \pm 0.08 \times 10^{-29}$, $3.73 \pm 0.30 \times 10^{-29}$, and $1.42 \pm 0.11 \times 10^{-29} \text{ cm}^2$ have been determined for σ_R for the powders of 1,3-cyclohexanedione (CHD), 3'-aminoacetophenone (AAP), 3'-hydroxyacetophenone (HOAP), diethyl acetamidomalonate (DEAM), isovanillin (ISOV), lactide (DODO), Meldrum's acid (MA), p-tolyl sulfoxide (PTSO), and vanillin (VAN) for the Raman modes at 1174, 714, 1663, 2943, 1671, 657, 641, 796, and 814 cm^{-1} , respectively. The values of Γ for these modes have been determined to be 8.4 ± 0.2 , 4.4 ± 0.1 , 4.5 ± 0.1 , 10.6 ± 0.2 , 4.5 ± 0.1 , 2.7 ± 0.1 , 6.8 ± 0.2 , 5.7 ± 0.2 , and $5.2 \pm 0.1 \text{ cm}^{-1}$, respectively.

ACKNOWLEDGMENTS

The authors wish to thank Dr. T.Y. Fan for his comments regarding this report.

TABLE OF CONTENTS

	Page
Abstract	ii
Acknowledgments	iii
List of Illustrations	v
List of Tables	vii
1. INTRODUCTION	1
2. EXPERIMENTAL	2
3. THEORETICAL	4
4. RESULTS AND DISCUSSION	5
5. SUMMARY	14
References	15

LIST OF ILLUSTRATIONS

Figure No.		Page
1	Schematic of the optical setup used for the measurement of the Raman cross sections.	2
2	Raman spectrum of neat diethyl phthalate (DEP) obtained using 100 mW of 785-nm pump power.	5
3	Raman spectrum of neat dimethyl phthalate (DMP) obtained using 100 mW of 785-nm pump power.	5
4	Raman spectrum of neat ethyl cinnamate (EC) obtained using 100 mW of 785-nm pump power.	6
5	Raman spectrum of neat propylene carbonate (PC) obtained using 100 mW of 785-nm pump power.	6
6	Raman spectrum of neat tripropyl phosphate (TPP) obtained using 100 mW of 785-nm pump power.	7
7	Raman spectrum of 1,3-cyclohexanedione (CHD) powder obtained using 100 mW of 785-nm pump power.	8
8	Raman spectrum of 3'-aminoacetophenone (AAP) powder obtained using 100 mW of 785-nm pump power.	9
9	Raman spectrum of 3'-hydroxyacetophenone (HOAP) powder obtained using 100 mW of 785-nm pump power.	9
10	Raman spectrum of diethyl acetamidomalonate (DEAM) powder obtained using 100 mW of 785-nm pump power.	10
11	Raman spectrum of isovanillin (ISOV) powder obtained using 100 mW of 785-nm pump power.	10
12	Raman spectrum of lactide (DODO) powder obtained using 100 mW of 785-nm pump power.	11

13	Raman spectrum of Meldrum's acid (MA) powder obtained using 100 mW of 785-nm pump power.	11
14	Raman spectrum of p-tolyl sulfoxide (PTSO) powder obtained using 100 mW of 785-nm pump power.	12
15	Raman spectrum of vanillin (VAN) powder obtained using 100 mW of 785-nm pump power.	12

LIST OF TABLES

Table No.		Page
1	Values of the Molecular Mass M and Density ρ Along with CAS Number and Chemical Formula	3
2	Values of the Linewidth Γ (FWHM) and Raman Cross Section σ_R for the 1041-, 1040-, 1638-, 849-, and 726-cm ⁻¹ Modes of Neat DEP, DMP, EC, PC, and TPP, Respectively	8
3	Values of the Linewidth Γ (FWHM), N_R , and Raman Cross Section σ_R for the 1174-, 714-, 1663-, 2943-, 1671-, 657-, 641-, 796-, and 814-cm ⁻¹ Modes of CHD, AAP, HOAP, DEAM, ISOV, DODO, MA, PTSO, and VAN, Respectively	13

1. INTRODUCTION

The values of the absolute Raman cross sections σ_R of solids, liquids, and gases are important for the science and technology of these materials. Values of the absolute σ_R of relatively few materials have been reported. These include benzene [1,2], benzenethiol [3], bis-2-chlorethyl sulfide (HD) [4], carbon monoxide [5], carbon dioxide [5], diamond [6–8], diisopropyl methylphosphonate (DIMP) [4], dimethyl methylphosphonate (DMMP) [4], ethyl dimethylphosphoramidocyanidate (GA) [4], ethyl S-2-diisopropylamino ethylmethylphosphonothioate (VX) [4], gallium phosphide [9,10], hydrogen sulfide [5], isopropyl methylphosphonofluoridate (GB) [4], nitrogen [5], oxygen [5], and silicon [11,12].

Here, we report the measurement of the absolute Raman cross sections of fourteen materials, which substantially expands the number of materials for which such measurements have been made. These are diethyl phthalate (DEP), dimethyl phthalate (DMP), ethyl cinnamate (EC), propylene carbonate (PC), 1,3-cyclohexanedione (CHD), 3'-aminoacetophenone (AAP), 3'-hydroxyacetophenone (HOAP), diethyl acetamidomalonate (DEAM), isovanillin (ISOV), lactide (DODO),¹ Meldrum's acid (MA),¹ p-tolyl sulfoxide (PTSO), tripropyl phosphate (TPP), and vanillin (VAN).

¹ Lactide and Meldrum's acid are synonyms for 3,6-dimethyl-1,4-dioxane-2,5-dione and 2,2-dimethyl-1,3-dioxane-4,6-dione, respectively.

2. EXPERIMENTAL

A schematic of the optical setup for the measurement of Raman cross sections is shown in Figure 1.




Figure 1. Schematic of the optical setup used for the measurement of the Raman cross sections.

The acronym PL is for pump laser. HWP is a half-wave plate for setting the polarization of the pump laser. BS is a beamsplitter for reflecting the pump laser beam towards the sample and for transmitting the backscattered Raman light. L1 is a lens for focusing the pump laser on the sample and for the collimation of the Raman scattered light. LWPFs are long-wave-pass filters for blocking the pump light. L2 is a lens for focusing the Raman light on the entrance slit of the Raman spectrometer (RS).

The pump laser is a 784.9-nm, 1400-mW cw single-mode (both transverse and longitudinal), linearly polarized Sacher Lasertechnik diode laser model SYS-420-0785-1400 (Sacher Lasertechnik, Marburg, Germany/Buena Park, California). The HWP is Edmund Optics part number 46-561 (Edmund Optics, Barrington, New Jersey). The BS is Semrock part number LPD01-785RS-25×36×1.1 with 803–1214 nm passband (Semrock, Rochester, New York). The lens L1 is a 12-mm-diameter, 20-mm-effective-focal-length (EFL), near-infrared (NIR) achromat Edmund Optics part number 45-792. The long-wave-pass filters are 25-mm-diameter Semrock part number LP02-785RE-25 with 790–1770 nm passband. The lens L2 is a 25-mm-diameter, 40-mm-EFL, NIR achromat Edmund Optics part number 45-801. The Raman spectrometer is a 500-mm-focal-length, $f\#$ 6.5, Princeton Instruments model SP-2556, which is equipped with the Princeton Instruments camera PIXIS 100BR (Princeton Instruments, Trenton, New Jersey). The grating in the spectrometer has 1200 grooves/mm. The blaze wavelength of the grating is 750 nm.

All the chemicals used in this work were purchased from Sigma-Aldrich. The values of the molecular mass M and density ρ of these materials are given in Table 1. The values of ρ for the liquids

are obtained from their Material Safety Data Sheets (MSDS). The values of ρ of the powders were deduced from the measured mass and volume contained in the near infrared (NIR) 1.0-mm path length optical cells, Spectrocell part no. R-4001-T (Spectrocell Inc., Orelan, Pennsylvania) with fused silica windows.

Table 1
Values of the Molecular Mass M and Density ρ Along with CAS Number and Chemical Formula

Material	CAS No.	Phase	Chemical Formula	M (g/mol)	ρ (g/cm ³)
Diethyl phthalate (DEP)	84-66-2	Liquid	C ₁₂ H ₁₄ O ₄	222.24	1.12
Dimethyl phthalate (DMP)	131-11-3	Liquid	C ₁₀ H ₁₀ O ₄	194.18	1.19
Ethyl cinnamate (EC)	103-36-6	Liquid	C ₁₁ H ₁₂ O ₂	176.21	1.05
Propylene carbonate (PC)	108-32-7	Liquid	C ₄ H ₆ O ₃	102.09	1.19
Tripropyl phosphate (TPP)	513-08-6	Liquid	C ₉ H ₂₁ O ₄ P	224.23	1.01
1,3-Cyclohexanedione (CHD)	504-02-9	Powder	C ₆ H ₈ O ₂	112.13	0.29
3'-Aminoacetophenone (AAP)	99-03-6	Powder	C ₈ H ₉ NO	135.16	0.54
3'-Hydroxyacetophenone (HOAP)	121-71-1	Powder	C ₈ H ₈ O ₂	136.15	0.50
Diethyl acetamidomalonate (DEAM)	1068-90-2	Powder	C ₉ H ₁₅ NO ₅	217.22	0.74
Isovanillin (ISOV)	621-59-0	Powder	C ₈ H ₈ O ₃	152.15	0.78
Lactide (DODO)	95-96-5	Powder	C ₆ H ₈ O ₄	144.13	0.81
Meldrum's acid (MA)	2033-24-1	Powder	C ₆ H ₈ O ₄	144.13	0.58
p-Tolyl sulfoxide (PTSO)	1774-35-2	Powder	C ₁₄ H ₁₄ OS	230.33	0.39
Vanillin (VAN)	121-33-5	Powder	C ₈ H ₈ O ₃	152.15	0.51

The purity of the chemicals used in this work is as follows: DMP 99.5%, DEP 99.9%, EC 99.7%, PC 99.97%, CHD >99.9%, AAP 99.3%, HOAP 99.9%, DEAM 99.9%, ISOV 99.96%, DODO 99.60%, MA >99.9%, PTSO 98.2%, TPP 99.7%, and VAN 99.9%.

3. THEORETICAL

The Raman cross section σ_R (cm^2) of a Raman mode ν_R is given by

$$\sigma_R = \left(\frac{4\pi}{K} \right) \left(\frac{M}{\rho N_A} \right) \left(\frac{N_R}{N_P} \right), \quad (1)$$

where K is a constant that takes into account the transmittance of the L1 lens (93%) and the fused silica front window of the optical cell (96.5%) for the pump light and the Raman scattered light, and the transmittance of the beamsplitter (98%), four LWPFS (92%), and the L2 lens (93%) for the Raman scattered light. The value of K has been determined to be 0.68 ± 0.02 . M (g/mol) is molecular mass, ρ (g/cm^3) is the density, N_A is the Avogadro constant, N_R is the number of Raman photons ($\text{s}^{-1} \cdot \text{cm}^{-1} \cdot \text{sr}^{-1} \cdot \text{mW}^{-1}$) integrated over the Raman mode, and N_P is the number of pump photons ($\text{s}^{-1} \cdot \text{mW}^{-1}$) equal to 3.96×10^{15} . Using the values of K , N_A , and N_P , Eq. (1) may be written as

$$\sigma_R = 7.75 \pm 0.11 \times 10^{-39} \left(\frac{MN_R}{\rho} \right). \quad (2)$$

We neglect the effect of the elastic scattering of the pump light on the measurement of the Raman cross section.

4. RESULTS AND DISCUSSION

Figures 2 through 6 show the Raman spectra (Raman signal/data point versus Raman shift) of neat DEP, DMP, EC, PC, and TPP. Data-point spacing is 0.42 to 0.20 cm^{-1} for the 200 to 3200 cm^{-1} Raman shift.




Figure 2. Raman spectrum of neat diethyl phthalate (DEP) obtained using 100 mW of 785-nm pump power.




Figure 3. Raman spectrum of neat dimethyl phthalate (DMP) obtained using 100 mW of 785-nm pump power.




Figure 4. Raman spectrum of neat ethyl cinnamate (EC) obtained using 100 mW of 785-nm pump power.




Figure 5. Raman spectrum of neat propylene carbonate (PC) obtained using 100 mW of 785-nm pump power.




Figure 6. Raman spectrum of neat tripropyl phosphate (TPP) obtained using 100 mW of 785-nm pump power.

All the Raman spectra were measured using 100 mW of the pump power incident upon the lens L1. Each spectrum is the average of three scans for the 125 to 3200 cm^{-1} range, which consists of nine segments. Each segment is the average of 20 frames with each frame obtained with an integration time of 1.0 s. Thus, it takes 9 minutes for the acquisition of each spectrum. Using the 785-nm pump laser, the spectral resolution of the Raman spectrometer (FWHM) was determined to be $2.0 \pm 0.1 \text{ cm}^{-1}$.

The values of N_R for the 1041-, 1040-, 1638-, 849-, and 726- cm^{-1} modes of DEP, DMP, EC, PC, and TPP are $1.25 \pm 0.06 \times 10^7$, $1.98 \pm 0.10 \times 10^7$, $2.38 \pm 0.12 \times 10^8$, $1.76 \pm 0.09 \times 10^7$, and $8.87 \pm 0.44 \times 10^6$ photons ($\text{s}^{-1} \cdot \text{cm}^{-1} \cdot \text{sr}^{-1} \cdot \text{mW}^{-1}$), respectively; the values of N_R were obtained after subtracting the background. The values of σ_R have been determined by substituting the values of M , N_R , and ρ in Eq. (2). The accuracy of σ_R for neat DEP, DMP, EC, PC, and TPP is estimated to be $\pm 6\%$, which is due to the $\pm 5\%$, $\pm 3\%$, and $\pm 1.0\%$ uncertainty in the values of N_R , K , and ρ , respectively. The values of the linewidth Γ (full width at half maximum) and σ_R for neat DEP, DMP, EC, PC, and TPP are given in Table 2.

Figures 7 through 15 show the Raman spectra of the CHD, AAP, HOAP, DEAM, ISOV, DODO, MA, PTSO, TPP, and VAN powders.

Table 2

Values of the Linewidth Γ (FWHM) and Raman Cross Section σ_R for the 1041-, 1040-, 1638-, 849-, and 726- cm^{-1} Modes of Neat DEP, DMP, EC, PC, and TPP, Respectively

Material	Mode (cm^{-1})	Γ (cm^{-1})	σ_R (cm^2)
DEP	1041	6.6 ± 0.1	$1.92 \pm 0.12 \times 10^{-29}$
DMP	1040	7.0 ± 0.1	$2.50 \pm 0.15 \times 10^{-29}$
EC	1638	11.3 ± 0.2	$3.10 \pm 0.19 \times 10^{-28}$
PC	849	15.8 ± 0.2	$1.17 \pm 0.07 \times 10^{-29}$
TPP	726	29.0 ± 0.3	$1.53 \pm 0.09 \times 10^{-29}$




Figure 7. Raman spectrum of 1,3-cyclohexanedione (CHD) powder obtained using 100 mW of 785-nm pump power.




Figure 8. Raman spectrum of 3'-aminoacetophenone (AAP) powder obtained using 100 mW of 785-nm pump power.




Figure 9. Raman spectrum of 3'-hydroxyacetophenone (HOAP) powder obtained using 100 mW of 785-nm pump power.




Figure 10. Raman spectrum of diethyl acetamidomalonate (DEAM) powder obtained using 100 mW of 785-nm pump power.




Figure 11. Raman spectrum of isovanillin (ISOV) powder obtained using 100 mW of 785-nm pump power.




Figure 12. Raman spectrum of lactide (DODO) powder obtained using 100 mW of 785-nm pump power.




Figure 13. Raman spectrum of Meldrum's acid (MA) powder obtained using 100 mW of 785-nm pump power.




Figure 14. Raman spectrum of *p*-tolyl sulfoxide (PTSO) powder obtained using 100 mW of 785-nm pump power.




Figure 15. Raman spectrum of vanillin (VAN) powder obtained using 100 mW of 785-nm pump power.

The large background in the Raman spectra of AAP and ISOV in Figures 8 and 11, respectively, is most likely due to fluorescence. The values of N_R were obtained after subtracting the background. The accuracy of σ_R of the powders is estimated to be $\pm 8\%$, which is due to the $\pm 5\%$, $\pm 3\%$, and $\pm 5.0\%$

uncertainty in the values of N_R , K , and ρ , respectively. The effect of the elastic scattering of the pump beam, neglected in this work, may substantially decrease the estimated accuracy of σ_R . Also, the values of σ_R obtained in this work represent the lower bound because of the neglect of the elastic scattering of the pump beam. The values of Γ , N_R , and σ_R for CHD, AAP, HOAP, DEAM, ISOV, DODO, MA, PTSO, and VAN are given in Table 3.

Table 3

Values of the Linewidth Γ (FWHM), N_R , and Raman Cross Section σ_R for the 1174-, 714-, 1663-, 2943-, 1671-, 657-, 641-, 796-, and 814- cm^{-1} Modes of CHD, AAP, HOAP, DEAM, ISOV, DODO, MA, PTSO, and VAN, Respectively

Material	Mode (cm^{-1})	Γ (cm^{-1})	N_R (Photons/s.cm.sr.mW)	σ_R (cm^2)
CHD	1174	8.4 ± 0.2	$4.90 \pm 0.25 \times 10^7$	$1.47 \pm 0.12 \times 10^{-28}$
AAP	714	4.4 ± 0.1	$8.63 \pm 0.42 \times 10^6$	$1.67 \pm 0.13 \times 10^{-29}$
HOAP	1663	4.5 ± 0.1	$2.44 \pm 0.12 \times 10^7$	$5.15 \pm 0.40 \times 10^{-29}$
DEAM	2943	10.6 ± 0.2	$1.43 \pm 0.07 \times 10^7$	$3.25 \pm 0.26 \times 10^{-29}$
ISOV	1671	4.5 ± 0.1	$1.04 \pm 0.05 \times 10^8$	$1.57 \pm 0.13 \times 10^{-28}$
DODO	657	2.7 ± 0.1	$1.50 \pm 0.08 \times 10^7$	$2.07 \pm 0.17 \times 10^{-29}$
MA	641	6.8 ± 0.2	$5.17 \pm 0.26 \times 10^6$	$1.00 \pm 0.08 \times 10^{-29}$
PTSO	796	5.7 ± 0.2	$8.15 \pm 0.41 \times 10^6$	$3.73 \pm 0.30 \times 10^{-29}$
VAN	814	5.2 ± 0.1	$6.16 \pm 0.31 \times 10^6$	$1.42 \pm 0.11 \times 10^{-29}$

5. SUMMARY

We have measured the values of the absolute Raman cross sections of several chemicals for 785-nm excitation with an accuracy of $\pm 9\%$ for the liquids and $\pm 13\%$ for the powders. The measured values of the Raman cross sections are of the order of 10^{-28} to 10^{-29} cm².

REFERENCES

- [1] M.O. Trulson and R.A. Mathies, “Raman cross section measurements in the visible and ultraviolet using an integrating cavity: Application to benzene, cyclohexane, and cacodylate,” *J. Chem. Phys.* 1986, 84 (4): 2068–2074.
- [2] K.T. Schomaker, J.K. Delaney, and P.M. Champion, “Measurements of the Raman cross section of benzene,” *J. Chem. Phys.* 1986, 85 (8): 4240–4247.
- [3] R.L. Aggarwal, L.W. Farrar, E.D. Diebold, and D.L. Polla, “Measurement of the absolute Raman scattering cross section of the 1584-cm⁻¹ band of benzenethiol and the surface-enhanced Raman scattering cross section enhancement factor for femtosecond laser-nanostructured substrates,” *J. Raman Spectrosc.* 2009, 40 (9): 1331–1333.
- [4] S.D. Christensen, “Raman cross sections of chemical agents and simulants,” *Appl. Spectrosc.* 1988, 42 (2): 318–321.
- [5] D.G. Fouche and R.K. Chang, “Relative Raman cross section for N₂, O₂, CO, CO₂, SO₂, and H₂S,” *Appl. Phys. Lett.* 1971, 18 (12): 579–580.
- [6] A.K. McQuillan, W.R.L. Clements, and B.P. Stoicheff, “Stimulated Raman emission in diamond: Spectrum, gain, and angular distribution,” *Phys. Rev. A* 1970, 1 (3): 628–635.
- [7] M.H. Grimsditch and A.K. Ramdas, “Brillouin scattering in diamond,” *Phys. Rev. B* 1975, 11 (8): 3139–3148.
- [8] R.L. Aggarwal, L.W. Farrar, S.K. Saikin, X. Andrade, A. Aspuru-Guzik, and D.L. Polla, “Measurement of the absolute Raman cross section of the optical phonons in type Ia natural diamond,” *Solid State Commun.* 2012, 152 (3): 204–209.
- [9] J.M. Calleja, H. Vogt, and M. Cardona, *J. Phys. Colloq. C* 1981, 6: 487; *Phil. Mag. A* 1982, 45: 239.
- [10] R.L. Aggarwal, L.W. Farrar, and D.L. Polla, “Measurement of the absolute Stoke Raman cross sections of the longitudinal optical (LO) phonons of room-temperature GaP,” *Solid State Commun.* 2009, 149 (33): 1330–1332.
- [11] M. Grimsditch and M. Cardona, “Absolute cross-section for Raman scattering by phonons in silicon,” *Phys. Status Solidi B* 1980, 102 (1): 155–161.
- [12] R.L. Aggarwal, L.W. Farrar, S.K. Saikin, A. Aspuru-Guzik, and D.L. Polla, “Measurement of the absolute Raman cross section of the optical phonon in silicon,” *Solid State Commun.* 2011, 151 (7): 553–556.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 13 September 2013		2. REPORT TYPE Technical Report		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Measurement of the Absolute Raman Cross Sections of Diethyl Phthalate, Dimethyl Phthalate, Ethyl Cinnamate, Propylene Carbonate, Tripropyl Phosphate, 1,3-Cyclohexanedione, 3'-Aminoacetophenone, 3'-Hydroxyacetophenone, Diethyl Acetamidomalonate, Isovanillin, Lactide, Meldrum's Acid, p-Tolyl Sulfoxide, and Vanillin				5a. CONTRACT NUMBER FA8721-05-C-0002	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) R.L. Aggarwal, L.W. Farrar, B.G. Saar, T.H. Jeys, and R.B. Goodman				5d. PROJECT NUMBER 1567	
				5e. TASK NUMBER 610	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) MIT Lincoln Laboratory 244 Wood Street Lexington, MA 02420-9108				8. PERFORMING ORGANIZATION REPORT NUMBER TR-1175	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Joint Program Executive Office for Chemical and Biological Defense JPM NBC CA 2800 Bush River Rd. Aberdeen Proving Ground, MD 21010-5424				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We have measured the Raman spectra of several chemicals for 785-nm excitation using a Raman spectrometer with focal length of 500 mm, f-number of 6.5, and resolution of $2.0 \pm 0.1 \text{ cm}^{-1}$. Values of $1.92 \pm 0.12 \times 10^{-29}$, $2.50 \pm 0.15 \times 10^{-29}$, $3.10 \pm 0.19 \times 10^{-28}$, $1.17 \pm 0.07 \times 10^{-29}$, and $1.53 \pm 0.09 \times 10^{-29} \text{ cm}^2$ have been determined for the Raman cross section σ_R of neat Diethyl phthalate (DEP), dimethyl phthalate (DMP), ethyl cinnamate (EC), propylene carbonate (PC), and tripropylene phosphate (TPP) for the Raman modes at 1041, 1040, 1638, 849, and 726 cm^{-1} , respectively. The values of the linewidth (FWHM) Γ for these modes have been determined to be 6.6 ± 0.1 , 7.0 ± 0.1 , 11.3 ± 0.2 , 15.8 ± 0.2 , and $29.0 \pm 0.3 \text{ cm}^{-1}$, respectively. Values of $1.47 \pm 0.12 \times 10^{-28}$, $1.67 \pm 0.13 \times 10^{-29}$, $5.15 \pm 0.40 \times 10^{-29}$, $3.25 \pm 0.26 \times 10^{-29}$, $1.57 \pm 0.13 \times 10^{-28}$, $2.07 \pm 0.17 \times 10^{-29}$, $1.00 \pm 0.08 \times 10^{-29}$, $3.73 \pm 0.30 \times 10^{-29}$, and $1.42 \pm 0.11 \times 10^{-29} \text{ cm}^2$ have been determined for σ_R for the powders of 1,3-cyclohexanedione (CHD), 3'-aminoacetophenone (AAP), 3'-hydroxyacetophenone (HOAP), diethyl acetamidomalonate (DEAM), isovanillin (ISOV), lactide (DODO), Meldrum's acid (MA), p-tolyl sulfoxide (PTSO), and vanillin (VAN) for the Raman modes at 1174, 714, 1663, 2943, 1671, 657, 641, 796, and 814 cm^{-1} , respectively. The values of Γ for these modes have been determined to be 8.4 ± 0.2 , 4.4 ± 0.1 , 4.5 ± 0.1 , 10.6 ± 0.2 , 4.5 ± 0.1 , 2.7 ± 0.1 , 6.8 ± 0.2 , 5.7 ± 0.2 , and $5.2 \pm 0.1 \text{ cm}^{-1}$, respectively.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as report	18. NUMBER OF PAGES 24	19a. NAME OF RESPONSIBLE PERSON
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code)

