

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 20-07-2012		2. REPORT TYPE Research		3. DATES COVERED (From - To) 1 September 2009 - 30 June 2012	
4. TITLE AND SUBTITLE Murine model of repeated exposures to conspecific trained aggressors simulates features of post-traumatic stress disorder				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 604110	
				5d. PROJECT NUMBER 374	
6. AUTHOR(S) Hammamieh, Rasha, Chakraborty, Nabarun, De Lima, Thereza C.M., Meyerhoff, James, Gautam, Aarti, Muhie, Seid, D'Arpa, Peter, and Lumley, Lucille				5e. TASK NUMBER GJ	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Center for Environmental Health Research, 568 Doughten Drive, Fort Detrick, MD 21702-5010; Federal University of Santa Catarina, Department of Pharmacology, Florianopolis, SC, Brazil; US Army Medical Research Institute of Chemical Defense, APG, MD; Naval Medical Research Center, Silver Spring MD				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command 504 Scott Street Fort Detrick, MD 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S) USAMRMC	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for Public Release; Unlimited Distribution.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We evaluated repeated exposures of mice to a trained aggressor mouse as a model (adapted from "social stress" models of traumatic stress) for aspects of post-traumatic stress disorder (PTSD). Using a "cagewithin-cage resident-intruder" protocol, subject C57BL/6J mice were exposed to aggressors for 6 h daily for 5 or 10 days. At one to three random times during each 6-h session, subjects were exposed directly to aggressor for 1 min or 10 bites, whichever came first. Behavioral, physiological, and histological changes associated with aggressor-exposure were assessed for up to 6 weeks. During aggressor exposure, subjects displayed less territorial behavior, gained weight, and increased body temperature. One day after the last aggressor exposure, inflammatory cardiac histopathologies were prevalent; after 10 days, only mild myocardial degeneration with fibrosis or fibroplasias was evident, while controls showed almost no cardiac abnormalities at any time. After 4 weeks, the medial prefrontal cortex of control mice showed increased dendritic spine density, but aggressor-exposed mice showed no increase. Behaviors affected by aggressor exposure were evaluated in a partition test wherein the subject mouse is separated from the aggressor by a fenestrated partition that permits sensory					
15. SUBJECT TERMS PTSD, stress effects, avoidance, dendritic spines, cardiovascular disease					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Carol O'Brien
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (Include area code) 301-619-7685

Reset



Research report

Murine model of repeated exposures to conspecific trained aggressors simulates features of post-traumatic stress disorder

Rasha Hammamieh^{a,*}, Nabarun Chakraborty^{a,1}, Thereza C.M. De Lima^b, James Meyerhoff^a, Aarti Gautam^a, Seid Muhie^a, Peter D'Arpa^a, Lucille Lumley^c, Erica Carroll^d, Marti Jett^a

^a US Army Center for Environmental Health Research, Fort Detrick, MD, USA

^b Federal University of Santa Catarina – Department of Pharmacology, Florianopolis, SC, Brazil

^c US Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, Aberdeen, MD, USA

^d Naval Medical Research Center, Silver Spring, MD, USA

HIGHLIGHTS

- ▶ We established a modified social stress mouse model adapting and revising the 'resident-intruder' model.
- ▶ The model used SJL strain mouse as the aggressor and C57BL/6J as the subject mouse.
- ▶ The aggressor-exposed stress inflicted PTSD-like negative behavioral traits on the subject mice.
- ▶ Stress inflicted physiological alteration and developed neurological atrophy.
- ▶ Stress specific cardiac myopathy and lymphohistiocytic myocarditis was observed.

ARTICLE INFO

Article history:

Received 1 February 2012

Received in revised form 7 July 2012

Accepted 12 July 2012

Available online 20 July 2012

Keywords:

PTSD

Stress effects

Avoidance

Dendritic spines

Cardiovascular disease

ABSTRACT

We evaluated repeated exposures of mice to a trained aggressor mouse as a model (adapted from "social stress" models of traumatic stress) for aspects of post-traumatic stress disorder (PTSD). Using a "cage-within-cage resident-intruder" protocol, subject C57BL/6J mice were exposed to aggressors for 6 h daily for 5 or 10 days. At one to three random times during each 6-h session, subjects were exposed directly to aggressor for 1 min or 10 bites, whichever came first. Behavioral, physiological, and histological changes associated with aggressor-exposure were assessed for up to 6 weeks. During aggressor exposure, subjects displayed less territorial behavior, gained weight, and increased body temperature. One day after the last aggressor exposure, inflammatory cardiac histopathologies were prevalent; after 10 days, only mild myocardial degeneration with fibrosis or fibroplasias was evident, while controls showed almost no cardiac abnormalities at any time. After 4 weeks, the medial prefrontal cortex of control mice showed increased dendritic spine density, but aggressor-exposed mice showed no increase. Behaviors affected by aggressor exposure were evaluated in a partition test wherein the subject mouse is separated from the aggressor by a fenestrated partition that permits sensory cues to pass but prevents direct physical interaction. For up to 4–6 weeks after the last aggressor exposure, subjects showed prolonged grooming, freezing, retarded locomotion and no tail rattling. PTSD and its co-morbidities are often consequent to repeated aggravated "social" assaults (e.g., combat) and manifest socially over time, suggesting the relevance of this repeated aggressor-exposure model to clinical aspects of PTSD.

Published by Elsevier B.V.

1. Introduction

Resurgent interest in post-traumatic stress disorder (PTSD) pathological mechanisms and medical countermeasures has been

stimulated by the substantial increase in the 2-year prevalence rate associated with the start of the Iraq War and the high incidence of PTSD in U.S. veterans of recent wars in Iraq and Afghanistan. Between 2002 and 2008, a study of 289,328 U.S. veterans who were first-time users of VA health care after military service showed 22% diagnosed with PTSD and 17% diagnosed with depression [1]. Notably, greater combat exposure has been associated with higher risk for PTSD [1], confirming earlier findings [2–4].

Significant controversy in the diagnosis of PTSD [5] has emphasized a pressing need to identify and validate disease-specific

* Corresponding author at: 568 Doughten Drive, Fort Detrick, MD 21702, USA. Tel.: +1 3016192338; fax: +1 3016197606.

E-mail address: Rasha.Hammamieh1@us.army.mil (R. Hammamieh).

¹ The authors contributed equally to this article.

diagnostic biomarkers. A number of animal models of PTSD have been investigated. While no animal model can be expected to fully capture the complexity of human cognitive function in psychiatric disorders, the shared basic emotional processes of humans and other mammals suggests that animal models can capture core endophenotypes of psychiatric disorders [6,7].

Currently, the clinical diagnosis of PTSD rests largely on reported symptoms and reported history. The essential features of PTSD as listed in DSM-IV (309.81) include the development of characteristic symptoms following an event that is an extreme traumatic stressor, which involves threatened death or serious injury to the self, or the witnessing or learning of actual death or threatened death or serious injury happening to others who are family members or close associates (Criterion A1). The traumatic event(s) must elicit intense fear, helplessness, or horror (Criterion A2) and produce symptoms that include persistent re-experiencing of the traumatic event(s) (Criterion B), persistent avoidance of stimuli associated with the trauma and numbing of general responsiveness (Criterion C), and persistent symptoms of increased arousal (Criterion D). These symptoms must be present for more than 1 month (Criterion E) and cause clinically significant distress or impairment in social, occupational, or other important areas of functioning (Criterion F) [8].

In mice, a wide variety of physiological and psychogenic stresses, including so-called “social stress,” restraint stress [9], foot shock [10], juvenile abuse [11], and artificial alteration of stress hormones like corticosterone [12] have been shown to produce behaviors reminiscent of PTSD. In “social stress” models, subordinate mice have been subjected to the inescapable exposure to: another mouse made aggressive by isolation and/or training [13,14]; a predator [15]; a predator’s odor from sources such as skin, fur, urine or feces [16]; or, synthetic odors derived from a predator’s feces or anal glands [17].

The quality of an animal model has been assessed using the criteria of face validity, predictive validity, and construct validity—i.e., how closely the model represents symptoms, efficacy of treatments, and cellular and molecular processes, respectively, of the human disease (discussed in [6]). *Face validity* of models for representing symptoms of psychiatric diseases is typically an assumption of validity. For example, fear of novelty and unprotected areas are assumed to represent anxiety, and the changed behaviors in the Learned Helplessness Model are described as depression-like symptoms. *Predictive validity* refers to the quality of a model for showing that drugs with efficacy in relieving human symptoms mitigate the animal behaviors that correlate with the human psychiatric symptoms (e.g., effects of antidepressants in Forced Swimming Test and Tail Suspension Test (as discussed in [6])). *Construct validity* refers to the model’s representation of cellular and molecular processes of the human disorder.

Our systems biology approach aims to ultimately identify objective diagnostic criteria for PTSD. Here we present initial studies of a mouse model that employs “cage-within-cage resident-intruder” exposures of subject mice to a trained aggressor mouse for 6 h daily for 5 or 10 days. At one to three random times during each 6-h session, subject mice are removed from the cage-within-cage and put into direct physical contact with the aggressor in its home cage for 1 min or 10 bites, whichever comes first. We chose this model of multiple stressor exposures in order to model the stress of the unpredictable threats of daily trauma encountered by warfighters.

During the schedules of exposure of subject mice to the aggressor and over the course of the recovery of the subject mice in their home cages, we evaluated behavioral, physiological, and histological alterations. After significant recovery periods, a number of the acute stress effects, including histological and neurological pathologies and behaviors interpreted to represent human fear

and anxiety, were persistent in aggressor-exposed mice, modeling some features of PTSD and co-morbidities.

2. Material and methods

2.1. Mice

2.1.1. Aggressor mice

The SJL albino male mice (6-week old and weighing 30–35 g) were housed individually in polycarbonate cages (48 cm × 27 cm × 20 cm) for 1 month prior to the experiment to induce aggressiveness due to isolation. They were then trained to assault intruders by occasional pairing with olfactory bulbectomized (OBx) male C57BL/6J mice. All mice had free access to food and water and were kept in a temperature-controlled room (21 ± 2 °C) on reverse 12/12 h light/dark cycle (lights off at 06:00 AM).

2.1.2. Subject mice

Male C57BL/6J mice (6-week old weighing 20–25 g) were single housed under the same environmental conditions as the aggressor mice in a different room for 1 week before initiating the experiments. Subject mice were kept in a separate room from aggressor mice. All animal experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at the Walter Reed Army Institute of Research and the Medstar Research Institute and were performed in a facility accredited by the Association for the Assessment and Accreditation of Laboratory Animal Care International (AAALAC). All mice were purchased from Jackson Laboratory, ME, USA.

2.2. Aggressor exposure

Aggressor exposure (Agg-E) sessions followed a modified “cage-within-cage resident-intruder” protocol in which experimental mice in a wire mesh cage (17.5 cm × 14 cm × 7.5 cm) were placed inside the aggressor’s large plastic cage (48 cm × 27 cm × 20 cm) for 6 h (Fig. 1A). The 6-h sessions were repeated daily for either 5 or 10 consecutive days, as shown in the protocol timeline (Fig. 1B). Control mice were kept inside the same type of wired mesh box inside the same type of larger cage with fresh bedding for 6 h without direct exposure to an aggressor on the same daily schedule. Control and experimental mice were deprived of food and water during the 6-h “cage-within-cage” sessions, while the aggressor mice were provided food and water ad libitum. At the end of each 6-h session, control and experimental mice were returned to their respective home cage with food and water available ad libitum until the next session. At one to three random times during the 6-h session, subject mice were removed from the wire mesh cage and exposed directly to the aggressor mouse for 1 min or until 10 bites were inflicted on the subject mouse, whichever came first. On average, subject mice received 10 bites within 48 s.

2.3. Physiological evaluations

2.3.1. Body weights and temperatures

Body weights and temperatures were recorded before and after each 6-h Agg-E session. The core body temperatures of the mice were recorded using Electronic ID Transponders™ (IPTT-300, Bio Medic Data Systems Inc., DE) subcutaneously implanted into the dorsal cervical region 3 days in advance of the first 6-h stress session. Body weights were measured by placing the mice on a standard laboratory scale (Sartorius, Edgewood, NY).

2.3.2. Spleen blood cell counts

Following the partition test, after euthanasia via cervical dislocation of five control and five Agg-E mice, spleen tissue was collected aseptically and transferred into cold media with antibiotics. Immediately, the tissue was mashed through a cell strainer (BD, Inc., MD) and centrifuged at 800 × g. The pellet was resuspended in 1 mL ACK lysis buffer (Lonza, VA) and incubated at room temperature for 5–10 min. Nine milliliters of RPMI-1640 was added followed by centrifugation and resuspension in RPMI-1640 media. A 200 µL aliquot was used for complete cell count using ADVIA® 120 Hematology System (Siemens Healthcare Diagnostics, Inc., NY).

2.4. Cardiac and neurohistological evaluations

2.4.1. Heart histopathology

Heart samples were collected from five control and five Agg-E euthanized mice and perfused in ice-cold 4% paraformaldehyde. Tissues were then stained, sliced, and mounted, and a board-certified veterinary pathologist, blinded from the animal condition, identified and analyzed the samples using brightfield optics. Scoring was between one and four; where one = minimum and four = marked. Primary characteristics evaluated were arterial thrombus, myocardial degeneration and infiltration, and lymphohistiocytic epicarditis, myocarditis and vasculitis.

2.4.2. Neurohistology

The brains were carefully removed from the skulls of euthanized mice, impregnated with Golgi-Cox staining solution, and processed following the protocol established by FD Neurotech, Inc. (Ellicott City, MD). Briefly, the impregnated brains




Fig. 1. Cage-within-cage configuration for aggressor exposure (Agg-E) sessions. (A) The home cage of the aggressor mouse (albino SJL) contains a small mesh cage where the subject mouse (C57BL/6J) is placed during each 6-h Agg-E session. At one to three random times during each 6-h session, the smaller cage is removed and the subject mouse is put into direct contact with the aggressor for 1 min or until the subject receives 10 bites, whichever comes first. The subject mouse is deprived of food and water while the aggressor is permitted free access to food and water during the 6-h session. The control mice had similar set up during 6-h sessions, but without an aggressor. (B) The study time line shows the 5- or 10-day aggressor exposure schedules (Y axis) when body weight and temperature and territorial urine markings were recorded. Before, during and after the 5- and 10-day Agg-E schedules, subject mice were single-housed in home cages (when not in a 6-h session). The dots indicate the times when the listed analyses were performed and mice were sacrificed (at 1 day and 1.5 weeks after the 5-day Agg-E schedule; and, at 1 day, 4 weeks, and/or 6 weeks after the 10-day Agg-E schedule).

were serially sectioned at a thickness of 100 μm . Every third cryostat section (intervals of 300 μm) was mounted on gelatin-coated slides, stained, dehydrated in ethanol, cleared in xylene, and cover-slipped in Permount. The spine density of mPFC was assessed using the Camera Lucida[®] technique as described by Wellman and Kolb [18,19] and performed at Sinq System, Inc. (Columbia, MD). The brain region of interest, the mPFC, defined as in Van de Werd et al., consisted of three sub-regions: infralimbic, prelimbic, and anterior cingulate cortices [20]. At least 36–40 neurons selected satisfied criteria of having a distinguishable (single) cell soma, and were located in such a way as to have a relatively large portion of their dendritic tree available for analysis.

2.5. Behavioral evaluations

2.5.1. Partition test ethogram evaluation (freezing, grooming, partition avoidance, and tail rattling)

We performed ethogram evaluations at 1 day and 1.5 weeks after the 5-day schedule and 1 day, 4 weeks, and 6 weeks after the 10-day schedule using a 5-min partition test [21,22], 1 h prior to euthanasia. The aggressor home cage (48 cm $\times$ 27 cm $\times$ 20 cm) was bisected with a plastic fenestrated partition (1 cm² holes) that permits the passage of sensory cues but prevents direct physical contact. The subject mouse was placed on the opposite side of the partition from the aggressor and videoed for the entire 5 min test. The videos were analyzed with Ethovision XT v.7 software (Noldus[®], Leesburg, VA, USA) using 15 samples/s, dynamic subtraction detection, object always darker than background, erosion and dilation filters of one pixel, and one sample interval for averaging filter. Grooming duration was taken as the total time spent licking the paws or washing the nose, face and other body parts. Tail rattling was observed as rapid vibration of the tail or vigorous tapping of the tail on the floor when facing the partition.

2.5.2. Urine marking test

Blotter papers (0.8 mm thickness) were placed under the cages of control and Agg-E mice during the 6-h cage-within-cage sessions. At the end of each day, the papers were collected and UV-scanned using Molecular Imager Fx[®] (BioRad, Hercules, CA). The number of urination marks and the areas contouring the urine markings were measured using Quantity One[®] software (BioRad, Hercules, CA). The process was repeated on alternate days until the end of the 10-day Agg-E schedule.

2.5.3. Statistical analysis

Statistical analyses were performed using GraphPad[®] version 5.0 software (GraphPad Software, Inc., CA). Experimental results were expressed as mean $\pm$ SEM. Linear regression was used to evaluate body weight, temperature, and urine

markings. Unpaired *t*-test with Welch's correction was used to assess differences between control and Agg-E mice at specific time points after the 5-day and 10-day schedules. Two-way ANOVA, followed by Bonferroni post-test, was used to identify interactions between control or Agg-E mouse behaviors with home-cage rest time.

3. Results

Each subject C57BL/6J mouse was exposed to an SJL albino male trained aggressor mouse for 6 h daily for 5 or 10 days using a modified "cage-within-cage resident-intruder" protocol (Fig. 1A). At one to three random times during each 6-h session, the subject was removed from its cage and exposed directly to the aggressor for 1 min or 10 bites, whichever came first. The aggressor delivered an average of 10 bites to the subject within 48 (± 22.5) s and the latency of first attack was less than 5 (± 1.0) s. Throughout the course of the Agg-E schedules, we recorded body weights and temperatures, and territorial urine markings. One day after the Agg-E schedules and during up to 6 weeks of home cage rest, we assessed changes in physiological, histological and behavioral parameters of Agg-E as compared to control mice (Fig. 1B). Control mice were confined within the cage-within-cage environment with fresh bedding, no aggressor present, and deprived of food and water, on the same 5- or 10-day schedule as the Agg-E mice.

3.1. Body weights and temperatures

Body weights and temperatures were recorded daily prior to each 6-h "cage-within-cage" session (Fig. 2). The average *body weights* of both control and Agg-E groups increased over the course of the 10-day schedule, but Agg-E mice gained significantly more weight (Fig. 2A; $p < 0.001$, slopes of the linear regression lines significantly differed). *Body temperature* of the Agg-E mice was significantly elevated over control mice (Fig. 2B; $p < 0.01$, linear regression).




Fig. 2. Body weight and body temperature are increased during 10 days of Agg-E. Body weights and body temperatures of individual mice were normalized by their own baseline levels before the start of the 10-day schedule. Mean $\pm$ SEM values are plotted ($n = 16$ – 21 mice per group). (A) *Body weight*: linear regression slopes of body weights of Agg-E (closed circles) and control (open circles) are different ($p < 0.05$), and Agg-E body weights differ from controls on days 5, 6, 9 and 10 (unpaired t -test with Welch's correction; * $p < 0.05$; *** $p < 0.001$). (B) *Body temperature* of Agg-E (closed circles) and controls (open circles). Linear regression elevated differently (unpaired t -test with Welch's correction; $p = 0.01$).

3.2. Territorial urine marking

Urine markings were recorded on days 1, 3, 5, 7 and 10 of the 10-day schedule (Fig. 3). After ranking the urine markings by area, the largest 90% were counted. Territorial urine markings of Agg-E mice were initially three-fold lower than control mice ($p = 0.03$, unpaired t -test with Welch's correction) on day 1 and remained lower over the 10-day schedule ($p = 0.02$, linear regression slopes differed), being significantly lower than control mice on day 10 ($p = 0.05$, unpaired t -test with Welch's correction).

3.3. Spleen blood cell counts

We measured blood cells in spleen tissue at 1 day after the 5- and 10-day schedules. Red blood cells (RBC), white blood cells (WBC), and platelets of the 5-day Agg-E mice were significantly increased compared to control mice (Fig. 4A–C; $p < 0.01$ for RBC and WBC, $p < 0.05$ for platelets; unpaired t -test with Welch's correction), as was hematocrit (not shown). After the 10-day Agg-E schedule, these blood cell types also appeared to be increased in Agg-E mice, but not to statistically significant levels. Additionally, basophils of Agg-E mice after the 5-day schedule were increased ($p < 0.01$, unpaired t -test with Welch's correction), and after the 10-day Agg-E schedule trended toward elevation as well ($p = 0.07$, unpaired t -test with Welch's correction) (Fig. 4D).




Fig. 3. Territorial urine marking of Agg-E mice is reduced during the Agg-E schedule. The largest 90% of urine markings were counted and plotted $\pm$ SEM on days 1, 3, 5, 7 and 10 of the schedule (unpaired t -test with Welch's correction; * $p < 0.05$; $n = 5$ mice per group). The slopes of the linear regression lines are also significantly different ($p < 0.05$).

3.4. Heart histopathology

Most of the Agg-E mice exhibited myocardial degeneration and/or lymphohistiocytic myocarditis 1 day after the 5-day (in six of eight mice) and 10-day (in eight of 11 mice) Agg-E schedules (Table 1). More mice of the 10-day Agg-E group showed myocardial degeneration as compared to the 5-day Agg-E group at this 1 day point. None of the 28 control mice showed myocardial degeneration or lymphohistiocytic myocarditis at any time (control data not shown). Additionally at the 1 day point, mice of both Agg-E schedules exhibited fibrinoid and/or lymphohistiocytic vasculitis (four of eight of 5-day Agg-E; and five of 11 of 10-day Agg-E). Again, none of the control mice exhibited either of these histopathologies at any time. Following the home cage rest periods, the inflammatory histopathologies had disappeared, but myocardial degeneration with fibrosis or fibroplasia persisted. More mice of the 5-day Agg-E group showed myocardial degeneration after 1.5 weeks of home cage rest than did mice of the 10-day Agg-E group after their longer 4-week rest; however, more mice of the 10-day Agg-E group showed myocardial degeneration at the 1 day point.

3.5. Dendritic spine density in the medial prefrontal cortex

The dendritic spine density of pyramidal neurons of the medial prefrontal cortex (mPFC) was evaluated at 1 day and 4 weeks after the 10-day Agg-E schedule (Fig. 5). At 1 day following repeated exposures, control and Agg-E mice showed no difference in dendritic spine density. Four weeks later, the dendritic spine density of control mice had increased significantly by 26% ($p < 0.01$) compared to the 1 day data, while the Agg-E mice showed only a 9% increase that was not significant. Accordingly, mPFC dendritic spine density of control mice tended to be greater than Agg-E mice at the 4-week point ($p = 0.085$, unpaired t -test with Welch's correction).

3.6. Behaviors in the partition test

Agg-E and control mice behaviors were evaluated in a partition test in which subject mice are placed in the aggressor home cage separated from the aggressor by a fenestrated partition that allows transmission of sensory information but prevents direct physical contact (Fig. 6). We evaluated freezing, avoidance of the aggressor, tail rattling (Fig. 7) and locomotion (Supplementary Fig. S1) in the partition zone, and grooming and total distance traveled in the entire subject side of the cage, during the 5-min partition test. The Agg-E and control group values for these behaviors, except for




Fig. 4. Spleen blood cell counts of Agg-E mice are increased. Cell counts for red blood cells (RBC), white blood cells (WBC), and platelets, and basophil percent cell count 1 day after the 5- and 10-day Agg-E schedules are shown. Control (diagonal hatching) and Agg-E (shaded) means are plotted $\pm$ SEM (unpaired *t*-test with Welch's correction, **p* < 0.05, ***p* < 0.01, #*p* < 0.1; *n* = 5 mice per group).

the tail rattling, were normalized to the value of the control group at each time point (Fig. 7A–C and Supplementary Figs. S1 and S2). ANOVA with Bonferroni post-test (BPT) was used to determine the significance of differences between the control and Agg-E groups at times after the last exposure session.

3.6.1. Freezing

Freezing of mice was computed as the fraction of total time spent immobile (<10% shift of body position between sequential frames of the 15 frames sampled/s) within the partition zone (Fig. 7A). At 1 day following the repeated exposures, mice exposed to the aggressor for 5 days froze more frequently than control mice (*p* < 0.05, BPT), and this behavior persisted after 1.5 weeks of home cage rest (*p* < 0.05, BPT). Mice exposed to the aggressor

for 10 days also froze more than control mice 1 day later (*p* < 0.001, BPT) and after 4 weeks of home cage rest (*p* < 0.05, BPT), but recovered to the control level of freezing in the partition zone after 6 weeks of home cage rest. Overall, freezing did not significantly change between 1 day and 1.5 weeks after the 5-day schedule ($F_{(1,54)} = 0.01$, *p* = 0.92; two-way ANOVA), but after the 10-day schedule became significantly less frequent over the course of home cage rest up to 6 weeks ($F_{(2,54)} = 3.62$, *p* = 0.03; two-way ANOVA).

3.6.2. Grooming

Grooming duration was measured by visual inspection of partition test videos (Fig. 7B). Grooming duration of 5-day Agg-E mice

Table 1

Heart histopathology following daily Agg-E for 5 or 10 days.

5 days of Agg-E	1 day post										1.5 weeks post					
Mouse ID #	1	2	3	4	5	6	7	8			1 ^a	2 ^a	3	4	5	
Myocardial degeneration							2					2		2	2	
Myocarditis, lymphohistiocytic	3	3	2	3			2	3								
Vasculitis, lymphohistiocytic	3		2				1									
Vasculitis, fibrinoid			2	2												
10 days of Agg-E	1 day post											4 weeks post				
Mouse ID #	1	2	3	4	5	6	7	8	9	10	11	1	2	3	4 ^a	5 ^a
Myocardial degeneration							2		3	1	2	2				
Myocarditis, lymphohistiocytic		2	2			2	2	1								
Vasculitis, lymphohistiocytic		2	2	2	2			1								
Vasculitis, fibrinoid		2	2													

Note: no control mice exhibited any of these histopathologies. Out of all of the 28 control mice, only one mouse at 1.5 weeks and another at 4 weeks showed some brown-black pigment. 1: minimal; 2: mild; 3: moderate; 4: marked.

^a Brown-black pigment.




Fig. 5. Dendritic spine density of pyramidal neurons in mPFC increases from 1 day to 4 weeks after the 10-day schedule in control but not Agg-E mice. Brains were collected at 1 day and 4 weeks after the 10-day schedule and Golgi stained. The density of dendritic spines of 36–40 neurons on 10 µm tissue slices was counted for each mouse. Mean dendritic spine density $\pm$ SEM is plotted (unpaired *t*-test with Welch's correction, * $p < 0.05$, # $p = 0.085$; $n = 5$ mice per group). Control (diagonal hatching) and Agg-E (shaded).

was not statistically different than control mice after either 1 day or 1.5 weeks; although there is a trend of increased grooming of Agg-E mice. In contrast, grooming duration of 10-day Agg-E mice was significantly greater than control mice after 1 day ($p < 0.0001$, BPT), was marginally greater after 4 weeks ($p = 0.09$, BPT), and was significantly greater after 6 weeks ($p = 0.05$, BPT). Over the 6 weeks of the home cage rest, the grooming duration of 10-day Agg-E mice was significantly greater than controls ($F_{(2,54)} = 4.69$, $p = 0.01$, two-way ANOVA).

3.6.3. Avoidance of aggressor (time spent per visit to the partition zone)

The time mice spent per visit to the partition zone was used as a measure of avoidance of the aggressor (Fig. 7C). Although Agg-E and control mice made a similar number of total visits to the partition zone, both 5- and 10-day Agg-E mice time spent significantly less time per visit to the partition zone than control mice after 1 day ($p < 0.01$ and $p < 0.001$, respectively, BPT). Five-day exposed mice did not show a significant change in visit time over the 1.5 week home cage rest ($F_{(1,53)} = 2.73$, $p = 0.1$; two-way ANOVA). But 10-day

Agg-E mice increased the time per visit to the partition zone over the 6-week course of home cage rest ($F_{(2,54)} = 5.73$, $p = 0.005$; two-way ANOVA).

3.6.4. Tail rattling

Tail rattling by Agg-E and control mice was monitored by visual inspection of the partition test videos. Mice exposed to the 10-day schedule of aggressor exposures did not show any tail rattling 1 day after the last session, with the exception of one mouse (denoted by a dot in the figure). At the 4- and 6-week recovery points, none of these 10-day Agg-E mice showed any tail rattling (Fig. 7D, star represents zero mice). Similarly, none of the mice exposed to the 5-day schedule of aggressor exposures showed any tail rattling 1 day later. After 1.5 weeks, only one Agg-E mouse exhibited tail rattling (denoted by a dot in figure). In contrast, 45% of control mice demonstrated tail rattling 1 day after the sham, 5-day Agg-E schedule, which increased to 83% of mice after 1.5 weeks of home cage rest. After the sham, 10-day Agg-E schedule, 33% of control mice demonstrated tail rattling at day 1, and after 4 and 6 weeks of home cage rest, the number of control mice exhibiting tail rattling increased to 67% and 100%, respectively.

4. Discussion

4.1. Ethological relevance of the model

We found that mice repeatedly confined cage-within-cage in an aggressor home cage and at random times repeatedly subjected to attack by the aggressor, exhibited acute stress-induced alterations of heart tissue and brain structure as well as alterations in behaviors associated with fear, hypo- and hyper-responding and anxiety, some of which persisted. We chose the aggressor-exposure procedure, adapted from social stress paradigms, because of its ethological relevance to the traumatic stress encountered by warfighters, associated with high incidence PTSD and comorbidities. Social stress models vary greatly in both the procedures for inducing stress and the consequent stress effects [23]. Social stress procedures have been evaluated as models of anxiety disorders, depression and PTSD. Combat veterans with PTSD often meet criteria for several comorbid mood and anxiety disorders, which may be a consequence of the combat arena where multiple exposures can potentially produce synergistic or sensitizing effects




Fig. 6. Partition test cage configuration. The aggressor home cage is divided in half by a fenestrated partition that permits transmission of sensory cues but prevents direct physical interaction between the SJL trained aggressor (Agg) and the subject C57BL/6J mouse. Freezing, aggressor avoidance, tail rattling, and locomotion were measured in the partition zone (hatched area). Grooming and total distance traveled during the 5-min partition test were measured throughout the entire subject side of the cage.




Fig. 7. Behavior evaluations using the partition test: Freezing, grooming, partition avoidance, and tail rattling. Mice were evaluated for behaviors in a 5-min partition test 1 day and 10 days after the 5-day schedule; and 1 day, 4 weeks and 6 weeks after the 10-day schedule. Control (diagonal hatching) and Agg-E (shaded) data were normalized to the control for each time point. Means $\pm$ SEM are plotted (Bonferroni post test of paired comparisons of two-way ANOVA data; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; $n = 6$ –17 mice per group). (A) *Freezing*: frequency of freezing in the partition zone. (B) *Grooming*: duration of self-grooming. (C) *Partition avoidance*: time spent per visit to the partition zone. (D) *Tail rattling*: percentage of the population demonstrating tail rattling. Stars represent zero mice and dots represent one mouse.

[24]. Highly significant relationships have been found between the cumulative number and event severity of post-disaster negative life events and the incidence rate and severity of avoidance-depression dimension of PTSD symptomatology [25]. Also, greater combat exposure has been associated with higher risk for PTSD [1–4].

4.2. Acute stress effects during or 1 day after repeated Agg-E

During the 5- and 10-day schedules, Agg-E mice gained more weight, increased body temperature, and produced fewer territorial urine markings. One day after the 5-day schedule, splenic blood cell counts of Agg-E mice were increased (RBC, WBC, platelets, and basophils). Cardiac histopathologies were also present 1 day after the 5- and 10-day schedules. Specifically, lymphocytic myocarditis, lymphocytic and fibrinoid vasculitis, as well as myocardial degeneration were found in Agg-E mice; none of these heart histopathologies were observed in any control mice. Altered behaviors of Agg-E mice 1 day after the last Agg-E session included: prolonged grooming throughout the entire subject side of the partitioned cage, reduced time spent per visit to the partition zone; and, reduced locomotion and tail rattling and increased freezing within the partition zone.

The *weight gain* of Agg-E mice that we observed persisted over the entire 10-day Agg-E schedule. In contrast, weight loss has been observed in some social defeat models, including those evaluated for aspects of ethologically relevant stresses associated with existing PTSD models [26–29]. However, other studies have shown weight gain. For example, following chronic social defeat stress for 10 days (5 min of direct aggressor exposure followed by housing across a plastic separator with holes for 24 h), C57BL/6J maintained

on standard chow gained weight [30]. In an adult male mouse resident-subordinate social stress model, body weight increased without an increase in food intake [31]. In contrast, the increased body mass and adiposity of Syrian hamsters subjected to Agg-E was accompanied by an increase in food intake [32]. C57BL/6 and BALB/c mice repeatedly exposed (10 days) to an aggressor for 3 min resulted in weight gain, whereas the same number of repeats of aggressor exposure for 10 min did not produce significant changes in weight in either strain [33]. Thus, conflicting effects of social defeat stress on body weight appear to be largely explained by the procedure used to induce stress and genetic strain differences. In humans, chronic stress induces either increased comfort food intake and body weight gain or decreased intake and body weight loss [34].

The *increased body temperature* that we observed has been previously observed in social defeat models [27,35]. Repeated social defeat of male NMRI mice increased their core body temperature and corticosterone, indicative of a chronic stress state [36]. In our study, body temperature was elevated across the 10-day Agg-E schedule, consistent with previous reports of social stress resulting in no adaptation to the tachycardic or hyperthermic responses [36].

We observed *reduced territorial urine markings* during the cage-within-cage sessions over the course of the 10-day schedule, while the control animals showed a high level of urine marking (a hyper-exploratory behavior) at the earlier part of the schedule that subsided, probably due to habituation. In a previous study mice that received 30 bites from trained aggressors during three 2-min encounters showed a deficit in territorial urine marking [37]. A similar deficit in territorial scent marking was also observed in socially stressed Mongolian gerbils [38,39].

The increased blood cell counts of Agg-E mice were determined from spleen sampled 1 day after the 5-day schedule and immediately after the stress of the partition test. The WBC elevation in Agg-E mice is a likely consequence of their fear associated with the conspecific aggressor present in the partition test, which increased plasma catecholamines that are known to transiently increase leukocyte redistribution and leukocyte blood levels within minutes [40]. The RBC elevation is also likely associated with the stress of the partition test as elevated RBC counts are associated with acute stress [41,42]. Only a non-significant trend toward heightened blood cell counts of Agg-E mice is evident the day after the 10-day schedule. Possible explanations for the greater difference in blood cell counts of Agg-E versus control mice after 5 days of Agg-E than after 10 days of Agg-E are that, over the longer stress period, control mice increased their allostatic load as a consequence of the greater number of repeated *sham* Agg-E exposures (e.g., 6 h daily fasting [43,44]), which may have heightened their stress response to the partition test, resulting in elevated RBC, WBC and platelets. Another possible explanation is that in Agg-E mice, the repeated catecholamine-induced redistribution of leukocytes became down-regulated (WBC and basophils) upon repeated stimulation.

Repeated stress and repeated stimulation of blood cells may lead to chronic blood cell dysfunction. Chronic stress has been associated with alterations in blood cell activities and numbers, such as increased platelet reactivity in war veterans with PTSD [45], and elevated leukocyte and total T-cell counts in patients with chronic, primarily combat-related PTSD 20 years after exposure to severe stress of combat [46]. In a repeated social stress model (C57BL/6J mice), exposure to the aggressor for 2, 4 or 6 consecutive days gradually increased the percentages and numbers of neutrophils and monocytes in the blood and the spleen [47]. Persistent stimulation of immune cells by repeated stress may lead to the development of glucocorticoid resistance of leukocytes [47] and other dysregulated immune parameters that have been associated with PTSD [48].

The lymphocytic myocarditis, lymphocytic and fibrinoid vasculitis, and myocardial degeneration in Agg-E mice at 1 day after the 5- and 10-day schedule were not observed in any of the control mice. Other models of stress have shown cardiac disturbances, but none to our knowledge have reported histopathologies such as we have observed. Chronic overcrowding stress of borderline hypertensive Wistar rats was associated with marked subcellular injury of endothelial cells in aorta with mitochondrial damage, presence of vacuoles, increased number of lysosomes, Weibel–Palade bodies, changes of intercellular connections, and local disruption of endothelium; only slight changes were seen in Wistar rats [49]. Repeated social stress of seven consecutive days in a rat resident-intruder model resulted in adrenal hypertrophy and heart rate variability, both of which were inhibited by an antagonist of the corticotropin-releasing factor-1 receptor [50]. Repeated exposure (10 consecutive times, on alternate days) to defeat by a conspecific in a rat model resulted in transient disturbance of heart rate circadian rhythmicity, moderate right ventricle hypertrophy, and the rats failed to develop habituation of cardiac autonomic responsiveness (tachycardia and vagal withdrawal) upon re-exposure to a homotypic acute stressor [51].

The cardiac histopathologies we have observed may be related to Takotsubo cardiomyopathy, also known as “broken heart syndrome”, or stress-induced cardiomyopathy, a syndrome characterized by cardiac-type chest pain and ECG changes, mimicking anterior myocardial infarction and apical aneurysmal dilatation of the left ventricle, which is triggered by a recent severe emotional or physical stressor, and thought to be likely due to catecholamine-mediated myocyte damage and microvascular dysfunction [52].

4.3. Stress effects persisting to 1.5–6 weeks

Some acute stress effects of the aggressor exposure schedules lessened or disappeared with single-housed home cage rest. But other effects became evident or persisted.

4.3.1. Persistent cardiac and neurological histopathologies

The cardiac inflammatory histopathologies disappeared at 1.5 and 4 weeks after the 5- and 10-day Agg-E schedules, respectively. But after these recovery periods, myocardial degeneration with fibroplasia or fibrosis was present in several mice, suggesting that the inflammatory histopathologies resolved into fibroplasia or fibrosis. Interestingly, Takotsubo cardiomyopathy is usually followed by complete recovery; however, in a follow-up of 4.4 ± 4.6 years of 100 patients, 31 patients had continued episodes of chest pain, and 10 patients had recurrent disease, though no difference in survival was found over this follow-up period of only 4 years [53]. Increasing evidence suggests that PTSD is associated with cardiovascular disease, including increased heart rate, elevated blood pressure, decreased heart rate variability, dyslipidemia, and low-grade systemic inflammation and hypercoagulability [54,55]. A review of 4328 male U.S. military veterans has reported that PTSD diagnosis is associated with increased risk of death from early-age heart disease [56].

Dendritic spine density in the mPFC was not different between controls and Agg-E mice at 1 day after the 10-day schedule, but 4 weeks later, dendritic spine density had significantly increased in the control but not in the Agg-E mice. In mouse models, socially stressed animals have exhibited decreased dendritic spine density in both hippocampus and mPFC, as well as an increased dendritic spine density in the basolateral amygdala [57]. Glucocorticoid stress hormones are known to target mPFC and either chronic stress or chronic glucocorticoid administration produces dendritic remodeling in prefrontal pyramidal neurons. Stress also causes increased release of the excitatory amino acid glutamate, which binds NMDA receptors in mPFC [58].

Alterations in mPFC and adjacent and connected neural structures have been revealed in a number of rat stress models [59]. Repeated and chronic restraint stress resulted in a reduced dendritic spine density of pyramidal cells of mPFC [60–62]. The combination of prenatal restraint stress and postnatal chronic mild stress reduced mPFC dendritic spine density [63]. Chronic stress of isolation rearing reduced the volume of the mPFC [64]. The mPFC infralimbic region of the rat in response to chronic restraint stress showed dendritic retraction and spine loss that co-occurred with receptor-mediated impairments to catecholaminergic facilitation of synaptic plasticity; post-stress recovery did not reverse distal dendritic retraction, but showed over extension of proximal dendritic arbors [65]. A competitive NMDA receptor antagonist administered during chronic restraint stress resulted in hypertrophy of apical dendrites, suggesting that NMDA receptor activation is crucial for stress-induced dendritic atrophy in mPFC [58].

PTSD in combat nurses and veterans, firefighters, and those with cumulative adverse life events has been associated with decreased volume of the mPFC [66–68]. PTSD has also been associated with mPFC hypo-responsiveness during symptomatic states and the performance of emotional cognitive tasks, which associated inversely with PTSD symptom severity [69].

4.3.2. Persistent altered behaviors

Behaviors altered in Agg-E mice also diminished or disappeared with home cage rest. But others persisted. Visits of Agg-E mice to the partition zone (partition avoidance) were shorter 1 day after the last 6-h session, and this effect disappeared with as little as 1.5 weeks of home cage rest. In contrast, grooming duration persisted after the 10-day Agg-E schedule, remaining elevated out to 6 weeks of home cage rest. Tail rattling in the partition zone, an

Table 2

Summary of acute and persistent stress effects of repeated aggressor exposure.

Parameter	Effect	Days of Agg-E	Acute	Persistence
Body weight	↑	10	During	nd
Body temperature	↑	10	During	nd
Urine marking	↓	10	During	nd
Blood cell counts	↑	5 and 10	1 day	nd
Heart inflammation or degeneration	↑↑	5 and 10	1 day	–
Heart degeneration	↑	5 and 10	1 day	1.5 and 4 weeks
mPFC dendritic spine density	↓ ^a	10	–	1 day to 4 weeks
Avoidance (time per visit to partition)	↑	5 and 10	1 day	–
Freezing (in partition zone)	↑	5 and 10	1 day	1.5 and out to 6 weeks
Grooming (in entire subject half cage)	↑	5 and 10	1 day	Out to 6 weeks
Tail rattling (in partition zone)	↓	5 and 10	1 day	1.5 and 6 weeks
Locomotion (in partition zone)	↓	5 and 10	1 day	Out to 6 weeks
Locomotion (in entire subject half cage)	↔	5 and 10	–	–

nd: not determined.

^a Increased in control between 1 day and 4 weeks but not in Agg-E; lower dendritic spine density of control suggested at 4 weeks ($p = 0.085$).

agonistic response to the aggressor, was suppressed in Agg-E mice throughout the post-Agg-E period; 100% of control mice displayed tail rattling while none of the 10-day Agg-E mice displayed this behavior after 6 weeks, and only two Agg-E mice ever displayed any tail rattling. Freezing of Agg-E mice in the partition zone was increased the day after the last Agg-E session and persisted to 4 weeks. A summary of all the persistent effects of the Agg-E stress is presented in Table 2.

Grooming duration remained elevated in Agg-E mice even after 6 weeks of home cage rest. Grooming in rodents has been used as a model of stress, anxiety and depression [70]. Repeated restraint stress increased grooming duration in rats, and the antidepressant desipramine decreased this effect, but in the chronic, mild, stress group, desipramine increased grooming frequency [71]. Grooming has been suggested to be relevant to overall increases in stereotypic behavior in depressed patients [72,73]. We measured persistent grooming of rostral regions in our study since anxious rodents generally groom their rostral regions more frequently than caudal regions [74]. Grooming is considered a stress- and arousal-related behavior [75].

Tail rattling of the Agg-E mice was not observed even after 6 weeks of recovery when most of the other behaviors returned to control levels. The tail rattling exhibited by control mice increased over the course of recovery, and was probably initially suppressed due to the stress of the novel environment [76], with food and water deprivation [77], social isolation, and restraint in the cage-within-cage setting (absent an aggressor). Tail rattling is generally agreed to be an agonistic interaction, though it has been variably interpreted as being an aggressive or a defensive behavior, and some consider it a non-specific index of stress and reaction to danger [78–88]. Our observations of a near total absence of tail rattling are in accordance with previous findings that Agg-E generally reduces aggression and increases defensiveness [23], and may be related to the avoidance-depression dimension of PTSD symptomatology.

Freezing in the partition zone in our study exhibited a graded response to the duration of the Agg-E schedule: 10-day Agg-E mice showed significantly more freezing than 5-day Agg-E mice on the day after Agg-E (unpaired t -test with Welch's correction $p = 0.005$). Freezing has been used as a measure of associative fear intensity, and higher freezing responses have been observed with increasing foot shock intensities [10,89,90]. Furthermore, freezing has been considered one component of a classic anxiogenic profile that resulted from 10 days of social defeat in a rat model, and the freezing of socially defeated mice in the presence of a male conspecific was attenuated by treatment with an antagonist of the arginine vasopressin receptor V1bR [91]. Consistent with our result, 10 and 20 days exposure of male C57BL/6J mice to aggression in

daily interactions led to the development of anxiety, assessed in the partition and plus-maze tests [92,93]. Mice defeated in these studies approached the partition less and spent less time per visit to the partition zone the day after the last stress session. This effect was more pronounced in subjects defeated in 20-day as compared with 10-day exposures to aggressive confrontations. In the elevated plus-maze, defeated mice showed fewer open and total entries than controls, and rarely passed from one enclosed arm to another. Thus, in this report, the plus-maze test result is consistent with the partition test freezing data [92]. We observed increased freezing frequency in the partition zone at 1.5 weeks after the 5-day Agg-E and 4 weeks after the 10-day Agg-E and over the 6 weeks (two-way ANOVA). Freezing is suggested to be a contextual conditioned fear response to the aggressor.

Locomotion of Agg-E mice was only altered in the partition zone and it persisted over the recovery periods. Repeated Agg-E has been previously associated with dramatically reduced exploration in the partition test [37]. We found that control and Agg-E mice traveled similar distances within the entire half cage in the partition test (Supplementary Fig. S1) and made similar numbers of visits to the partition zone (not shown). The only difference was that Agg-E mice traveled less within the partition zone (Supplementary Fig. S2), which may indicate their fear of the aggressor.

4.4. Relevance of Agg-E model to PTSD

Animal models relevant to PTSD have been discussed by Yehuda and Antelman [94], Rasmusson and Charney [95], Stam et al. [96], and Siegmund and Wotjak [6]. In our studies, the persistence of some acutely altered behaviors after home cage rest for up to 4–6 weeks suggests symptoms similar to those reported in PTSD and co-morbid depression and anxiety disorders. Our model includes uncontrollable and unpredictable stressors (randomly timed attacks by aggressors) which have been associated with models wherein animals are much more likely to develop behavioral and biochemical manifestations similar to core PTSD symptoms [95,97].

The Agg-E model that we used meets several criteria proposed for the establishment of an animal model of PTSD [6,94]. We assessed behavioral correlates of associative trauma-related memories (i.e., the aggressor), and found increased frequency of freezing, decreased locomotion, and decreased tail rattling in response to the conspecific aggressor (i.e., within the partition zone). Non-associative stimuli are the subject of future studies; however, other published data show that somewhat similar stressors sensitize animals to display anxiety-like behavior in response to novel stressors [26]. Our longitudinal study design revealed

desensitization with time to both freezing in the partition zone and partition avoidance (fear responses). The aversive stimulus (stressor), repeated Agg-E, induced the persistent phenotype we observed, and the intensity of the stressor (5 or 10 days of Agg-E) generally correlated with the severity of symptoms. Although we have not addressed more acute Agg-E, a single social defeat produces robust activation of the HPA-axis [98] and can have long-term consequences ranging to weeks [27]. The symptoms we have observed persisted for several weeks: cardiac degeneration, failure to show the normal increase mPFC dendritic spine density, more grooming, and less tail rattling, less locomotion and more freezing in the partition zone. These symptoms are interpreted to include exaggerated fear responses to the trauma-related cue of the conspecific aggressor across the partition. Although we have not observed hypervigilance, the Agg-E mice showed similar overall activity as controls, as the total locomotion of the two groups outside of the partition zone was similar. Hyperarousal has been suggested in rats exposed to repeated social defeat, as acoustic startle was enhanced and prolonged for up to 10 days [26]. In our study, grooming is considered an arousal-related behavior [75]. The PTSD-like symptoms we observed also included signs that may be interpreted as hypo-responding (emotional blunting, social withdrawal)—i.e., almost nonexistent tail rattling in the partition zone. We also observed considerable inter-individual variability of PTSD-like symptoms, which may possibly result from predispositions to vulnerability and resilience established prior to the experiment, such as during rearing. Predictive validity, i.e., that therapeutics effective in the treatment of PTSD should ameliorate at least part of the PTSD-like symptoms, was also not addressed in the current study. However, it has been noted that social defeat models respond to chronic, but not acute, administration of antidepressants, as is the case in humans; such *pharmacological* validity has not been observed in other stress models, which responded equally to both acute and chronic antidepressant treatments [99–101].

Our data support some of the validity criteria for an animal model of PTSD and general dimensions of traumatic psychosocial stress with regard to the “ethological validity” of the unpredictable and uncontrollable nature of the trauma, “face validity” of symptoms representing PTSD symptoms (associative memory fear, anxiety and hypo- and hyper-responding) and “construct validity” of representing cellular and molecular processes (cardiac pathologies and lack of expected increased dendritic spine density of pyramidal neurons of mPFC).

In other models of repetitive stress or prolonged glucocorticoid administration, neurologic structural changes have been observed, including reduced neurogenesis in the dentate gyrus [102], decreased hippocampus granule cells, and changed electrophysiological properties of hippocampal neurons (see [90] and references therein) as well as the changes to dendritic spine density in mPFC that we and others have observed [103]. Such structural changes are suggested to involve excessive excitatory transmission [90,104]. Recently, Wohleb et al. have shown that the acute stress of repeated social defeat enhanced the inflammatory profile of CD11b⁺ microglia and macrophages in the brain, induced c-Fos activation (in brain regions associated with threat and fear appraisal), anxiety-like behavior, as well as neuroinflammation, and these effects were blocked by β -adrenergic receptor antagonism and required a functional IL-1 receptor type-1 [105].

4.5. Summary

The repeated Agg-E procedure we used induced marked physiological changes consistent with repeated stimulation of the hypothalamic–pituitary axis and sympathetic nervous system that acutely altered the metabolism of heart and brain tissue that later

manifest as cardiac fibroplasia or fibrosis, reduced dendritic spine density of pyramidal neurons in the mPFC, and persistent behaviors indicative of fear, hypo-responding, hyper-responding, and anxiety, reminiscent of features of PTSD and comorbid dimensions.

Disclaimers

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as official Department of the Army position, policy, or decision, unless so designated by other official documentation.

Research was conducted in compliance with the Animal Welfare Act, and other Federal statutes and regulations relating to animals and experiments involving animals and adheres to principles stated in the Guide for the Care and Use of Laboratory Animals (NRC 2011) in facilities that are fully accredited by the Association for the Assessment and Accreditation of Laboratory Animal Care, International.

Citations of commercial organizations or trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

Acknowledgements

This research was sponsored by the U.S. Army Medical Research and Materiel Command and had the financial support of the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), which provided a sabbatical grant to Dr. T.C.M. De Lima. The authors would like to thank Stacy-Ann Miller, Monique Melige, Bintu Sowe, Seshamalani Srinivasan, and Meskerem Jibitu for their technical support and assistance.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bbr.2012.07.022>.

References

- [1] Seal KH, Metzler TJ, Gima KS, Bertenthal D, Maguen S, Marmar CR. Trends and risk factors for mental health diagnoses among Iraq and Afghanistan veterans using Department of Veterans Affairs health care, 2002–2008. *American Journal of Public Health* 2009;99:1651–8.
- [2] Adler AB, Huffman AH, Bliese PD, Castro CA. The impact of deployment length and experience on the well-being of male and female soldiers. *Journal of Occupational Health Psychology* 2005;10:121–37.
- [3] Milliken CS, Auchterlonie JL, Hoge CW. Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq war. *JAMA: The Journal of the American Medical Association* 2007;298:2141–8.
- [4] Kulka RA, Schlenger WE, Fairbank JA, Hough RL, Jordan BK, Marmar CR, et al. Trauma and the Vietnam War generation: report of findings from the National Vietnam veterans readjustment study. New York: Brunner/Mazel; 1990.
- [5] Ramchand R, Schell TL, Karney BR, Osilla KC, Burns RM, Caldarone LB. Disparate prevalence estimates of PTSD among service members who served in Iraq and Afghanistan: possible explanations. *Journal of Traumatic Stress* 2010;23:59–68.
- [6] Siegmund A, Wotjak CT. Toward an animal model of posttraumatic stress disorder. *Annals of the New York Academy of Sciences* 2006;1071:324–34.
- [7] Ohman A, Mineka S. Fears, phobias, and preparedness: toward an evolved module of fear and fear learning. *Psychological Review* 2001;108:483–522.
- [8] American Psychiatric Association. Diagnostic criteria from DSM-IV. Washington, D.C.: The Association; 1994.
- [9] Hunter RG, McCarthy KJ, Milne TA, Pfaff DW, McEwen BS. Regulation of hippocampal H3 histone methylation by acute chronic stress. *Proceedings of the National Academy of Sciences of the United States of America* 2009;106:20912–7.
- [10] Golub Y, Mauch CP, Dahlhoff M, Wotjak CT. Consequences of extinction training on associative and non-associative fear in a mouse model of Posttraumatic Stress Disorder (PTSD). *Behavioural Brain Research* 2009;205:544–9.

- [11] Siegmund A, Dahlhoff M, Habersetzer U, Mederer A, Wolf E, Holsboer F, et al. Maternal inexperience as a risk factor of innate fear and PTSD-like symptoms in mice. *Journal of Psychiatric Research* 2009;43:1156–65.
- [12] Brinks V, de Kloet ER, Oitzl MS. Corticosterone facilitates extinction of fear memory in BALB/c mice but strengthens cue related fear in C57BL/6 mice. *Experimental Neurology* 2009;216:375–82.
- [13] Pizarro JM, Lumley LA, Medina W, Robison CL, Chang WE, Alagappan A, et al. Acute social defeat reduces neurotrophin expression in brain cortical and subcortical areas in mice. *Brain Research* 2004;1025:10–20.
- [14] Robison CL, Meyerhoff JL, Saviolakis GA, Chen WK, Rice KC, Lumley LA. A CRH1 antagonist into the amygdala of mice prevents defeat-induced defensive behavior. *Annals of the New York Academy of Sciences* 2004;1032:324–7.
- [15] Baisley SK, Cloninger CL, Bakshi VP. Fos expression following regimens of predator stress versus footshock that differentially affect prepulse inhibition in rats. *Physiology and Behavior* 2011;104:796–803.
- [16] Power AE, McGaugh JL. Cholinergic activation of the basolateral amygdala regulates unlearned freezing behavior in rats. *Behavioural Brain Research* 2002;134:307–15.
- [17] Wallace KJ, Rosen JB. Neurotoxic lesions of the lateral nucleus of the amygdala decrease conditioned fear but not unconditioned fear of a predator odor: comparison with electrolytic lesions. *Journal of Neuroscience* 2001;21:3619–27.
- [18] Seib LM, Wellman CL. Daily injections alter spine density in rat medial prefrontal cortex. *Neuroscience Letters* 2003;337:29–32.
- [19] Kolb B, Gorny G, Soderpalm AH, Robinson TE. Environmental complexity has different effects on the structure of neurons in the prefrontal cortex versus the parietal cortex or nucleus accumbens. *Synapse* 2003;48:149–53.
- [20] Van De Werd HJ, Rajkowska G, Evers P, Uylings HB. Cytoarchitectonic and chemoarchitectonic characterization of the prefrontal cortical areas in the mouse. *Brain Structure & Function* 2010;214:339–53.
- [21] Kudryavtseva NN. A sensory contact model for the study of aggressive and submissive behavior in male mice. *Aggressive Behavior* 1991;17:285–91.
- [22] Kudryavtseva NN, Bakshantovskaya IV, Koryakina LA. Social model of depression in mice of C57BL/6J strain. *Pharmacology Biochemistry and Behavior* 1991;38:315–20.
- [23] Blanchard RJ, McKittrick CR, Blanchard DC. Animal models of social stress: effects on behavior and brain neurochemical systems. *Physiology and Behavior* 2001;73:261–71.
- [24] Yehuda R, Bierer LM, Pratchett LC, Pelcovitz M. Using biological markers to inform a clinically meaningful treatment response. *Annals of the New York Academy of Sciences* 2010;1208:158–63.
- [25] Maes M, Mylle J, Delmeire L, Janca A. Pre- and post-disaster negative life events in relation to the incidence and severity of post-traumatic stress disorder. *Psychiatry Research* 2001;105:1–12.
- [26] Pulliam JV, Dawagbreh AM, Alema-Mensa E, Plotsky PM. Social defeat stress produces prolonged alterations in acoustic startle and body weight gain in male Long Evans rats. *Journal of Psychiatric Research* 2010;44:106–11.
- [27] Koolhaas JM, Meerlo P, De Boer SF, Strubbe JH, Bohus B. The temporal dynamics of the stress response. *Neuroscience and Biobehavioral Reviews* 1997;21:775–82.
- [28] Bhatnagar S, Vining C, Iyer V, Kinni V. Changes in hypothalamic–pituitary–adrenal function, body temperature, body weight and food intake with repeated social stress exposure in rats. *Journal of Neuroendocrinology* 2006;18:13–24.
- [29] Blanchard DC, Spencer RL, Weiss SM, Blanchard RJ, McEwen B, Sakai RR. Visible burrow system as a model of chronic social stress: behavioral and neuroendocrine correlates. *Psychoneuroendocrinology* 1995;20:117–34.
- [30] Chuang JC, Cui H, Mason BL, Mahgoub M, Bookout AL, Yu HG, et al. Chronic social defeat stress disrupts regulation of lipid synthesis. *Journal of Lipid Research* 2010;51:1344–53.
- [31] Bartolomucci A, Pederzani T, Sacerdote P, Panerai AE, Parmigiani S, Palanza P. Behavioral and physiological characterization of male mice under chronic psychosocial stress. *Psychoneuroendocrinology* 2004;29:899–910.
- [32] Foster MT, Solomon MB, Huhman KL, Bartness TJ. Social defeat increases food intake, body mass, and adiposity in Syrian hamsters. *American Journal of Physiology – Regulatory, Integrative and Comparative Physiology* 2006;290:R1284–93.
- [33] Savignac HM, Finger BC, Pizzo RC, O’Leary OF, Dinan TG, Cryan JF. Increased sensitivity to the effects of chronic social defeat stress in an innately anxious mouse strain. *Neuroscience* 2011;192:524–36.
- [34] Dallman MF, Pecoraro N, Akana SF, La Fleur SE, Gomez F, Houshyar H, et al. Chronic stress obesity: a new view of “comfort food”. *Proceedings of the National Academy of Sciences of the United States of America* 2003;100:11696–701.
- [35] Meerlo P, Overkamp GJ, Koolhaas JM. Behavioural and physiological consequences of a single social defeat in Roman high- and low-avoidance rats. *Psychoneuroendocrinology* 1997;22:155–68.
- [36] Keeney AJ, Hogg S, Marsden CA. Alterations in core body temperature, locomotor activity, and corticosterone following acute and repeated social defeat of male NMRI mice. *Physiology and Behavior* 2001;74:177–84.
- [37] Lumley LA, Sipos ML, Charles RC, Charles RF, Meyerhoff JL. Social stress effects on territorial marking and ultrasonic vocalizations in mice. *Physiology and Behavior* 1999;67:769–75.
- [38] Shimozuru M, Kikusui T, Takeuchi Y, Mori Y. Scent-marking and sexual activity may reflect social hierarchy among group-living male Mongolian gerbils (*Meriones unguiculatus*). *Physiology and Behavior* 2006;89:644–9.
- [39] Shimozuru M, Kikusui T, Takeuchi Y, Mori Y. Social-defeat stress suppresses scent-marking and social-approach behaviors in male Mongolian gerbils (*Meriones unguiculatus*). *Physiology and Behavior* 2006;88:620–7.
- [40] Dhabhar FS, Malarkey WB, Neri E, McEwen BS. Stress-induced redistribution of immune cells – from barracks to battlefields: a tale of three hormones – Curt Richter Award Winner Original Research Article. *Psychoneuroendocrinology*, 2012 Jun 22. PMID: 22727761.
- [41] Brun JF, Khaled S, Raynaud E, Bouix D, Micallef JP, Orsetti A. The triphasic effects of exercise on blood rheology: which relevance to physiology and pathophysiology? *Clinical Hemorheology and Microcirculation* 1998;19:89–104.
- [42] Yalcin O, Bor-Kucukatatay M, Senturk UK, Baskurt OK. Effects of swimming exercise on red blood cell rheology in trained and untrained rats. *Journal of Applied Physiology* 2000;88:2074–80.
- [43] Nowland MH, Hugunin KM, Rogers KL. Effects of short-term fasting in male Sprague-Dawley rats. *Comparative Medicine* 2011;61:138–44.
- [44] Tamakoshi K, Yatsuya H, Kondo T, Hori Y, Zhang H, Ishikawa M, et al. U-shaped association between white blood cell count and fasting plasma glucose level. *Diabetes Care* 2003;26:950.
- [45] Vidovic A, Grubisic-Ilic M, Kozaric-Kovacic D, Gotovac K, Rakos I, Markotic A, et al. Exaggerated platelet reactivity to physiological agonists in war veterans with posttraumatic stress disorder. *Psychoneuroendocrinology* 2011;36:161–72.
- [46] Boscarino JA, Chang J. Higher abnormal leukocyte and lymphocyte counts 20 years after exposure to severe stress: research and clinical implications. *Psychosomatic Medicine* 1999;61:378–86.
- [47] Engler H, Bailey MT, Engler A, Sheridan JF. Effects of repeated social stress on leukocyte distribution in bone marrow, peripheral blood and spleen. *Journal of Neuroimmunology* 2004;148:106–15.
- [48] Pace TW, Heim CM. A short review on the psychoneuroimmunology of post-traumatic stress disorder: from risk factors to medical comorbidities. *Brain, Behavior, and Immunity* 2011;25:6–13.
- [49] Okruhlicova L, Dlugosova K, Mitasikova M, Bernatova I. Ultrastructural characteristics of aortic endothelial cells in borderline hypertensive rats exposed to chronic social stress. *Physiological Research/Academia Scientiarum Bohemoslovaca* 2008;57(Suppl. 2):S31–7.
- [50] Wood SK, McFadden KV, Grigoriadis D, Bhatnagar S, Valentino RJ. Depressive and cardiovascular disease comorbidity in a rat model of social stress: a putative role for corticotropin-releasing factor. *Psychopharmacology* 2012;222(2):325–36.
- [51] Carnevali L, Mastorci F, Graiani G, Razzoli M, Trombini M, Pico-Alfonso MA, et al. Social defeat and isolation induce clear signs of a depression-like state, but modest cardiac alterations in wild-type rats. *Physiology and Behavior* 2012;106:142–50.
- [52] Milinis K, Fisher M. Takotsubo cardiomyopathy: pathophysiology and treatment. *Postgraduate Medical Journal* 2010. PMID: 22647668.
- [53] Elesber AA, Prasad A, Lennon RJ, Wright RS, Lerman A, Rihal CS. Four-year recurrence rate and prognosis of the apical ballooning syndrome. *Journal of the American College of Cardiology* 2007;50:448–52.
- [54] von Kanel R, Abbas CC, Begre S, Gander ML, Saner H, Schmid JP. Association between posttraumatic stress disorder following myocardial infarction and liver enzyme levels: a prospective study. *Digestive Diseases and Sciences* 2010;55:2614–23.
- [55] Von Kanel R, Kraemer B, Saner H, Schmid JP, Abbas CC, Begre S. Posttraumatic stress disorder and dyslipidemia: previous research and novel findings from patients with PTSD caused by myocardial infarction. *The World Journal of Biological Psychiatry* 2010;11:141–7.
- [56] Boscarino JA. A prospective study of PTSD and early-age heart disease mortality among Vietnam veterans: implications for surveillance and prevention. *Psychosomatic Medicine* 2008;70:668–76.
- [57] Meyerhoff JL, Lumley LA, Robison CL, Gorny G, Saviolakis G, Kolb B. Chronic social stress in mice is associated with increased dendritic spine density in the amygdala. *Neuroscience Meeting* 2005, 872.5.
- [58] Martin KP, Wellman CL. NMDA receptor blockade alters stress-induced dendritic remodeling in medial prefrontal cortex. *Cerebral Cortex* 2011;21:2366–73.
- [59] Brown SM, Henning S, Wellman CL. Mild, short-term stress alters dendritic morphology in rat medial prefrontal cortex. *Cerebral Cortex* 2005;15:1714–22.
- [60] Radley JJ, Rocher AB, Rodriguez A, Ehlenberger DB, Dammann M, McEwen BS, et al. Repeated stress alters dendritic spine morphology in the rat medial prefrontal cortex. *The Journal of Comparative Neurology* 2008;507:1141–50.
- [61] Radley JJ, Rocher AB, Miller M, Janssen WG, Liston C, Hof PR, et al. Repeated stress induces dendritic spine loss in the rat medial prefrontal cortex. *Cerebral Cortex* 2006;16:313–20.
- [62] Radley JJ, Sisti HM, Hao J, Rocher AB, McCall T, Hof PR, et al. Chronic behavioral stress induces apical dendritic reorganization in pyramidal neurons of the medial prefrontal cortex. *Neuroscience* 2004;125:1–6.
- [63] Michelsen KA, van den Hove DL, Schmitz C, Segers O, Prickaerts J, Steinbusch HW. Prenatal stress and subsequent exposure to chronic mild stress influence dendritic spine density and morphology in the rat medial prefrontal cortex. *BMC Neuroscience* 2007;8:107.

- [64] Day-Wilson KM, Jones DN, Southam E, Cilia J, Totterdell S. Medial prefrontal cortex volume loss in rats with isolation rearing-induced deficits in prepulse inhibition of acoustic startle. *Neuroscience* 2006;141: 1113–21.
- [65] Goldwater DS, Pavlides C, Hunter RG, Bloss EB, Hof PR, McEwen BS, et al. Structural and functional alterations to rat medial prefrontal cortex following chronic restraint stress and recovery. *Neuroscience* 2009;164: 798–808.
- [66] Rauch SL, Shin LM, Segal E, Pitman RK, Carson MA, McMullin K, et al. Selectively reduced regional cortical volumes in post-traumatic stress disorder. *Neuroreport* 2003;14:913–6.
- [67] Shin LM, Wright CI, Cannistraro PA, Wedig MM, McMullin K, Martis B, et al. A functional magnetic resonance imaging study of amygdala and medial prefrontal cortex responses to overtly presented fearful faces in posttraumatic stress disorder. *Archives of General Psychiatry* 2005;62:273–81.
- [68] Ansell EB, Rando K, Tuit K, Guarnaccia J, Sinha R. Cumulative adversity and smaller gray matter volume in medial prefrontal, anterior cingulate, and insula regions. *Biological Psychiatry* 2012;72:57–64.
- [69] Shin LM, Rauch SL, Pitman RK. Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. *Annals of the New York Academy of Sciences* 2006;1071:67–79.
- [70] Kalueff AV, Tuohimaa P. Experimental modeling of anxiety and depression. *Acta neurobiologiae experimentalis* 2004;64:439–48.
- [71] D'Aquila PS, Peana AT, Carboni V, Serra G. Exploratory behaviour and grooming after repeated restraint and chronic mild stress: effect of desipramine. *European Journal of Pharmacology* 2000;399:43–7.
- [72] van Erp AM, Kruk MR, Meelis W, Willekens-Bramer DC. Effect of environmental stressors on time course, variability and form of self-grooming in the rat: handling, social contact, defeat, novelty, restraint and fur moistening. *Behavioural Brain Research* 1994;65:47–55.
- [73] Coscia EM, Fentress JC. Neurological dysfunction expressed in the grooming behavior of developing weaver mutant mice. *Behavior Genetics* 1993;23:533–41.
- [74] Denmark A, Tien D, Wong K, Chung A, Cachat J, Goodspeed J, et al. The effects of chronic social defeat stress on mouse self-grooming behavior and its patterning. *Behavioural Brain Research* 2010;208:553–9.
- [75] Carter M, de Lecea L. Hyperarousal and post-traumatic stress disorder: a role for the hypocretin system. In: Shiromani P, eDoux JE, Keane TM, editors. *Post-traumatic stress disorder: basic science and clinical practice*. Humana Press, A Part of Springer Science + Business Media, LLC; 2009. p. 201.
- [76] Misslin R, Herzog F, Koch B, Ropartz P. Effects of isolation, handling and novelty on the pituitary–adrenal response in the mouse. *Psychoneuroendocrinology* 1982;7:217–21.
- [77] Oarada M, Nikawa T, Kurita N. Effect of timing of food deprivation on host resistance to fungal infection in mice. *British Journal of Nutrition* 2002;88:151–8.
- [78] Bigi S, Huber C, De Acetis L, Alleva E, Dixon AK. Removal of the submaxillary salivary glands first increases and then abolishes the agonistic response of male mice in repeated social encounters. *Physiology and Behavior* 1994;55:13–9.
- [79] Corridi P, Chiarotti F, Bigi S, Alleva E. Familiarity with conspecific odor and isolation-induced aggressive behavior in male mice (*Mus domesticus*). *Journal of Comparative Psychology* 1993;107:328–35.
- [80] Godar SC, Bortolato M, Frau R, Dousti M, Chen K, Shih JC. Maladaptive defensive behaviours in monoamine oxidase A-deficient mice. *The International Journal of Neuropsychopharmacology* 2011;14:1195–207.
- [81] Krsiak M, Sulcova A, Donat P, Tomasikova Z, Dlohozkova N, Kosar E, et al. Can social and agonistic interactions be used to detect anxiolytic activity of drugs? *Progress in Clinical and Biological Research* 1984;167:93–114.
- [82] Lumley LA, Robison CL, Slusher BS, Wozniak K, Dawood M, Meyerhoff JL. Reduced isolation-induced aggressiveness in mice following NAALADase inhibition. *Psychopharmacology* 2004;171:375–81.
- [83] Meek LR, Myren K, Sturm J, Burau D. Acute paternal alcohol use affects offspring development and adult behavior. *Physiology and Behavior* 2007;91:154–60.
- [84] Palanza P, Parmigiani S, vom Saal FS. Urine marking and maternal aggression of wild female mice in relation to anogenital distance at birth. *Physiology and Behavior* 1995;58:827–35.
- [85] Patel A, Siegel A, Zalcman SS. Lack of aggression and anxiolytic-like behavior in TNF receptor (TNF-R1 and TNF-R2) deficient mice. *Brain, Behavior, and Immunity* 2010;24:1276–80.
- [86] Sukhotina IA, Beshpalov AY. Effects of the NMDA receptor channel blockers memantine and MK-2/579 on morphine withdrawal-facilitated aggression in mice. *Psychopharmacology* 2000;149:345–50.
- [87] Terranova ML, Laviola G, de Acetis L, Alleva E. A description of the ontogeny of mouse agonistic behavior. *Journal of Comparative Psychology* 1998;112:3–12.
- [88] Yoshimura H, Ogawa N. Pharmacological analysis of agonistic behavior between resident and intruder mice: effect of anticholinergic drugs. *Japanese Journal of Pharmacology* 1982;32:1111–6.
- [89] Siegmund A, Wotjak CT. A mouse model of posttraumatic stress disorder that distinguishes between conditioned and sensitized fear. *Journal of Psychiatric Research* 2007;41:848–60.
- [90] Golub Y, Kaltwasser SF, Mauch CP, Herrmann L, Schmidt U, Holsboer F, et al. Reduced hippocampus volume in the mouse model of posttraumatic stress disorder. *Journal of Psychiatric Research* 2011;45:650–9.
- [91] Litvin Y, Murakami G, Pfaff DW. Effects of chronic social defeat on behavioral and neural correlates of sociality: vasopressin, oxytocin and the vasopressin-ergic V1b receptor. *Physiology and Behavior* 2011;103:393–403.
- [92] Avgustinovich DF, Gorbach OV, Kudryavtseva NN. Comparative analysis of anxiety-like behavior in partition and plus-maze tests after agonistic interactions in mice. *Physiology and Behavior* 1997;61:37–43.
- [93] Kudryavtseva NN, Bondar NP, Avgustinovich DF. Effects of repeated experience of aggression on the aggressive motivation and development of anxiety in male mice. *Neuroscience and Behavioral Physiology* 2004;34: 721–30.
- [94] Yehuda R, Antelman SM. Criteria for rationally evaluating animal models of posttraumatic stress disorder. *Biological Psychiatry* 1993;33:479–86.
- [95] Rasmussen AM, Charney DS. Animal models of relevance to PTSD. *Annals of the New York Academy of Sciences* 1997;821:332–51.
- [96] Stam R, Bruijnzeel AW, Wiegant VM. Long-lasting stress sensitization. *European Journal of Pharmacology* 2000;405:217–24.
- [97] Foa EB, Zinbarg R, Rothbaum BO. Uncontrollability and unpredictability in post-traumatic stress disorder: an animal model. *Psychological Bulletin* 1992;112:218–38.
- [98] Huhman KL, Moore TO, Ferris CF, Mougey EH, Meyerhoff JL. Acute and repeated exposure to social conflict in male golden hamsters: increases in plasma POMC-peptides and cortisol and decreases in plasma testosterone. *Hormones and Behavior* 1991;25:206–16.
- [99] Berton O, McClung CA, Dileone RJ, Krishnan V, Renthall W, Russo SJ, et al. Essential role of BDNF in the mesolimbic dopamine pathway in social defeat stress. *Science* 2006;311:864–8.
- [100] Tsankova NM, Berton O, Renthall W, Kumar A, Neve RL, Nestler EJ. Sustained hippocampal chromatin regulation in a mouse model of depression and antidepressant action. *Nature Neuroscience* 2006;9:519–25.
- [101] Golden SA, Covington 3rd HE, Berton O, Russo SJ. A standardized protocol for repeated social defeat stress in mice. *Nature Protocols* 2011;6:1183–91.
- [102] Dranovsky A, Hen R. Hippocampal neurogenesis: regulation by stress and antidepressants. *Biological Psychiatry* 2006;59:1136–43.
- [103] Hammack SE, Cooper MA, Lezak KR. Overlapping neurobiology of learned helplessness and conditioned defeat: implications for PTSD and mood disorders. *Neuropharmacology* 2012;62:565–75.
- [104] Kim JJ, Foy MR, Thompson RF. Behavioral stress modifies hippocampal plasticity through N-methyl-D-aspartate receptor activation. *Proceedings of the National Academy of Sciences of the United States of America* 1996;93:4750–3.
- [105] Wohleb ES, Hanke ML, Corona AW, Powell ND, Stiner LM, Bailey MT, et al. beta-Adrenergic receptor antagonism prevents anxiety-like behavior and microglial reactivity induced by repeated social defeat. *Journal of Neuroscience* 2011;31:6277–88.