

**A New Method for the Production of Tetranitroglycoluril
From Imidazo-[4,5-*d*]-Imidazoles With the Loss of
Dinitrogen Oxide**

by William M. Sherrill and Eric C. Johnson

ARL-TR-6829

February 2014

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Army Research Laboratory

Aberdeen Proving Ground, MD 21005-5066

ARL-TR-6829**February 2014**

A New Method for the Production of Tetranitroglycoluril From Imidazo-[4,5-*d*]-Imidazoles With the Loss of Dinitrogen Oxide

William M. Sherrill

Weapons and Materials Research Directorate, ARL

Eric C. Johnson

Bowhead Science and Technology

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) February 2014		2. REPORT TYPE Final		3. DATES COVERED (From - To) May 2013–September 2013	
4. TITLE AND SUBTITLE A New Method for the Production of Tetranitroglycoluril From Imidazo-[4,5-d]-Imidazoles With the Loss of Dinitrogen Oxide				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) William M. Sherrill and Eric C. Johnson*				5d. PROJECT NUMBER AH-43	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory ATTN: RDRL-WML-C Aberdeen Proving Ground, MD 21005-5066				8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-6829	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES *Bowhead Science and Technology, LLC., 103 Bata Blvd, Belcamp, MD 21017.					
14. ABSTRACT A new method for the preparation of tetranitroglycoluril (TNGU) (also known as Sorguyl), is described in which imidazo-[4,5-d]-imidazoles are nitrated with the elimination of dinitrogen oxide (N ₂ O) to generate TNGU. This method of TNGU synthesis results in material that is less sensitive than material produced via alternative routes.					
15. SUBJECT TERMS TNGU, Sorguyl, DINGU, synthesis, explosive					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 26	19a. NAME OF RESPONSIBLE PERSON William M. Sherrill
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 410-278-8608

Contents

List of Figures	iv
List of Tables	iv
Acknowledgments	v
1. Introduction	1
2. Computational Analysis	2
3. Synthesis	3
4. Mechanism	4
4.1 Scheme 1—Proposed Mechanism for the Intramolecular Decomposition and Extrusion of N ₂ O	6
4.2 Scheme 2—Proposed Mechanism for the Intermolecular Decomposition and Extrusion on N ₂ O	6
5. Sensitivity	8
6. Experimental	9
7. Conclusions	10
8. References	11
Appendix. FTIR Data	13
List of Symbols, Abbreviations, and Acronyms	15
Distribution List	17

List of Figures

Figure 1. Structures of high explosives TNGU, RDX, and HMX.....	1
Figure 2. TNGU showing location of bridgehead and carbonyl positions.	2
Figure 3. Structure of starting material imidazo imidazoles 4-6 and target 7.....	2
Figure 4. SPM of target material 7 compared to RDX and HMX.	3
Figure 5. Postulated intermediate in the formation of 1.	5
Figure 6. Literature examples of substrates believed to extrude N ₂ O.	5
Figure 7. Reaction headspace FTIR monitoring setup.....	7
Figure 8. Concentration of N ₂ O in headspace of reactions over time.	8

List of Tables

Table 1. Performance predictions from Cheetah 7.0.	3
Table 2. Summary of nitration conditions and results.	4
Table 3. Sensitivity data.....	8

Acknowledgments

The authors wish to thank Mr. Roy Maulbetsch, Mr. Terry Piatt, and Ms. Lori Pridgeon for general equipment assistance and supply management. The authors also wish to thank Drs. Eric Bukowski, Kimberly Spangler, Brian Roos, Thuvan Piehler, Joseph Banning, and Mr. Stephen Aubert for procedural input and technical review. The authors also wish to express their gratitude to Drs. Edward Byrd and Betsy Rice for making available their computer codes and related scripts.

INTENTIONALLY LEFT BLANK.

1. Introduction

Tetranitroglycoluril* (TNGU) **1** is a powerful high explosive that was first publicly described in the 1970s by the French (1–3). This material, with a density of 2.01 g/cm³ and heat of formation (ΔH_f) of 50 kJ/mol makes it a powerful high explosive similar to the standard military explosives 1,3,5-trinitroperhydro-1,3,5-triazine (RDX, **2**) and octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (HMX, **3**) (4) (figure 1).




Figure 1. Structures of high explosives TNGU, RDX, and HMX.

Compound **1** has several undesirable characteristics that have impeded its adoption into military formulations. The first of these is its inherent water lability. The dinitrourea moiety present in the molecule readily hydrolyzes even on exposure to atmospheric moisture. This means formulations containing TNGU will rapidly degrade upon exposure to normal atmospheric conditions (5). While this feature is usually detrimental in an explosive, TNGU containing formulations are being investigated for use in potential self-remediating munitions to lessen the hazards associated with unexploded ordinance (UXO) (6).

Another detrimental feature of TNGU is the inherent sensitivity of the material that is synthesized and isolated according to literature procedures (6). TNGU as isolated using these methods has a 50% initiation (H_{50}) value for impact sensitivity of 4.10 inches (in) while RDX and HMX are 9.08 in, respectively. Additionally, TNGU has a friction measurement of 54 Newtons (N) while RDX and HMX both are approximately 120 N. Taken together, both of these numbers place TNGU in a category of materials that are usually deemed too sensitive to handle or formulate.

* Also known as Sorguyl.

Recent attempts to rectify both the hydrolytic instability and the sensitivity of TNGU through substitution of the bridgehead hydrogens resulted only in material that was energetically less powerful and still of comparably sensitivity and stability (7). This led to the idea of using substitution of the oxygen from the carbonyl position in the dinitrourea moiety as an additional means of potentially tuning both the sensitivity and the hydrolytic instability of the molecule (figure 2). To this end, several different imidazo imidazoles **4-6** were prepared using known literature procedures (8) and then attempts were made to further nitrate them to produce a novel material **7** (figure 3). Both **5** and **6** are noted in the Kony paper (8) to be explosives in their own right; however, complete sensitivity analysis was never reported. A separate report details the synthesis of these materials and found that while both **5** and **6** may be considered energetic materials, both are significantly less sensitive than RDX (9).




Figure 2. TNGU showing location of bridgehead and carbonyl positions.




Figure 3. Structure of starting material imidazo imidazoles 4-6 and target 7.

2. Computational Analysis

Having determined the sensitivity of these materials, further nitration was then attempted in an effort to synthesize **7**. The target **7** is predicted by quantum mechanical models developed by Rice (10–14) to have a density of 2.032 g/cm³ and a ΔH_f of +617.01 kJ/mol. From these numbers, the predicted performance of this material exceeds HMX by almost 30% making it a highly desirable target (table 1).

Table 1. Performance predictions from Cheetah 7.0 (4).

Substance	ρ^a	ΔH_f (kJ/mol)	P_{cl}^d (GPa)	D_v^e (km/s)	ΔH_d^f (kJ/mL)	OB ^g (%)
7	2.03 ^b	617.01 ^b	42.51	9.621	12.17	+11.70
TNGU ^c	2.01	50.0	41.77	9.566	11.78	+4.97
RDX ^c	1.816	70.01	33.46	8.862	10.42	-21.61
HMX ^c	1.90	75.02	37.19	9.246	11.00	-21.61

^aDensity. ^bPredicted using the methods of Rice (10–14). ^c ΔH_f and density numbers obtained from Cheetah 7.0 database.

^dChapman-Jouguet pressure. ^eDetonation velocity. ^fHeat of detonation. ^gOxygen balance.

Using visualization software the theoretical electron surface potential map (SPM) of **7** was examined for areas of large predicted electron buildup or deficiency. The areas of high-electron density in figure 4 are represented in blue, while areas of electron deficiency are shown in red. It is currently believed that molecules exhibiting large areas of charge separation are generally less stable than molecules with an SPM showing a well distributed surface charge. Upon examining the SPM of **7**, it can be observed that the center of the molecule exhibits a higher area of electron density as compared to RDX and HMX suggesting that **7** may be more sensitive.




Figure 4. SPM of target material **7** compared to RDX and HMX.

3. Synthesis

With the predicted performance and sensitivity in hand, the synthesis of **7** was attempted from all three starting materials **4**, **5**, and **6**. None of the standard suite of acidic nitration conditions of 100% nitric acid (HNO₃), mixed sulfuric acid (H₂SO₄)/<98% HNO₃, and acetic anhydride (Ac₂O)/100% HNO₃ were found to be strong enough conditions to convert the lesser nitrated materials to the desired target. When subjecting materials **5** and **6** to >98% HNO₃, no reaction is observed; however, when using **4**, the material is recovered as the nitrate salt, **5**. All compounds

4-6 exhibit decomposition in mixed H₂SO₄/HNO₃ conditions and no products from these reactions were successfully recovered. Compounds **4** and **5** are both converted to the dinitro compound **6** when subjected to Ac₂O/HNO₃ conditions, but were unable to be nitrated further even with extended reaction times and heating. It was discovered that **4-6** can be further nitrated by employing a nitration mixture using trifluoroacetic anhydride (TFAA) in conjunction with HNO₃. In this case, when **4**, **5**, or **6** is dissolved in <98% HNO₃ and TFAA is added slowly, a white powder slowly precipitates over time with the evolution of gas bubbles. After several hours, the powder was isolated and analyzed to find the material was not **7** as anticipated, but rather TNGU. This was further confirmed by obtaining a nuclear magnetic resonance (NMR) spectrum showing perfect overlap of signals from a mixture of **1** prepared via the nitration of **4-6** and of TNGU prepared via the literature method (6). All three starting materials **4-6** yield TNGU as the product under the TFAA/HNO₃ conditions with reasonable yields although **5** consistently has product yields between 70% to 80% (table 2).

Table 2. Summary of nitration conditions and results.

Substance	>98% HNO ₃	>98% H ₂ SO ₄ />98% HNO ₃	Ac ₂ O/<98% HNO ₃	TFAA/<98% HNO ₃
4	5	Decomposes	6	1
5	No RXN	Decomposes	6	1
6	No RXN	Decomposes	No RXN	1

4. Mechanism

Of the myriad of possible scenarios, it is postulated that two are the most likely. The first being the α -carbon* of **7** is so electron withdrawn that the molecule is rapidly and completely hydrolyzed to TNGU on exposure to moisture. This theory was quickly dismissed by repeating the experiments using meticulously dried starting materials and glassware in conjunction with completely anhydrous reagents under an inert gas atmosphere. Workup was done by allowing the powder to settle to the bottom of the nitration mixture and decanting the mother liquor. The product was then washed extensively with anhydrous dichloromethane (DCM) and immediately analyzed. Analysis of the material obtained showed it to be identical to material isolated in the original experiments indicating that hydrolysis of **7** is not responsible for the formation of **1** under the reaction conditions or under workup.

The second scenario employs intra or intermolecular degradation with elimination of dinitrogen oxide (N₂O) as the driving force behind the formation of **1**. Once the material is subjected to the

* α designates the carbonyl carbon of the imidazo imidazole.

nitration conditions, it is plausible that each of the materials **4-6**, proceeds through an intermediate resembling **8** (figure 5). The α -carbon of this species is severely electron deficient making it highly susceptible to nucleophilic attack.




Figure 5. Postulated intermediate in the formation of **1**.

This type of reactivity with the elimination of N_2O has been documented previously by Wright and McKay (15) in their attempt to further nitrate **9** while only recovering **11** (figure 6). They proposed an intermediate **10**, which could easily rearrange under the nitration conditions with elimination of N_2O to ultimately form **11**. Additionally, this type of reactivity has been used in the synthesis of 2,6-dioxo-1,3,4,5,7,8-hexanitrodecahydro-1H,5H-diimidazo[4,5-b:4',5'-e]pyrazine (HHTDD) **13** starting from **12** (5).





Figure 6. Literature examples of substrates believed to extrude N_2O .


The intramolecular decomposition mechanism is envisioned to employ the oxygen of the nitrimino moiety as the nucleophile and then proceed through a four-membered intermediate **14**. (Scheme 1). The subsequent rearrangement of the four-membered cycle results in the liberation of N_2O and a nitrourea type species **6**. This mechanism is plausible whether the starting intermediate is a dinitro- or trinitroimidazo imidazole.

4.1 Scheme 1—Proposed Mechanism for the Intramolecular Decomposition and Extrusion of N₂O



Alternatively, the intermolecular mechanism starts from the same intermediate; however, it uses an additional molecule of **8** as the oxygen source (Scheme 2).

4.2 Scheme 2—Proposed Mechanism for the Intermolecular Decomposition and Extrusion on N₂O



There are two possible ways to test the plausibility of the degradation theory. The first is ¹⁸O labeled HNO₃ to pinpoint the source of the oxygen nucleophile. This is prohibitively expensive due to the cost of ¹⁸O labeled HNO₃. The second, and most feasible, is to monitor the headspace of the nitration mixture for the appearance of a higher than baseline concentration of N₂O. N₂O gas exhibits a very specific doublet in the infrared (IR) at approximately 2,235 cm⁻¹. Using headspace Fourier transform infrared (FTIR) spectroscopy it was possible to monitor the concentration of N₂O in the headspace of a nitration mixture without substrate present. This was used as the baseline for the subsequent experiments. The setup for the headspace monitoring experiment is shown in figure 7. For comparison, a spectrum of pure N₂O is available in the appendix of this report.




Figure 7. Reaction headspace FTIR monitoring setup.

The highest concentration of N_2O measured in the headspace of the baseline nitration mixture never exceeded 0.05 absorbance units (AU) indicating the amount of N_2O produced from the nitration mixture itself is minimal under the conditions used in the experiment. When glycoluril is added to the nitration mixture, the measured concentration of N_2O increases slightly; however, the maximum concentration still never exceeded 0.20 AU. When compounds **4**, **5**, or **6**, are subjected to the nitration conditions, the maximum concentration of N_2O observed in the headspace is more than five times higher than the maximum concentration observed for glycoluril (figure 8). The fact that TNGU is the only product recovered from the nitration of **4-6** together with those compounds exhibiting a maximum N_2O headspace concentration of 1.45 AU, a marked increase over reactions containing glycoluril, are strong indicators of a reaction mechanism similar to what is depicted in Scheme 1 and Scheme 2 is in operation.




Figure 8. Concentration of N_2O in headspace of reactions over time.

5. Sensitivity

It is interesting to note that the sensitivity of **1** produced via the nitration of **4-6** with TFAA/ HNO_3 is markedly improved over **1** produced using the literature (6) Ac_2O/HNO_3 nitration of glycoluril. As of yet, the reason for this increased stability is not completely understood although it is postulated that morphological differences between the materials may be responsible. TNGU produced via the (6) method has a drop height of 4.10 in on our impact testing apparatus while that produced via the TFAA method has a drop height of 11.07 in. The friction and electrostatic discharge (ESD) measurement between both samples remain comparable.

Table 3. Sensitivity data.

Substance	Impact (in) ^a	Friction (N) ^b	ESD (J) ^c
TFAA/ HNO_3 1	11.07	70	3.25
Ac_2O/HNO_3 1	4.10	54	3.25
RDX	9.08	120	0.625
HMX	9.60	120	0.025

^a H_{50} impact height determined on an apparatus using a 2-kg weight by the Langlie one-shot method (16). ^bFriction determined on a Julius Peter's BAM friction apparatus. ^cESD determined using an ABL Laboratories ESD apparatus.

6. Experimental

H₅₀ values for drop weight testing were determined using the Langlie one-shot method on a tester dropping a 5-lb weight from a maximum height of 60 in (16). Friction sensitivity measurements were determined on a BAM friction apparatus and ESD was determined using an ABL Laboratories ESD apparatus. NMR experiments were done using an Anasazi Instruments 90-megahertz (MHz) nuclear magnetic resonance (NMR). FTIR was obtained from a Brüker Alpha-T fitted with a 5-cm path length gas flow cell in the transmission module, or a diamond-attenuated total reflectance (ATR) module. Differential scanning calorimetry was conducted on a TA instruments Q10 or Q20 calorimeter. All chemicals were obtained from Sigma-Aldrich, St. Louis, MO, and were used as received.

Note: While these compounds were prepared without incident according the following procedures, these materials are energetic and should be prepared and handled cautiously by trained personnel.

Synthesis of 4-6

The syntheses of these compounds were carried out according to the methods detailed by Kony (8). These procedures were quickly and accurately reproduced to yield **4-6** and analysis of the resulting products matched the data given in his publication. Sensitivity analysis of these materials is available in a separate report (9).

Alternate preparation of 5

Kony originally prepared **5** by the action of 100% HNO₃ on the purified hydrochloride salt **4**. It was discovered that **5** is most easily prepared by taking the crude **4** obtained from the procedures described in the Kony report and dissolving it in 70% HNO₃. Once all the material was dissolved, it could be crash precipitated by pouring over ice followed by purification from dilute HNO₃ as described by Kony.

Synthesis of 1 from 4-6

Representative procedure from 4: 100 mg (0.45 mmol) of **4** was dissolved in 2 milliliters (mL) of 100% HNO₃ at 0 °C under an inert nitrogen (N₂) atmosphere. To this solution was added 2 mL of TFAA drop wise at such a rate as to keep the temperature of the solution below 10 °C. After the addition was completed, the reaction was allowed to warm to room temperature and stirred for 2 h at which point stirring was stopped and the precipitant was allowed to settle. The mother liquor was decanted and the residual solid was washed with anhydrous DCM (5 × 5 mL). The solid **1** was then dried under vacuum (10 torr) at ambient temperature for 2 h. The total amount of **1** recovered from this process was 0.701 g (0.22 mmol, 49%).

From 5: 1 was prepared from **5** according to the preceding representative procedure. The total amount of **1** recovered from 100 mg (0.40 mmol) of **5** was 104 mg (0.32 mmol, 80%).

From 6: 1 was prepared from **6** according to the representative procedure. The total amount of **1** recovered from 100 mg (0.43 mmol) of **6** was 0.892 mg (0.28 mmol, 65%).

Scale up to 2 g of 1 from 5: 2.027 g (8.16 mmol) of nitrate salt **5** was dissolved into 40 mL of 100% HNO₃ at 0 °C under an N₂ atmosphere. To this solution was added drop wise 40 mL of TFAA at a rate to keep the temperature under 10 °C. Upon completion of the addition, the material was allowed to warm to ambient temperature and stirring was continued for 2 h. After the time had elapsed, the suspension was cooled to 0 °C, stirring was stopped and the suspended solid was allowed to settle. The mother liquor was decanted and the remaining solid was washed with DCM (5 × 20 mL). The material was then vacuum dried (10 torr 20 °C) for 2 h. The total amount of **1** recovered from this process was 1.89 g (5.86 mmol, 72%)

Procedure for headspace analysis

The Brüker Alpha-T was fitted with the transmission module and in it was placed a 5-cm-long gas flow cell fitted with zinc selenide (ZnSe) windows. A sample injection port covered with a septum was fitted to the intake side of the gas supply line approximately 6 in prior to the cell. The exhaust of the cell was vented into the fume hood. N₂ carrier gas at 12 mL/minute (min) was flowed through the cell at all times while measuring the concentration of N₂O in the headspace. A background of pure N₂ was used and a sample of pure N₂O was injected into the cell to confirm the setup was functioning correctly. A 6-mL sample of the reaction headspace from a 100-mg scale reaction was injected and analyzed at 3-min intervals. The amount of N₂O present in the sample was determined by measuring the absorbance unit values for the N₂O peak of the doublet at approximately 2,235 cm⁻¹.

7. Conclusions

A new method for the synthesis of **1** has been discovered from materials other than glycoluril. This new method causes the substrates to lose N₂O during the nitration process and the product obtained via this new route is more stable to external insult than material produced according to traditional literature methods while still having analytics matching that of the material produced according to the literature procedures.


8. References

1. Boileau, J.; Emeury, J. M. L.; Kehren, J. P. A. *Tetranitroglycoluril for Explosives*, Ger. Offen. 2,435,651, Société Nationale de Pouders et Explosifs, Paris, France, **1975**.
2. Boileau, J.; Emeury, J. M. L.; Kehren, J. P. A. *Tetranitroglycoluril and Method of Preparation Thereof*, Société Nationale des Poudres et Explosifs, Paris, France, U.S. Patent 4,487,938, **1984**.
3. Boileau, J.; Carail, M.; Wimmer, E.; Gallo, R.; Pierrot, M. Dérivés nités acétylés du glycoluril, *Propellants Explos. Pyrotech* **1985**, *10*, 118–120.
4. Bastea, S.; Fired, L. E.; Glaesemann, K. R.; Howard, W. M.; Kuo, I. W.; Souers, P. C.; Vitello, P. A. *Cheetah 7.0 User's Manual*, LLNL-SM-599073, Lawrence Livermore National Laboratory, Livermore, CA, **2012**.
5. Chapman, R. D.; Quintana, R. L.; Baldwin, L. C.; Hollins, R. A. Cyclic Dinitroureas as Self-Remediating Munition Charges. SERDP Project WP-1624 Final Report February **2009**.
6. Merritt, A. R.; Hoang, H. Q.; Young, K. J.; Baldwin, L. C.; Ferguson, B. P.; Chapman, R. D.; Quintana, R. L. *Self Remediating Explosives Based on Tetranitroglycoluril (TNGU) 5th JANNAF Combustion, 33rd Airbreathing Propulsion, 33rd Exhaust Plume and Signatures, 27th Propulsion Systems Hazards Joint Subcommittee Meeting*. Hyatt Regency Monterey, CA, 3–6 December **2012**.
7. Sherrill, W. M.; Johnson, E. C.; Paraskos, A. J. Synthesis and Characterization of Mono-, Di-, and Tetranitrated 7,8-Disubstituted Glycolurils. *Propellants Explos. Pyrotech.* ASAP **23 August 2013**.
8. Kony, M.; Dagley, I. J. Synthesis of Octahydro-2,5-bis(nitrimino)imidazo-[4,5-d]imidazole. *Heterocycles* **1994**, *38*, 595–600.
9. Johnson, E. C.; Sherrill, W. M. *Evaluation of Sensitivity and Explosive Performance Estimates of N,N'-(Tetrahydroimidazo[4,5-D] Imidazole-2,5(1H,3H)-Diylidene)Dinitramide and 5-(Nitroimino)Hexahydroimidazo [4,5-d]Imidazol-2(1H)-Iminium Nitrate*; ARL-TR-6524; U.S. Army Research Laboratory: Aberdeen Proving Ground, MD, July **2013**.
10. Byrd, E. F. C.; Rice, B. M., Improved Prediction of Heats of Formation of Energetic Materials Using Quantum Mechanical Calculations. *J. Phys. Chem. A* **2006**, *110*, 1005–1013.
11. Rice, B. M.; Pai, S. V.; Hare, J. Predicting Heats of Formation of Energetic Materials Using Quantum Mechanical Calculations. *Combust. Flame* **1999**, *118*, 445–458.

12. Rice, B. M.; Hare, J. J.; Byrd, E. F. C. Accurate Predictions of Crystal Densities Using Quantum Mechanical Molecular Volumes. *J. Phys. Chem A* **2007**, *111*, 10874.
13. Byrd, E. F. C.; Rice, B. M. A Comparison of Methods to Predict Solid Phase Heats of Formation of Molecular Energetic Salts. *J. Phys. Chem. A* **2009**, *113*, 345.
14. Rice, B. M.; Byrd, E. F. C. Evaluation of Electrostatic Descriptors for Predicting Crystalline Density. *J. Comput. Chem* **2013**, *34*, 2146–2151.
15. McKay, A. F.; Wright, G. F. The Nitration Products of 2-Nitramino-D2-1,3-diazacycloalkenes. *J. Am. Chem. Soc.* **1948**, *70*, 3990.
16. Langlie, H. J. A Reliability Test for “One-Shot” Items. Technical Report U-1792 Aeronutronic Division of Ford Motor Company, Newport Beach, CA, USA, **1965**.

Appendix. FTIR Data

This appendix appears in its original form, without editorial change.



C:\Program Files\OPUS_65\MEAS\MEAS\FLOWCELL\NNO 1.0

NNO 1

Sample Compartment

27/06/2013

List of Symbols, Abbreviations, and Acronyms

Ac ₂ O	acetic anhydride
ATR	attenuated total reflectance
AU	absorbance units
DCM	dichloromethane
D _v	detonation velocity
ESD	electrostatic discharge
FTIR	Fourier transform infrared
H ₂ SO ₄	sulfuric acid
H ₅₀	50% initiation height
HHTDD	2,6-dioxo-1,3,4,5,7,8-hexanitrodecahydro-1H,5H-diimidazo[4,5-b:4',5'-e]pyrazine
HMX	octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine
HNO ₃	nitric acid
in	inch
IR	infrared
MHz	megahertz
min	minute
mL	milliliter
N	newton
N ₂	nitrogen
N ₂ O	dinitrogen oxide
NMR	nuclear magnetic resonance
OB	oxygen balance
P _{cj}	Chapman-Jouguet pressure
RDX	1,3,5-trinitroperhydro-1,3,5-triazine

SPM	surface potential map
TFAA	trifluoroacetic anhydride
TNGU	tetranitroglycoluril
UXO	unexploded ordinance
ZnSe	zinc selenide
ΔH_f	heat of formation
ΔH_d	heat of detonation
α	designates the carbonyl carbon of the imidazo imidazole
ρ	density

NO. OF
COPIES ORGANIZATION

1 DEFENSE TECHNICAL
(PDF) INFORMATION CTR
DTIC OCA

1 DIRECTOR
(PDF) US ARMY RESEARCH LAB
IMAL HRA

1 DIRECTOR
(PDF) US ARMY RESEARCH LAB
RDRL CIO LL

1 GOVT PRINTG OFC
(PDF) A MALHOTRA

2 WEAPONS DEV & INTEGRATION
(PDF) DIRCTRT AMRDEC
RDMR WDN
J NEIDERT
G DRAKE

1 US ARMY ARDEC
(PDF) AMSRD AAR AEE W
R DAMAVARAPU

1 LOS ALAMOS NATL LAB
(PDF) D CHAVEZ

1 LAWRENCE LIVERMORE NATL LAB
(PDF) P PAGORIA

3 NAVAL SURFACE WARFARE CTR
(PDF) A PEARSALL
B SLEADD
P CARUANA

NO. OF
COPIES ORGANIZATION

3 NAVAL AIR WARFARE CTR
(PDF) CODE 474100D
A MERRITT
H HUANG
K YOUNG

ABERDEEN PROVING GROUND

14 DIR USARL
(PDF) RDRL CI
P PLOSTINS
RDRL WM
P BAKER
B FORCH
RDRL WML
M ZOLTOSKI
RDRL WMM
J ZABINSKI
RDRL WML B
E BYRD
B RICE
N TRIVEDI
RDRL WML C
S AUBERT
K MCNESBY
B ROOS
W SHERRILL
K SPANGLER
RDRL WML H
J NEWILL

INTENTIONALLY LEFT BLANK.