

Beyond the Spin Model Approximation for Ramsey Spectroscopy

A. P. Koller,¹ M. Beverland,² A. V. Gorshkov,³ and A. M. Rey¹

¹*JILA, NIST, and Department of Physics, University of Colorado Boulder, Colorado 80309, USA*

²*Institute for Quantum Information and Matter, California Institute of Technology, MC 305-16, Pasadena, California 91125, USA*

³*Joint Quantum Institute, NIST, and University of Maryland, College Park, Maryland 20742, USA*

(Received 3 December 2013; revised manuscript received 31 January 2014; published 26 March 2014)

Ramsey spectroscopy has become a powerful technique for probing nonequilibrium dynamics of internal (pseudospin) degrees of freedom of interacting systems. In many theoretical treatments, the key to understanding the dynamics has been to assume the external (motional) degrees of freedom are decoupled from the pseudospin degrees of freedom. Determining the validity of this approximation—known as the spin model approximation—has not been addressed in detail. Here we shed light in this direction by calculating Ramsey dynamics exactly for two interacting spin-1/2 particles in a harmonic trap. We focus on *s*-wave-interacting fermions in quasi one- and two-dimensional geometries. We find that in one dimension the spin model assumption works well over a wide range of experimentally relevant conditions, but can fail at time scales longer than those set by the mean interaction energy. Surprisingly, in two dimensions a modified version of the spin model is exact to first order in the interaction strength. This analysis is important for a correct interpretation of Ramsey spectroscopy and has broad applications ranging from precision measurements to quantum information and to fundamental probes of many-body systems.

DOI: 10.1103/PhysRevLett.112.123001

PACS numbers: 32.30.-r, 06.30.Ft, 34.20.Cf

Ramsey spectroscopy, a technique initially designed to interrogate microwave atomic clocks, has become an important modern tool for probing dynamics of interacting many-body systems with internal (pseudospin) degrees of freedom. Ramsey spectroscopy applies [see Fig. 1(a)] two strong resonant pulses to a system initially prepared in a well-defined pseudospin state, separated by a dark time of free evolution. The first pulse initializes the pseudospin dynamics by preparing the system in a nontrivial superposition of eigenstates; i.e., it introduces a quantum quench [1]. The second pulse reads the coherence or correlations developed during the dark time. Recently, Ramsey spectroscopy has been proposed for extracting real-space and time correlations [2–6], characterizing topological order [7,8], measuring spin diffusion dynamics in bosonic [9–14] and fermionic systems [15–18], and as a means to probe many-body interactions in atomic, molecular, and trapped ion systems [19–27].

Generally speaking, Ramsey spectroscopy measures the collective pseudospin and traces out other external degrees of freedom involved during the free evolution. In most atomic setups the latter are associated with motional degrees of freedom in the harmonic trapping potential and/or lattice potential confining the atoms. The external degrees of freedom can affect the spin dynamics in a nontrivial way, however. A great simplification could be gained if it were possible to decouple the motional and spin degrees of freedom, and reduce the many-body dynamics down to those extracted from a pure interacting spin model. Evidence that this scenario is possible, even far from quantum degeneracy, has been reported in recent

experiments [9,13,14,25–28], where the observed spin dynamics corresponded to those of a pure spin Hamiltonian. These observations are opening a path for the investigation of quantum magnetism in atomic systems without the need for ultralow temperatures. It is thus important to determine the parameter regime in which a pure interacting-spins picture is valid.

In this Letter we provide insight on the validity of a pure spin model description of Ramsey spectroscopy by performing exact calculations for fermions with *s*-wave interactions and an internal pseudospin-1/2 degree of freedom, confined in quasi-1D and quasi-2D harmonic traps. We show that the large degeneracy of the harmonic oscillator spectrum can limit the validity of the spin model to time scales less than the inverse interaction strength, due to resonant collisionally induced excitation of spatial modes [see Fig. 1(b)]. Cold atom experiments are protected from this problem if the temperature is high enough that atoms probe the actual Gaussian shape of the potential which breaks the harmonic spectrum degeneracy. This was shown to be the case for




FIG. 1 (color online). (a) Ramsey spectroscopy of two interacting spin-1/2 particles. (b) In a harmonic trap the spectrum degeneracy allows near-resonant mode-changing collisions coupled to the spin dynamics.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 26 MAR 2014		2. REPORT TYPE		3. DATES COVERED 00-00-2014 to 00-00-2014	
4. TITLE AND SUBTITLE Beyond the Spin Model Approximation for Ramsey Spectroscopy				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) California Institute of Technology, Institute for Quantum Information and Matter, Pasadena, CA, 91125				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

example in Refs. [9,25,26] where a pure spin model well described the experimental observations. At very low temperatures, Pauli blocking can also prevent mode changing collisions, as recently observed in Ref. [29]. However, the degeneracy is a concern for intermediate temperatures at which the set of populated levels is effectively harmonic. Here we show that surprisingly, in two dimensions and to first order in the interaction strength, the full two-particle dynamics can be described in terms of an effective spin model with appropriate parameters. Our two-body calculations are not only a first step towards understanding the interplay between spin and particle motion in generic many-body ensembles, but are also directly applicable to optical clocks that interrogate an array of 1D tube-shaped traps, each with fewer than three atoms [28,30,31].

Physical situation.—Consider two fermions with internal degrees of freedom $\{\uparrow, \downarrow\}$ corresponding, for instance, to the $^1S_0 - ^3P_0$ electronic levels in alkaline-earth-based optical lattice clocks, and assume their interactions are primarily described by an s -wave pseudopotential. The atoms are also illuminated by a laser beam detuned by $\delta = \omega_L - \omega_0$ from the atomic transition ω_0 , with wave vector $\vec{k}$ and bare Rabi frequency Ω . The two-particle Hamiltonian is then given by $\hat{H}(\vec{x}_1, \vec{x}_2) = \sum_{i=1,2} \hat{H}_L(\vec{x}_i) + \hat{H}_D(\vec{x}_1, \vec{x}_2)$:

$$\begin{aligned} \hat{H}_L(\vec{x}_i) &= -\frac{\hbar\Omega}{2} e^{-i(\omega_L t - \vec{k} \cdot \vec{x}_i)} \hat{\sigma}_i^+ + \text{H.c.}, \\ \hat{H}_D(\vec{x}_1, \vec{x}_2) &= H_{sp}(\vec{x}_1) + H_{sp}(\vec{x}_2) + g\hat{P}_s \delta(\vec{r}) \frac{\partial}{\partial r} r. \end{aligned} \quad (1)$$

Here, $\hat{H}_L(\vec{x}_i)$ describes the atom-laser interaction: $\hat{\sigma}_i^+$ is the spin raising operator acting on atom i , and H.c. is the Hermitian conjugate. $H_{sp}(\vec{x}_i) = -\hbar^2/(2M)\nabla_i^2 + V(\vec{x}_i) + (\hbar\omega_0/2)\hat{\sigma}_i^z$ is the single particle Hamiltonian with an external potential V , assumed for simplicity to be independent of the internal state and separable. $H_{sp}(\vec{x}_i)$ has eigenfunctions $\phi_{\mathbf{n}}(\vec{x}_i)$ and eigenenergies $E_{\mathbf{n}}$ with $\mathbf{n} = \{n_x, n_y, n_z\}$. M is the particle's mass and $\hat{\sigma}^z$ the Pauli matrix. $\vec{r} = \vec{x}_1 - \vec{x}_2$ is the relative coordinate, $g = 4\pi\hbar^2 a_s^{\uparrow\downarrow}/M$ and $a_s^{\uparrow\downarrow}$ the 3D s -wave scattering length. $\hat{P}_s = |s\rangle\langle s|$ is the projector into the singlet state, $|s\rangle = (1/\sqrt{2})(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$. Only fermions in the singlet state can interact, while spin triplet states, $|t_{\downarrow\downarrow}\rangle = |\downarrow\downarrow\rangle$, $|t_{\downarrow\uparrow}\rangle = (1/\sqrt{2})(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$, and $|t_{\uparrow\uparrow}\rangle = |\uparrow\uparrow\rangle$ cannot experience s -wave interactions.

The spin model.—The assumptions of the spin model are: if there are no degeneracies in the two-atom non-interacting spectrum, i.e., $(E_{\mathbf{m}} + E_{\mathbf{n}}) = (E_{\mathbf{m}'} + E_{\mathbf{n}'})$ occurs only for $(\mathbf{m}, \mathbf{n}) = (\mathbf{m}', \mathbf{n}')$ or $(\mathbf{m}, \mathbf{n}) = (\mathbf{n}', \mathbf{m}')$, and interactions are treated as a perturbation, scattering processes that change the single-particle modes become off resonant and atoms remain frozen during the dynamics. In this case interactions are diagonal in the single-particle

basis and for particles in modes $(\mathbf{m}, \mathbf{n})$ they are fully characterized by the interaction energy

$$\hbar U_{\uparrow\downarrow}^{nm} = g \int d^3\vec{x} |\phi_{\mathbf{n}}(\vec{x})|^2 |\phi_{\mathbf{m}}(\vec{x})|^2. \quad (2)$$

Fermions with s -wave interactions in one dimension.—

We begin with the case of two atoms tightly confined transversally in their ground state and with dynamics only along the z direction, where they experience a 1D harmonic trapping potential with angular trapping frequency ω_z . The two atoms are initially prepared in the state $(1/\sqrt{2})(|n_1, n_2\rangle - |n_2, n_1\rangle)|t_{\downarrow\downarrow}\rangle$.

The atoms are assumed to be in the Lamb-Dicke regime, with Lamb-Dicke parameter $\eta = k_z a_{ho}/\sqrt{2} \ll 1$. $a_{ho} = \sqrt{\hbar/M\omega_z}$ is the harmonic oscillator length, and k_z the projection of the probe laser wave vector along z . Mode changes during the laser interrogation can be suppressed if the laser detuning from the atomic transition δ and the bare Rabi frequency Ω satisfy $\delta, \eta\Omega \ll \omega_z$. In this regime the mode dependence of the Rabi frequencies is $\Omega_n = \Omega e^{-(\eta^2/2)} L_n^0(\eta^2)$ [32]. The Hamiltonian in the rotating frame of the laser [21–23,25] under the spin model approximation can be written as $\hat{H}_{sm}^{n_1, n_2} = \hat{H}_L^{n_1, n_2} + \hat{H}_D^{n_1, n_2}$, where

$$\begin{aligned} \hat{H}_L^{n_1, n_2} &= \hbar\Delta\Omega^{n_1, n_2} \frac{(\hat{\sigma}_1^x - \hat{\sigma}_2^x)}{2} - \hbar\bar{\Omega}^{n_1, n_2} \hat{s}_x, \\ \hat{H}_D^{n_1, n_2} &= 2\hbar u_{\uparrow\downarrow}^{n_1, n_2} \hat{P}_s - \hbar\delta\hat{s}_z. \end{aligned} \quad (3)$$

$\hat{H}_L^{n_1, n_2}$ acts only during the two laser pulses, and $\hat{H}_D^{n_1, n_2}$ acts only during the dark time. Here, $\hat{s}_{x,y,z} = (\hat{\sigma}_1^{x,y,z} + \hat{\sigma}_2^{x,y,z})/2$ are collective spin operators and $\bar{\Omega}^{n_1, n_2} = (\Omega_{n_1} + \Omega_{n_2})/2$ is the mean Rabi frequency. $\Delta\Omega^{n_1, n_2} = (\Omega_{n_1} - \Omega_{n_2})/2$ arises from the excitation inhomogeneity and can transfer some of the initial triplet population to the singlet, allowing interactions. The interaction energy $\hbar u_{\uparrow\downarrow}^{n_1, n_2} = \hbar U_{\uparrow\downarrow}^{nm}$ in Eq. (2) with $\mathbf{n} = \{0, 0, n_1\}$ and $\mathbf{m} = \{0, 0, n_2\}$. We can ignore the detuning and interactions during the laser pulses if the pulses are short compared to the time scales set by those energies. We also ignore single-particle energies which are constants and do not contribute to the dynamics.

The spin model assumptions break down in a harmonic trap due to the degeneracy of the noninteracting two-atom spectrum: even weak interactions can transfer atoms initially in modes $\{n_1, n_2\}$ to the various degenerate configurations $\{n_1 + k, n_2 - k\}$ (for integer k) during the dynamics. To account for these mode changes, we take advantage of the exact eigenfunctions and eigenvalues of $\hat{H}_D(\vec{x}_1, \vec{x}_2)$ in Eq. (1) for two atoms with s -wave interactions in a harmonic trap [33]. These solutions exploit the separability of the Hamiltonian in the center-of-mass coordinate R and relative coordinate r . There is no degeneracy in the relative coordinate degree of

freedom. See Supplemental Material [34] for straightforward expressions for the change of basis. Equivalent expressions are given in [35].

Ramsey dynamics in the spin model approximation.—Denoting τ the Ramsey dark time, the population difference between the two spin states measured after the second pulse takes the generic form

$$\langle \hat{s}_z \rangle(\tau) = A(\tau) \cos(\delta\tau) + B(\tau) \sin(\delta\tau) + C(\tau). \quad (4)$$

$A(\tau)$, $B(\tau)$, and $C(\tau)$ have the form $A(\tau) = I_1(\tau)f_1 + f_2$, $B(\tau) = I_2(\tau)f_3$, $C(\tau) = I_3(\tau)f_4 + f_5$, where $I_i(\tau)$ depend on the dark time physics, and f_i are independent of the dark time physics and depend only on the laser pulse quantities $\{\Delta\theta_{j=1,2}^{n_1,n_2}, \bar{\theta}_{j=1,2}^{n_1,n_2}\}$ (see Supplemental Material [34]). $\Delta\theta_j^{n_1,n_2} = \Delta\Omega^{n_1,n_2}t_j$ and $\bar{\theta}_j^{n_1,n_2} = \bar{\Omega}^{n_1,n_2}t_j$, with $t_{1,2}$ the pulse durations. In the spin model approximation, the dark time functions depend simply on interactions: $I_1^{\text{sm}} = I_3^{\text{sm}} = \cos(u_{\uparrow\downarrow}^{n_1,n_2}\tau)$, $I_2^{\text{sm}} = \sin(u_{\uparrow\downarrow}^{n_1,n_2}\tau)$.

Ramsey dynamics in the weakly interacting regime ($u_{\uparrow\downarrow}^{1,0} \ll \omega_z$).—For weakly interacting atoms ($u_{\uparrow\downarrow}^{1,0} \ll \omega_z$), we are able to write the dynamics (beyond the spin model approximation) in a closed analytic form [34]. These expressions for the dynamics are exact for times $\tau \ll \omega_z/(u_{\uparrow\downarrow}^{1,0})^2$:

$$I_1^{\text{exact}} = I_3^{\text{exact}} = 2 \sum_{n_r=0, \text{even}}^{n_1+n_2} |d_{n_r}^{n_1,n_2}|^2 \cos \left[\frac{\Delta E_s(n_r)}{\hbar} \tau \right],$$

$$I_2^{\text{exact}} = 2 \sum_{n_r=0, \text{even}}^{n_1+n_2} |d_{n_r}^{n_1,n_2}|^2 \sin \left[\frac{\Delta E_s(n_r)}{\hbar} \tau \right]. \quad (5)$$

Here, $d_{n_r}^{n_1,n_2}$ are the change of basis coefficients defined in Supplemental Material [34]. Comparing Eq. (5) to the spin model solution, we see the single frequency $u_{\uparrow\downarrow}^{n_1,n_2}$ in the spin-model dynamics gets replaced by a sum over many frequencies $\Delta E_s(n_r)/\hbar$ in the exact dynamics. These frequencies are associated with the first order correction of the eigenenergies due to interactions [33]: $\Delta E_s(n_r) = \hbar u_{\uparrow\downarrow}^{1,0} (\Gamma(n_r/2 + 1/2)/\sqrt{\pi} \Gamma(n_r/2 + 1)) (1 + \mathcal{O}(u_{\uparrow\downarrow}^{1,0}/\omega_z))$. The many frequencies that appear come from the resonant mode-changing processes. States with odd n_r do not experience s -wave interactions and do not contribute.

When we compare the exact dynamics to those predicted by the spin model we find that they agree for short times, $\tau u_{\uparrow\downarrow}^{1,0} \ll 1$. The spin model fails at longer times, however, when leakage of the population to other modes in the individual-particle coordinate basis becomes significant [See Fig. 2(a)]. This is reflected in the behavior of the angular frequency shift $\Delta\omega(\tau)$ —an important quantity for atomic clock experiments—defined as $\Delta\omega(\tau)\tau = -\arctan[B(\tau)/A(\tau)]$, which is the observed change in the atomic transition due to interactions [see Fig. 3(a)]. The failure of the spin model at




FIG. 2 (color online). Ramsey dynamics [see Eq. (4)] with $\delta = 0$: (a) 1D spin model, exact solution, and projection of population onto initial mode (here $n_1 = 10$ and $n_2 = 0$), with $u_{\uparrow\downarrow}^{1,0} \approx 0.2\omega_z$. Dephasing of the exact dynamics results from mode changes. (b) Thermal averages in two dimensions: spin model vs effective spin model, at different temperatures, with $u_{\uparrow\downarrow}^{1,0} \approx 0.04\omega_z$. For both figures: $\theta_1 = \theta_2 = \pi/3$, with thermally averaged inhomogeneity $\langle \Delta\Omega \rangle / \langle \Omega \rangle = 0.3$. $\theta_i = \Omega t_i$ are bare pulse areas.

times longer than the inverse interaction strength limits its applicability to model the new generation of atomic clocks that use ultra coherent lasers [36,37], allowing interrogation times exceeding a few seconds. A spin model treatment will be insufficient when conditions are such that the atoms see an almost purely harmonic potential.

Ramsey dynamics in the strongly interacting regime ($u_{\uparrow\downarrow}^{1,0} \gtrsim \omega_z$).—The spin model fails when $u_{\uparrow\downarrow}^{1,0} \gtrsim \omega_z$. To maintain the separation between interaction-induced effects and laser-induced effects, we imagine interactions set to be weak during the laser pulses and suddenly increased after the first pulse using, for example, a Feshbach resonance [25,38,39] [40]. For this situation, we can solve for the dynamics, given an initial pair of modes, although there is no closed form solution (the dark time functions $I_i(\tau)$ are more complicated, but the laser dependence through f_j remains the same as in the previous cases). We find that, in the limit of strong interactions ($u_{\uparrow\downarrow}^{1,0} \gg \omega_z$), the population imbalance exhibits periodic oscillations at the axial trapping frequency ω_z , in contrast with the spin model prediction of much faster oscillations at the interaction frequency (see Supplemental Material [34]). The frequency shift (proportional to this oscillation frequency), saturates to a value on the order of ω_z , instead of increasing without




FIG. 3 (color online). Thermally averaged frequency shifts: (a) 1D spin model and exact solution vs population excitation fraction (number of atoms in $\uparrow$ divided by the total number of atoms) after the first pulse, at intermediate and long times. Here, $u_{\uparrow\downarrow} \equiv u_{\uparrow\downarrow}^{1,0} \approx 0.2\omega_z$. (b) Frequency shifts for 2D spin model vs effective spin model, with $u_{\uparrow\downarrow} \equiv u_{\uparrow\downarrow}^{1,0} \approx 0.04\omega_z$. For both figures: $T = 208\hbar\omega_z/k_B$, $\theta = \theta_1 = \theta_2$, $\omega_z = \omega_{\perp} = 700 \times 2\pi$ Hz, and thermally averaged inhomogeneity $\langle\Delta\Omega\rangle/\langle\Omega\rangle = 0.3$. $\theta_i = \Omega t_i$ are bare pulse areas.

bound. These results reflect the fact that for strong interactions (unitarity), the fermions maximally repel each other, and the trap energy becomes the only relevant energy scale in the system. This behavior, expected to be a universal result, should apply even in the many-body case as seen in Refs. [19,20].

References [21,25] showed that s -wave frequency shifts can be cancelled by setting the second pulse area to $\bar{\theta}_2^{n_1, n_2} = \pi/2$. This result, obtained using the spin model, survives the inclusion of resonant mode changes even for strong interactions during the dark time, since the dependence of the dynamics on the functions f_i , and thus $\bar{\theta}_2^{n_1, n_2}$, remains the same even when interactions are strong.

Fermions with s -wave interactions in two dimensions.—For an anisotropic 2D harmonic potential with no accidental degeneracies, the treatment will be similar to the 1D case. An isotropic 2D harmonic potential, however, is more difficult to treat, due to the large degeneracy. In two dimensions the spin model remains the same as Eq. (3), with populated modes now $n_i = \{n_{xi}, n_{yi}, 0\}$, and interaction energy $\hbar u_{\uparrow\downarrow}^{\vec{n}_1, \vec{n}_2} = \hbar U_{\uparrow\downarrow}^{n_1, n_2}$ in Eq. (2). To go beyond the

spin model we use polar relative coordinates to eliminate much of the degeneracy. For noninteracting particles, the eigenfunctions can be parametrized by quantum numbers n and m , with energy $E = \hbar\omega_{\perp}(2n + |m| + 1)$ and angular momentum component $L_z = \hbar m$, where $\omega_{\perp}$ is the 2D oscillator frequency. S -wave interactions only affect states with $m = 0$, and this subset of states contains no degeneracy (other than the center-of-mass degeneracy). To first order in perturbation theory the interaction energy shift is independent of the radial quantum number n : $\Delta E = (g/2\sqrt{2}\pi^{3/2}a_z a_{\perp}^2)(1 + \mathcal{O}(u_{\uparrow\downarrow}^{1,0}/\omega_{\perp}))$, where a_z and $a_{\perp}$ are the oscillator lengths along the tightly confined z direction and the weakly confined x and y directions, respectively, and in 2D $u_{\uparrow\downarrow}^{1,0} \equiv u_{\uparrow\downarrow}^{\vec{n}_1, \vec{n}_2}$ with $\vec{n}_1 = (1, 0, 0)$, $\vec{n}_2 = (0, 0, 0)$. This result is striking: despite the large degeneracy in two dimensions, each interacting state with $m = 0$ receives the same energy shift to first order in perturbation theory, and accumulates the same phase during the dark time. An effective spin model, with diagonal matrix element $2\hbar u_{\uparrow\downarrow}^{\vec{n}_1, \vec{n}_2}$ replaced by ΔE , will be *exact* for the $m = 0$ states, to first order in the interaction strength. We can replace $\hat{H}_D^{n_1, n_2}$ in Eq. (3) with

$$\hat{H}_{D, \text{esm}}^{\vec{n}_1, \vec{n}_2} = \Delta E \hat{P}_{m=0} - \hbar \delta \delta_z, \quad (6)$$

where $\hat{P}_{m=0}$ projects onto interacting states with $m = 0$. For a properly symmetrized initial state $\Psi_{\mathbf{n}_1, \mathbf{n}_2}(\vec{x}_1, \vec{x}_2)$ in modes $(\mathbf{n}_1, \mathbf{n}_2)$, we denote the fraction of the population with $m = 0$ in the relative coordinate by $\mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0}$, which can be calculated as $\mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0} = \int d^3\vec{x}_1 d^3\vec{x}_2 |\Psi_{\mathbf{n}_1, \mathbf{n}_2}(\vec{x}_1, \vec{x}_2)|^2 \times \delta(\vec{x}_1 - \vec{x}_2) = (4\sqrt{2}\pi^{3/2}/g)\hbar u_{\uparrow\downarrow}^{\vec{n}_1, \vec{n}_2}$, where $u_{\uparrow\downarrow}^{\vec{n}_1, \vec{n}_2}$ is the 2D interaction energy calculated from Eq. (2). The dark time dynamics of this effective spin model are simple: $I_1^{\text{esm}} = I_3^{\text{esm}} = (1 - \mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0}) + \mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0} \cos(\Delta E \tau / \hbar)$, $I_2^{\text{esm}} = \mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0} \sin(\Delta E \tau / \hbar)$.

In the original spin model, $\hbar u_{\uparrow\downarrow}^{\vec{n}_1, \vec{n}_2}$ is used as the interaction energy. We see that this parameter appears in the effective spin model to quantify the population of interacting modes ($\mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0}$), instead of their energy. This dramatic result is seen in Fig. 2(b), comparing thermal averages of the previously implemented spin model with the new effective spin model. Oscillations during the dynamics remain at the same frequency ΔE at higher temperatures, but the amplitude of the oscillations, proportional to $\mathcal{P}_{\mathbf{n}_1, \mathbf{n}_2}^{m=0}$, decreases. The previously implemented spin model, on the other hand, predicts smaller interaction energies (slower oscillations) at higher temperatures. The frequency shift predicted by the original spin model is only valid at short times [see Fig. 3(b)].

Summary and outlook.—We test the validity of a spin model treatment for Ramsey spectroscopy with exact calculations for two pseudospin-1/2 fermions in a harmonic

trap. In one dimension the spin model treatment breaks down for dark times on the order of the inverse interaction strength, and for strong interactions. In two dimensions we find an effective spin model which is exact to first order in perturbation theory, and whose dynamics can be quite different from those predicted by a spin model treatment. Future theoretical treatments of interacting systems probed by Ramsey spectroscopy must take these effects into account to correctly describe dynamics outside of the short-time and weakly interacting regimes.

We thank M. Foss-Feig, G. Campbell, K. Hazzard, A. Ludlow, J. von Stecher, K. O'Hara, M. Martin, and the JILA Sr clock experimental team for feedback. This work was supported by NSF-PIF, NSF JILA-PFC-1125844, NSF JQI-PFC-0822671, NSF IQIM-PFC-1125565, ARO-DARPA-OLE, AFOSR, NIST, and the Lee A. DuBridge and Gordon and Betty Moore Foundations. A. K. was supported by the Department of Defense through the NDSEG program.

-
- [1] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, *Rev. Mod. Phys.* **83**, 863 (2011).
- [2] M. Knap, A. Kantian, T. Giamarchi, I. Bloch, M. D. Lukin, and E. Demler, *Phys. Rev. Lett.* **111**, 147205 (2013).
- [3] T. Kitagawa, S. Pielawa, A. Imambekov, J. Schmiedmayer, V. Gritsev, and E. Demler, *Phys. Rev. Lett.* **104**, 255302 (2010).
- [4] M. Knap, A. Shashi, Y. Nishida, A. Imambekov, D. A. Abanin, and E. Demler, *Phys. Rev. X* **2**, 041020 (2012).
- [5] A. Widera, O. Mandel, M. Greiner, S. Kreim, T. W. Hansch, and I. Bloch, *Phys. Rev. Lett.* **92**, 160406 (2004).
- [6] A. Kuklov, N. Prokof'ev, and B. Svistunov, *Phys. Rev. A* **69**, 025601 (2004).
- [7] M. Atala, M. Aidelsburger, J. T. Barreiro, D. Abanin, T. Kitagawa, E. Demler, and I. Bloch, [arXiv:1212.0572](https://arxiv.org/abs/1212.0572).
- [8] D. A. Abanin, T. Kitagawa, I. Bloch, and E. Demler, *Phys. Rev. Lett.* **110**, 165304 (2013).
- [9] H. K. Pechkis, J. P. Wrubel, A. Schwettmann, P. F. Griffin, R. Barnett, E. Tiesinga, and P. D. Lett, *Phys. Rev. Lett.* **111**, 025301 (2013).
- [10] J. M. McGuirk, H. J. Lewandowski, D. M. Harber, T. Nikuni, J. E. Williams, and E. A. Cornell, *Phys. Rev. Lett.* **89**, 090402 (2002).
- [11] D. M. Harber, H. J. Lewandowski, J. M. McGuirk, and E. A. Cornell, *Phys. Rev. A* **66**, 053616 (2002).
- [12] H. J. Lewandowski, D. M. Harber, D. L. Whitaker, and E. A. Cornell, *Phys. Rev. Lett.* **88**, 070403 (2002).
- [13] C. Deutsch, F. Ramirez-Martinez, C. Lacroûte, F. Reinhard, T. Schneider, J. N. Fuchs, F. Piéchon, F. Laloë, J. Reichel, and P. Rosenbusch, *Phys. Rev. Lett.* **105**, 020401 (2010).
- [14] W. Maineult, C. Deutsch, K. Gibble, J. Reichel, and P. Rosenbusch, *Phys. Rev. Lett.* **109**, 020407 (2012).
- [15] X. Du, L. Luo, B. Clancy, and J. E. Thomas, *Phys. Rev. Lett.* **101**, 150401 (2008).
- [16] F. Piéchon, J. N. Fuchs, and F. Laloë, *Phys. Rev. Lett.* **102**, 215301 (2009).
- [17] S. S. Natu and E. J. Mueller, *Phys. Rev. A* **79**, 051601 (2009).
- [18] M. Koschorreck, D. Pertot, E. Vogt, and M. Khl, *Nat. Phys.* **9**, 405 (2013).
- [19] M. W. Zwierlein, Z. Hadzibabic, S. Gupta, and W. Ketterle, *Phys. Rev. Lett.* **91**, 250404 (2003).
- [20] S. Gupta, Z. Hadzibabic, M. W. Zwierlein, C. A. Stan, K. Dieckmann, C. H. Schunck, E. G. M. van Kempen, B. J. Verhaar, and W. Ketterle, *Science* **300**, 1723 (2003).
- [21] K. Gibble, *Phys. Rev. Lett.* **103**, 113202 (2009).
- [22] A. M. Rey and A. V. Gorshkov, and C. Rubbo, *Phys. Rev. Lett.* **103**, 260402 (2009).
- [23] Z. H. Yu and C. J. Pethick, *Phys. Rev. Lett.* **104**, 010801 (2010).
- [24] J. W. Britton, B. C. Sawyer, A. C. Keith, C. C. J. Wang, J. K. Freericks, H. Uys, M. J. Biercuk, and J. J. Bollinger, *Nature (London)* **484**, 489 (2012).
- [25] E. L. Hazlett, Y. Zhang, R. W. Stites, K. Gibble, and K. M. O'Hara, *Phys. Rev. Lett.* **110**, 160801 (2013).
- [26] M. J. Martin, M. Bishof, M. D. Swallows, X. Zhang, C. Benko, J. von Stecher, A. V. Gorshkov, A. M. Rey, and J. Ye, *Science* **341**, 632 (2013).
- [27] B. Yan, S. A. Moses, B. Gadway, J. P. Covey, K. R. A. Hazzard, A. M. Rey, D. S. Jin, and J. Ye, *Nature (London)* **501**, 521 (2013).
- [28] M. D. Swallows, M. Bishof, Y. G. Lin, S. Blatt, M. J. Martin, A. M. Rey, and J. Ye, *Science* **331**, 1043 (2011).
- [29] J. Krauser, U. Ebling, N. Flaschner, J. Heinze, K. Sengstock, M. Lewenstein, A. Eckardt, and C. Becker, [arXiv:1307.8392](https://arxiv.org/abs/1307.8392).
- [30] N. D. Lemke, J. von Stecher, J. A. Sherman, A. M. Rey, C. W. Oates, and A. D. Ludlow, *Phys. Rev. Lett.* **107**, 103902 (2011).
- [31] N. Hinkley, J. A. Sherman, N. B. Phillips, M. Schioppo, N. D. Lemke, K. Beloy, M. Pizzocaro, and C. W. Oates and A. D. Ludlow, *Science* **341**, 1215 (2013).
- [32] G. K. Campbell, M. M. Boyd, J. W. Thomsen, M. J. Martin, S. Blatt, M. D. Swallows, T. L. Nicholson, T. Fortier, C. W. Oates, S. A. Diddams, N. D. Lemke, P. Naidon, P. Julienne, J. Ye, and A. D. Ludlow, *Science* **324**, 360 (2009).
- [33] T. Busch, B. G. Englert, K. Rzazewski, and M. Wilkens, *Found. Phys.* **28**, 549 (1998).
- [34] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.112.123001> for change of basis formulas, derivation of Ramsey dynamics, and further discussion of strong interactions.
- [35] G. Quémener and J. L. Bohn, *Phys. Rev. A* **83**, 012705 (2011).
- [36] T. L. Nicholson, M. J. Martin, J. R. Williams, B. J. Bloom, M. Bishof, M. D. Swallows, S. L. Campbell, and J. Ye, *Phys. Rev. Lett.* **109**, 230801 (2012).
- [37] B. J. Bloom, T. L. Nicholson, J. R. Williams, S. L. Campbell, M. Bishof, X. Zhang, W. Zhang, S. L. Bromley, and J. Ye, [arXiv:1309.1137](https://arxiv.org/abs/1309.1137). [*Nature (London)* (to be published)].
- [38] P. Makotyn, C. E. Klauss, D. L. Goldberger, E. A. Cornell, and D. S. Jin, [arXiv:1308.3696](https://arxiv.org/abs/1308.3696).
- [39] K. O'Hara, (private communication).
- [40] During the pulses we require $u_{\uparrow\downarrow}^{1,0} \ll \Omega$ to ignore interactions, $\eta\Omega \ll \omega_z$ to ignore laser induced mode changes, and $\eta^2\Omega \gg u_{\uparrow\downarrow}^{1,0}$ to populate all of the interacting modes in the singlet. These three conditions cannot simultaneously be satisfied unless $u_{\uparrow\downarrow}^{1,0} \ll \omega_z$.