

Directionality of Flare-Accelerated α Particles at the Sun

Gerald H. Share, Ronald J. Murphy

*E.O. Hulburt Center for Space Research, Naval Research Laboratory, Washington, DC
20375 USA*

gerald.share@nrl.navy.mil, ronald.murphy@nrl.navy.mil

David M. Smith, Robert P. Lin

*Space Sciences Laboratory and Dept. of Physics, University of California, Berkeley, CA
94720, U.S.A.*

dsmith@ssl.berkeley.edu, rlin@ssl.berkeley.edu

Brian R. Dennis, Richard A. Schwartz

NASA Goddard Space Flight Center, Greenbelt, MD 20771, U.S.A.

Brian.R.Dennis@nasa.gov, richard.schwartz@gsfc.nasa.gov

ABSTRACT

RHESSI has observed the ${}^7\text{Be}$ and ${}^7\text{Li}$ γ -ray lines from fusion of accelerated α particles with ambient helium produced during the 2002 July 23 solar flare that erupted at a heliocentric angle of 73° . This is the first observation of the lines with a germanium spectrometer. We have fit the measured lines with calculated shapes for different energy and angular distributions for the interacting α particles. A particle distribution from saturated pitch-angle scattering in the corona provides the best fit, but isotropic, downward-isotropic, or fan-beam distributions also provide acceptable fits. We can rule out a downward-beamed distribution with 99.99% confidence. *RHESSI* measurements of other Doppler-shifted nuclear de-excitation lines are consistent with a forward isotropic distribution of interacting particles in a magnetic loop tilted by $\sim 40^\circ$ from the normal to the solar surface toward Earth. Such a distribution is also consistent with the α - ${}^4\text{He}$ line shape with $\sim 15\%$ confidence.

Subject headings: Sun: flares—Sun: particle emission—Sun: X-rays, gamma rays

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 2003	2. REPORT TYPE	3. DATES COVERED 00-00-2003 to 00-00-2003	
4. TITLE AND SUBTITLE Directionality of Flare-Accelerated a Particles at the Sun		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory, E. O. Hulburt Center for Space Research, 4555 Overlook Avenue, SW, Washington, DC, 20375		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited			
13. SUPPLEMENTARY NOTES Ap.J. Letters, 595, L89 (2003).			
14. ABSTRACT RHESSI has observed the 7Be and 7Li -ray lines from fusion of accelerated particles with ambient helium produced during the 2002 July 23 solar are that erupted at a heliocentric angle of 73 . This is the rst observation of the lines with a germanium spectrometer. We have t the measured lines with calculated shapes for di erent energy and angular distributions for the interacting particles. A particle distribution from saturated pitch-angle scattering in the corona provides the best t, but isotropic, downward-isotropic, or fan-beam distributions also provide acceptable ts. We can rule out a downward-beamed distribution with 99.99% con dence. RHESSI measurements of other Doppler-shifted nuclear de- excitation lines are consistent with a forward isotropic distribution of interacting particles in a magnetic loop tilted by 40 from the normal to the solar surface toward Earth. Such a distribution is also consistent with the -4He line shape with 15% con dence.			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)
			18. NUMBER OF PAGES 10
			19a. NAME OF RESPONSIBLE PERSON

1. INTRODUCTION

The shape of γ -ray lines emitted in solar flares provides information on the angular distribution of energetic particles interacting in the chromosphere. Ramaty & Crannell (1976) performed the first calculations of Doppler shifts from recoil of excited ^{16}O nuclei after impact by protons and α particles. Kozlovsky & Ramaty (1977) recognized that the shapes of γ -ray lines at 0.429 and 0.478 MeV can be used to infer the angular distributions of flare-accelerated α particles where they interact. These lines result from de-excitation of ^7Be and ^7Li , respectively, produced in fusion of accelerated α particles with ambient ^4He (Kozlovsky & Ramaty 1974). Ramaty, Kozlovsky, & Lingenfelter (1979) compiled the fundamental cross sections for determining the production of γ rays and included the kinematics to calculate the shapes and shifts of γ -ray lines. This compilation, which has recently been updated (Kozlovsky, Murphy, & Ramaty 2002), formed the basis of a Monte Carlo code used in several solar studies. Murphy, Kozlovsky, & Ramaty (1988) used this code to calculate the expected shapes of the α - ^4He lines for four different angular distributions of particles: isotropic, fan beam, broadened fan beam, and downward beam. They presented line profiles for flares at Sun center and at the limb.

Murphy et al. (1990) continued this study using a magnetic loop model for transport of the ions, including the effects of mirroring and MHD pitch-angle scattering. Pitch-angle scattering can be characterized by the average distance for particle isotropization, assumed to be independent of energy (Hua et al. 1989). With no PAS, particles with large initial pitch angles mirror and tend to interact at their mirror points where the density is greatest. Particles with pitch angles too small to mirror enter the “loss cone” and either undergo a nuclear reaction as they are moving downward or are thermalized. Without PAS, most particles are moving parallel to the solar surface at their mirror points when they interact (i.e., a “fan beam”). Scattering causes the loss cone to be continuously repopulated and results in more downward-directed interacting particles. As PAS is increased, the distribution becomes even more downward-directed until saturation is reached. At saturation, particles are scattered into the loss cone as quickly as they are removed by interactions (Hua et al. 1989). Murphy et al. (1990) also folded their calculations through the instrument response of the *Solar Maximum Mission* (*SMM*) Gamma-Ray Spectrometer (GRS) and compared the results with spectra observed from the 1981 April 27 solar flare. These comparisons provided convincing evidence for the presence of the α - ^4He fusion lines in that flare. Their fits also suggested that a downward beam of particles did not fit the data as well as an isotropic or fan-beam geometry.

Share & Murphy (1997) compared the ^7Be and ^7Li line profiles calculated by Murphy, Kozlovsky, & Ramaty (1988) with the spectra from 19 solar flares observed by the *SMM*

Gamma Ray Spectrometer. They found that a downward beam of α particles could be ruled out with high confidence in two disk flares and that isotropic and fan-beam distributions of interacting particles were both consistent with all the data. In a subsequent study in which the flare spectra were summed by heliocentric angle, Share et al. (2002) concluded that the resolution of the *SMM* NaI detector was not sufficient to unambiguously distinguish between isotropic, downward-isotropic, and fan-beam distributions of interacting α particles.

In this letter we present *RHESSI* high-resolution observations of the α - ^4He fusion lines emitted during the 2002 July 23 solar flare. We compare these observations with line-shape calculations to estimate the angular distribution of interacting α particles. We compare our results on the angular distribution with those obtained for the same flare from red shifts of de-excitation lines (Smith et al. 2003) and from the delayed time profile of the neutron-capture line (Murphy et al. 2003).

2. α -HELIUM LINE OBSERVATION AND MODELS

Lin et al. (2002) and Smith et al. (2002) discuss the *RHESSI* experiment and its spectroscopic capabilities. Soft X-ray emission from the X4.8 class flare (AR0039; S13E72) on 2002 July 23 was observed by GOES beginning at 00:18 UT and peaking at 00:34 UT. *RHESSI* observed the flare until about 01:16 UT when it was occulted by the Earth. We accumulated 960 s of spectral data from 00:27:20 to 00:43:20 UT to study the solar α - ^4He lines. We estimated the background during the flare using 960-s spectral accumulations on the previous and subsequent days (± 15 orbits) when the satellite passed over similar geographic locations (Smith et al. 2003; Share et al. 2003). In Figure 1 we plot the count spectrum observed by *RHESSI* between 370 and 600 keV, after subtracting this background. The spectrum is dominated by the positron-electron annihilation line and the bremsstrahlung continuum.

We constructed a model solar photon spectrum, passed it through the instrument response function, and fit the background-corrected data from 150 to 8500 keV (Lin et al. 2003; Share et al. 2003). This model spectrum included a double power-law bremsstrahlung spectrum, the solar annihilation line and its positronium continuum, a nuclear de-excitation line function made up of 15 narrow and broad Gaussians (Smith et al. 2003), the neutron-capture line (Murphy et al. 2003), and a model for the α - ^4He fusion line complex. We plot the calculated line shapes of this complex for three representative angular distributions of interacting α particles in Figure 2 for a flare at a heliocentric angle of 73° . The most distinctive shape is for a downward beam, for which the ^7Be and ^7Li lines would be clearly resolved by *RHESSI*. A downward-isotropic distribution (i.e. one which is isotropic in the hemisphere towards the solar surface and has no component away from the Sun) exhibits a broad struc-

ture from ~ 400 keV to 500 keV, peaking near 450 keV where the two lines sum together. At this heliocentric angle the downward-isotropic and fully-isotropic distributions of interacting particles produce line shapes that are indistinguishable. The line shape for a fan-beam distribution, expected for particles interacting at their mirror points (Murphy, Kozlovsky, & Ramaty 1988; Murphy et al. 1990), is almost indistinguishable from the downward-isotropic distribution.

We have fit the *RHESSI* count spectrum shown in Figure 1 using these and other angular distributions for α particles with power-law energy spectra having indices ranging from 3.5 to 4.5. We find no significant change in the fits for the different spectral indices. The fit over the full range in energy, from 150 keV to 8500 keV, is acceptable with $\chi^2/\text{dof} = 0.91$ for 767 degree of freedom. Share et al. (2003) list the best fitting parameters derived from this fit. Even though the bremsstrahlung continuum for this flare was very strong and dominated over the α - ^4He lines, the presence of these lines is required with 99.99% confidence. We base this conclusion on the increase of 16 in χ^2 when we force the flux in the α - ^4He lines to 0, with all other parameters free to vary. The positronium continuum, lying just below the 511-keV annihilation line, has a distinctly different shape from the α - ^4He fusion line complex and is readily separated. We have only been able to set an upper limit on this continuum for the flare (Share et al. 2003). We have multiplied our best-fitting line shape by a factor of five for clarity and plotted it separately in Figure 1. This shape was derived for an interacting angular distribution of α particles after they undergo saturated pitch-angle scattering in the corona (Hua et al. 1989; Murphy et al. 2003); this calculation has only been done for magnetic loops perpendicular to solar surface at the footpoints. We also plot the count spectrum derived by passing the fitted bremsstrahlung, annihilation, and nuclear line components (excluding the α - ^4He lines) through the instrument response function. This reveals the relative contribution of the α - ^4He line complex to the spectrum.

In Figure 3 we plot the *RHESSI* count spectrum from 370 to 500 keV, after removing all the fitted components except for the α - ^4He line complex; the solid curve is the calculated best-fitting line shape after passing it through the instrument response function. The spectrum is relatively structureless but exhibits evidence for a peak near 450 keV that is expected for α particles undergoing saturated scattering in the corona. The downward-isotropic and isotropic distributions produce similar shapes that are not significantly worse fits to the data; a fan-beam distribution would have a 15% chance of fitting the count spectrum as well as the others, based on $\Delta\chi^2$. In contrast, we find that a downward beam of α particles interact to produce a line shape with two relatively narrow peaks near ~ 418 and 467 keV (see Figure 2) that is a poor fit to the *RHESSI* count spectrum; there is only an $\sim 10^{-4}$ probability ($\Delta\chi^2 = 14.7$) that such a distribution would fit as well as the others.

3. DISCUSSION

RHESSI has detected the γ -ray line complex from ${}^7\text{Be}$ and ${}^7\text{Li}$ produced when flare-accelerated α particles interacted with ${}^4\text{He}$ in the chromosphere of the 2002 July 23 flare that occurred at a heliocentric angle of 73° . For a magnetic loop perpendicular to the solar surface at the footpoints, the calculated line shapes for most expected angular distributions of interacting α particles are similar, except for a downward beam which produces two distinct relatively narrow line features. We passed these line shapes through the instrument response function and found that only the downward beam is an unacceptable fit to the *RHESSI* count spectrum (0.01 % probability of fitting as well as the best-fitting shape based on the χ^2 statistic). The best-fitting line shape is produced by interacting α particles that undergo saturated pitch-angle scattering in the corona (Hua et al. 1989; Murphy et al. 2003). Fits to shapes produced by generic isotropic, downward-isotropic, and fan-beam distributions are also acceptable ($>15\%$ probability of fitting as well as the best fit). These results are consistent with earlier studies of the α - ${}^4\text{He}$ line made using *SMM* data (Share & Murphy 1997; Share et al. 2002).

Measurements of the red-shifts of ${}^{12}\text{C}$, ${}^{16}\text{O}$, and ${}^{20}\text{Ne}$ from 19 flares observed by *SMM*, grouped by heliocentric angle, placed additional constraints on the angular distributions of interacting particles. Isotropic and fan-beam angular distributions could be ruled out with confidence $>99\%$, while a downward beam was excluded at 99.5% confidence (Share et al. 2002). The best fits were obtained for interacting particles that suffered significant pitch-angle scattering in the corona or for a generic isotropic distribution of particles that primarily interact in the forward hemisphere. In all these cases it was assumed that the magnetic loops containing the accelerated particles were perpendicular to the solar surface at the footpoints.

There are two papers in this issue that also address the angular distribution of interacting ions in the 2003 July 23 flare (Smith et al. 2003; Murphy et al. 2003). Smith et al. (2003) have studied the line shifts and shapes of six de-excitation lines >800 keV and find that the lines are red-shifted from their laboratory values between 0.1 and 0.8 %. These shifts are significantly larger than that produced by the distribution favored by the *SMM* analysis, i.e. an isotropic particle distribution that primarily interacts in the forward hemisphere perpendicular to the solar surface. Smith et al. (2003) offer two suggestions that could explain these large redshifts: 1) the magnetic loop containing the particles might be inclined by $\sim 40^\circ$ toward Earth and 2) the interacting particles might be highly beamed normal to the solar surface. The first suggestion is consistent with our α - ${}^4\text{He}$ line measurements; we find that a forward isotropic distribution interacting in such a tilted loop would fit the data as well or better than our most favored distribution $\sim 15\%$ of the time based on a $\Delta\chi^2$

analysis. In contrast, as discussed above, a narrow beam is inconsistent with the *RHESSI* spectrum with high confidence (99.99%).

Murphy et al. (2003) studied the time history of the 2.223 MeV neutron-capture line to infer the angular distribution of interacting particles that produce the neutrons. They used the same physically-based model cited above, that includes pitch-angle scattering and transport of the interacting particles, to calculate the time history. Their calculations have only been performed for magnetic loops that are perpendicular to the solar surface at the foot-points. They find that the *RHESSI* data are fit best by an interacting particle distribution produced with a modest amount of pitch-angle scattering; they can not rule out saturated pitch-angle scattering that produces distributions of particles primarily interacting in the forward direction at a heliocentric angle of 73° . The best-fitting models to the neutron-capture line observations are also consistent with the α - ^4He line measurements ($>68\%$ confidence). Murphy et al. (2003) note that had they been able to perform their calculations for a magnetic loop tilted toward Earth, as suggested by the de-excitation line measurements (Smith et al. 2003), more forward-directed interacting particle distributions would have fit the data better.

In conclusion, we can conceive of only one scenario that is consistent with the high-energy de-excitation line, neutron capture line, and α - ^4He line measurements of the angular distribution of interacting particles for the 2003 July 23 flare. This requires a magnetic loop that is tilted toward Earth and a broad forward-directed particle distribution that impacts the solar atmosphere.

This work was supported NASA DPR W19746 and DPR W10049 at NRL. The work at the UCB and NASA Goddard was supported by NASA contract NAS 5-98033.

REFERENCES

- Hua, X. -M., Ramaty, R., & Lingenfelter, R. E. 1989, *ApJ*, 341, 516
- Kozlovsky, B. & Ramaty, R. 1974, *ApJ*, 191, L43
- Kozlovsky, B. & Ramaty, R. 1977, *Ap. Lett. Comm.*, 19, L19
- Kozlovsky B, Murphy R. J., Ramaty R. 2002, *ApJS*, 141, 523
- Lin, R. P., et al. 2002, *Solar Physics*, 210, 3
- Lin, R. P., et al. 2003, *ApJ*, this issue

Murphy, R.J., Kozlovsky, B., & Ramaty, R. 1988, ApJ, 331, 1029

Murphy, R. J., Hua, X.-M., Kozlovsky, B., & Ramaty, R. 1990, ApJ, 351, 299

Murphy, R. J., et al. 2003, ApJ, this issue

Ramaty, R., Kozlovsky, B., & Lingenfelter, R. E. 1979, ApJS, 40, 487

Ramaty, R. & Crannell C. J. 1976, ApJ, 203, 766

Share, G. H. & Murphy, R. J. 1997, ApJ, 485, 409

Share, G. H., Murphy, R. J., Kiener, J., & de Séréville, N. 2002, ApJ, 573, 464

Share, G. H., et al. 2003, ApJ, this issue

Smith D. M., et al. 2002, Solar Physics, 210, 33

Smith, D. M., et al. 2003, ApJ, this issue




Fig. 1.— Background-corrected count spectrum accumulated from 00:27:20 to 00:43:20 UT. Solid curve is the best fit to the data. Dotted curve shows the *RHESSI* response to the best-fit α -⁴He line model for an α -particle distribution that undergoes saturated pitch-angle scattering in the corona, multiplied by a factor of 5 for clarity. Dashed curve shows the response to the fitted bremsstrahlung, annihilation, and nuclear- line components (excluding the α -⁴He lines).




Fig. 2.— Comparison of the calculated α - ^4He line shapes for a flare at a heliocentric angle of 73° . Solid curve, downward-isotropic distribution (similar to one for α particles undergoing saturated pitch angle scattering in the corona). Dotted curve, fan-beam distribution. Dashed curve, downward beam.




Fig. 3.— Count spectrum of the α - ^4He line measured by *RHESSI* derived by subtracting the best-fitting bremsstrahlung and other components found to contribute to the background-corrected spectrum. The curve shows the *RHESSI* response to the line shape derived for an α -particle distribution that undergoes saturated pitch-angle scattering in the corona for magnetic loops perpendicular to the solar surface at a heliocentric angle of 73° .