

UMIACS-TR 89-123
CS-TR 2364

DECEMBER 1989

TWO SIMPLE RESIDUAL BOUNDS
FOR THE EIGENVALUES OF
HERMITIAN MATRICES

G. W. STEWART*

ABSTRACT

Let A be Hermitian and let the orthonormal columns of X span an approximate invariant subspace of X . Then the residual $R = AX - XM$ ($M = X^H AX$) will be small. The theorems of this paper bound the distance of the spectrum of M from the spectrum of A in terms of appropriate norms of R .

*Department of Computer Science and Institute for Advanced Computer Studies, University of Maryland, College Park, MD 20742. This work was supported in part by the Air Force Office of Sponsored Research under Contract AFOSR-87-0188.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE DEC 1989		2. REPORT TYPE		3. DATES COVERED 00-00-1989 to 00-00-1989	
4. TITLE AND SUBTITLE Two Simple Residual Bounds for the Eigenvalues of Hermitian Matrices				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Maryland, College Park, Department of Computer Science, Institute for Advanced Computer Studies, College Park, MD, 20742				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Let A be Hermitian and let the orthonormal columns of X span an approximate invariant subspace of X. Then the residual $R = AX - XM$ $M = XhAX$ will be small. The theorems of this paper bound the distance of the spectrum of M from the spectrum of A in terms of appropriate norms of R.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

TWO SIMPLE RESIDUAL BOUNDS FOR THE EIGENVALUES OF HERMITIAN MATRICES

G. W. STEWART

Let A be a Hermitian matrix with eigenvalues $\lambda_1 \geq \cdots \geq \lambda_n$. If X is a matrix with orthonormal columns that spans an invariant subspace of A and

$$M = X^H A X, \tag{1}$$

then $AX - XM = 0$.

Now suppose that the columns of X span an *approximate* invariant subspace of A . Then the matrix

$$R = AX - XM$$

will be small, say in the spectral norm $\|\cdot\|$ defined by $\|R\| = \max_{\|x\|=1} \|Rx\|$, where $\|x\|$ is the Euclidean norm of x .¹ If the eigenvalues of M are $\mu_1 \geq \cdots \geq \mu_k$, then we should expect the μ_i to be near k of the λ_i . The problem treated in this note is to derive a bound in terms of the matrix R .

An important result, due to Kahan [3] (see also [6, p.219]) states that there are eigenvalues $\lambda_{j_1}, \dots, \lambda_{j_k}$ of A such that

$$|\mu_i - \lambda_{j_i}| \leq \|R\|, \quad i = 1, \dots, k. \tag{2}$$

If nothing further is known about the spectrum of A , this bound is generally satisfactory, although it can be improved somewhat [5]. However, it frequently happens (e.g., in the Lanczos algorithm or simultaneous iteration [6, Ch.13-14]) that we know that $n - k$ of the eigenvalues of A are well separated from the eigenvalues of M : specifically, if we know that

$$\begin{aligned} &\text{there is a number } \delta > 0 \text{ such that exactly} \\ &n - k \text{ of the eigenvalues of } A \text{ lie outside the} \\ &\text{interval } [\mu_k - \delta, \mu_1 + \delta], \end{aligned} \tag{3}$$

then the bound in (2) can be replaced by a bound of order $\|R\|^2$. Bounds of the kind have been given by Temple, Kato, and Lehman (see [6, Ch.10] and [1, §6.5]).

¹In fact, the choice (1) of M minimizes $\|R\|$, although we will not make use of this fact here.

Early bounds of this kind, dealt only with a single eigenvalue and eigenvector. Lehman's bounds are in some sense optimal, but are quite complicated.

The purpose of this note is to give two other bounds derived from bounds on the accuracy of the column space of X as an invariant subspace of A . They are very simple to state and yet are asymptotically sharp. In addition they can be established by appealing to results readily available in the literature.

Theorem 1. *With the above definitions, assume that A and M satisfy (3). If*

$$\rho \equiv \frac{\|R\|}{\delta} < 1,$$

then there is an index j such that $\lambda_j, \dots, \lambda_{j+k-1} \in (\mu_k - \delta, \mu_1 + \delta)$ and

$$|\mu_i - \lambda_{j+i-1}| \leq \frac{1}{1 - \rho^2} \frac{\|R\|^2}{\delta}, \quad i = 1, \dots, k.$$

Proof. Let $(X \ Y)$ be unitary. Then

$$\begin{pmatrix} X^H \\ Y^H \end{pmatrix} A(X \ Y) = \begin{pmatrix} M & S^H \\ S & N \end{pmatrix}$$

where $\|S\| = \|R\|$. By the “sin Θ ” theorem of Davis and Kahan [2] there is a matrix P satisfying

$$\|P(I + P^H P)^{\frac{1}{2}}\| \leq \rho. \quad (4)$$

such that the columns of

$$\hat{X} = (X + YP)(I + P^H P)^{-\frac{1}{2}}$$

(which are orthonormal) span an invariant subspace of A . From (4) it follows that

$$\frac{\|P\|}{\sqrt{1 + \|P\|^2}} \leq \rho,$$

and since $\rho < 1$

$$\|P\| \leq \frac{\rho}{\sqrt{1 - \rho^2}}. \quad (5)$$

Let $\hat{Y} = (Y - XP^H)(I + PP^H)^{-\frac{1}{2}}$. Then $(\hat{X} \ \hat{Y})$ is unitary. Since the columns of $\hat{X}$ span an invariant subspace of A , we have $\hat{Y}^H A \hat{X} = 0$. Hence

$$\begin{pmatrix} \hat{X}^H \\ \hat{Y}^H \end{pmatrix} A(\hat{X} \ \hat{Y}) = \begin{pmatrix} \hat{M} & 0 \\ 0 & \hat{N} \end{pmatrix}.$$

In [7] it is shown that

$$\hat{M} = (I + P^H P)^{\frac{1}{2}}(M + S^H P)(I + P^H P)^{-\frac{1}{2}}.$$

The eigenvalues of $\hat{M}$ are eigenvalues of A . Since $\rho < 1$ it follows from (2), they lie in the interval $(\mu_k - \delta, \mu_1 + \delta)$, and hence are $\lambda_j, \dots, \lambda_{j+k-1}$ for some index j . By a result of Kahan [4] on non-Hermitian perturbations of Hermitian matrices,

$$|\mu_i - \lambda_{j+i-1}| \leq \|(I + P^H P)^{\frac{1}{2}}\| \|(I + P^H P)^{-\frac{1}{2}}\| \|S\| \|P\|, \quad i = 1, \dots, k.$$

The theorem now follows on noting that $\|(I + P^H P)^{-\frac{1}{2}}\| \leq 1$ and inserting the bound (5) for $\|P\|$. ■

Two remarks. First, the theorem extends to operators in Hilbert space, provided X (now itself an operator) has a finite dimensional domain. Second, the bound is asymptotically sharp, as may be seen by letting $X = (1 \ 0)^T$ and

$$A = \begin{pmatrix} 0 & \epsilon \\ \epsilon & 1 \end{pmatrix}$$

(the eigenvalues of A are asymptotic to ϵ^2 and $1 - \epsilon^2$).

The requirement (3) unfortunately does not allow the eigenvalues of M to be scattered through the spectrum of A . If we pass to the Frobenius norm defined by $\|X\|_F^2 = \text{trace}(X^H X)$, then we can obtain a Hoffman-Wielandt type residual bound. Specifically, if

$$\delta = \min\{|\lambda_i - \mu_j| : \lambda_i \in \lambda(A), \mu_j \in \lambda(M)\} > 0, \quad (6)$$

then a variant of the $\sin \Theta$ theorem shows that there is a matrix P satisfying

$$\|P(I + P^H P)^{\frac{1}{2}}\| \leq \|P(I + P^H P)^{\frac{1}{2}}\|_F \leq \frac{\|R\|_F}{\delta}$$

such that the columns of

$$\hat{X} = (X + YP)(I + P^H P)^{-\frac{1}{2}}$$

span an invariant subspace of A . By a variant of Kahan's theorem due to Sun [9, 8], the eigenvalues $\lambda_{j_1}, \dots, \lambda_{j_k}$ of $\hat{M}$ may be ordered so that

$$\sqrt{\sum_{i=1}^k (\mu_i - \lambda_{j_i})^2} \leq \|(I + P^H P)^{\frac{1}{2}}\| \|(I + P^H P)^{-\frac{1}{2}}\| \|S\|_F \|P\|.$$

Hence we have the following theorem.

Theorem 2. *With the above definitions, assume that A and M satisfy (6). If*

$$\rho_F \equiv \frac{\|R\|_F}{\delta} < 1,$$

then there are eigenvalues $\lambda_{j_1}, \dots, \lambda_{j_k}$ of A such that

$$\sqrt{\sum_{i=1}^k (\mu_i - \lambda_{j_i})^2} \leq \frac{1}{1 - \rho_F^2} \frac{\|R\|_F^2}{\delta}.$$

References

- [1] F. Chatelin (1983). *Spectral Approximation of Linear Operators*. Academic Press, New York.
- [2] C. Davis and W. M. Kahan (1970). “The Rotation of Eigenvectors by a Perturbation. III.” *SIAM Journal on Numerical Analysis*, **7**, 1–46.
- [3] W. Kahan (1967). “Inclusion Theorems for Clusters of Eigenvalues of Hermitian Matrices.” Technical report, Computer Science Department, University of Toronto.
- [4] W. Kahan (1975). “Spectra of Nearly Hermitian Matrices.” *Proceedings of the American Mathematical Society*, **48**, 11–17.
- [5] N. J. Lehmann (1963). “Optimale Eigenwerteinschiessungen.” *Numerische Mathematik*, **5**, 246–272.
- [6] B. N. Parlett (1980). *The Symmetric Eigenvalue Problem*. Prentice-Hall, Englewood Cliffs, New Jersey.
- [7] G. W. Stewart (1971). “Error Bounds for Approximate Invariant Subspaces of Closed Linear Operators.” *SIAM Journal on Numerical Analysis*, **8**, 796–808.
- [8] G. W. Stewart and Ji guang Sun (1990). *Matrix Perturbation Theory*. Academic Press, Boston. In production.
- [9] J.-G. Sun (1984). “On the Perturbation of the Eigenvalues of a Normal Matrix.” *Math. Numer. Sinica*, **6**, 334–336.