

Data Assimilation with Regional Lagrangian Models

A. F. Bennett

College of Oceanic & Atmospheric Sciences

Oregon State University

104 Ocean Admin Bldg

Corvallis, OR 97331-5503

phone: (541) 737-2849 fax: (541) 737-2064 email: bennett@oce.orst.edu

Award #: N000149910040

<http://www.oce.orst.edu/faculty/bennett.html>

LONG-TERM GOALS

The long term goal is regional modeling and data assimilation, allowing us to test models, and get good estimates of the state of the ocean. These 'inverse' estimates use the data to overcome errors in initial conditions, in boundary conditions, and in dynamical parameterizations.

OBJECTIVES

The objective is to assimilate float or drifter data from both coastal waters and the deep ocean, over a period of at least three months, into a stratified primitive equation model.

APPROACH

Regional primitive equation models are ill-posed, if the domain is Eulerian. However, they are well-posed if the domain is Lagrangian (Bennett and Chua, 1999). It is therefore possible to find efficient solutions of the inverse or data assimilation problem in Lagrangian domains, using the methods developed here in the last decade. My postdoctoral fellow Jodi Mead had committed the last 24 months to computing solutions of the inverse problem in a Lagrangian domain together with synthetic data. She will continue work on this project for at least the next 12 months.

WORK COMPLETED

Beta-plane tests of the viscous shallow water equations in Lagrangian coordinates have included integrations of uniformly rotating flow with open boundaries, and spatially periodic integrations on the time and space scales of North Atlantic mesoscale variability.

The results of the integrations are in close agreement (3%) with those of Eulerian integrations of an otherwise identical model, for 100 days or more.

For variational assimilation of real N. Atlantic float and hydrography data, the tangent linear form, and adjoint, of the viscous Lagrangian model has been derived in spherical polar coordinates. These operators are being implemented iteratively in a fully nonlinear assimilation scheme.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE SEP 2000		2. REPORT TYPE		3. DATES COVERED 00-00-2000 to 00-00-2000	
4. TITLE AND SUBTITLE Data Assimilation with Regional Lagrangian Models				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) College of Oceanic & Atmospheric Sciences,,Oregon State University,104 Ocean Admin Bldg,,Corvallis,OR,97331				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

With help from A Macdonald of WHOI, we have used N. Atlantic hydrographic data from "HYDROBASE" to compute Montgomery potentials on isopycnal surfaces. These will provide 'first guess' or 'background' boundary conditions and initial conditions for assimilations of real float tracks and depths.

A presentation of this work will be given at the meeting "Lagrangian Analysis and Predictability of Coastal and Ocean Dynamics", October 2-6, 2000 in Ischia, Italy.

A rational derivation of open boundary conditions has been made for diffusive hydrostatic and nonhydrostatic dynamics, as no derivation could be found in the published literature.

RESULTS

With a uniformly rotating flow, the viscous Lagrangian model accurately describes a moving domain for time periods of at least 160 days. This is a nested experiment, and the values at the open boundaries are interpolated from a known Eulerian field. Such interpolation anticipates the use of real hydrographic data provided on Eulerian grids.

The integrations with time and space scales of North Atlantic mesoscale variability yielded a highly nonlinear flow regime, with complex float trajectories. Real float data are of comparable complexity. Successful assimilation of simulated data of this type should lead to successful assimilation of the real data.

IMPACT/APPLICATIONS

There is little published material on the Lagrangian form of the equations of fluid dynamics, or suitable numerics, most likely because the Lagrangian viewpoint is uncommon. Semi-Lagrangian numerics are common for efficient time integrations in numerical weather prediction, but they involve interpolation to a fixed grid, which is not desirable for assimilation of float or drifter data. Semi-Lagrangian methods are awkward in this context, but may be computationally competitive. The Lagrangian viewpoint will give new insights into ocean circulation, both as a forward model and as a basis for assimilating data.

TRANSITIONS

Not applicable

RELATED PROJECTS

We are implementing semi-Lagrangian methods to determine if they conserve the well-posedness of the open boundaries. We are also exploring Lattice Boltzmann methods for well-posed regional modeling and variational assimilation in Eulerian domains.

REFERENCES

A. F. Bennett and B. S. Chua, 1999: Open boundary conditions for Lagrangian geophysical fluid dynamics, *Journal of Computational Physics*, 153, 418-436.

PUBLICATIONS

A.F.Bennett, 2000: Open boundary conditions for diffusive and nonhydrostatic numerical weather prediction models. *Monthly Weather Review* (in preparation).

J. L. Mead and A.F. Bennett, 1999: Towards regional assimilation of data: The Lagrangian form of the reduced gravity model and its inverse. *Journal of Marine Systems* (to appear).