

Award Number: W81XWH-08-1-0384

TITLE: The Contribution of Genotype to Heterotopic Ossification after Orthopaedic Trauma

PRINCIPAL INVESTIGATOR: Erika J. Mitchell, M.D.

CONTRACTING ORGANIZATION: Vanderbilt University Medical Center
Nashville, TN 37232

REPORT DATE: June 2012

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE June 2012		2. REPORT TYPE Final		3. DATES COVERED 15 May 2008 – 14 May 2012	
4. TITLE AND SUBTITLE The Contribution of Genotype to Heterotopic Ossification after Orthopaedic Trauma				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-08-1-0384	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Erika J. Mitchell, Patrick Norris, Judith Jenkins, John Morris E-Mail: erika.j.mitchell@vanderbilt.edu; erimitchell@lumc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Vanderbilt University Medical Center Nashville, TN 37232				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Purpose, scope, significant findings, progress Heterotopic ossification has been associated with numerous factors, however, there has been no conclusive evidence that a given environmental element is causative. We propose that certain individuals are genetically predisposed toward an altered sympathetic response to trauma which not only contributes to post-traumatic morbidity but also the formation of heterotopic bone. We have genotyped 2869 patients admitted to a Level I trauma center ICU for alleles that may be associated with bone healing, autonomic regulation and inflammation. Our preliminary results in a subset of 1313 patients have shown an association of a minor allele of the $\beta 2$ adrenergic receptor with HO formation. Furthermore, the patients who formed HO also were more likely to have had a prolonged ICU stay and days on a ventilator independent of a higher ISS score which was also associated with HO. Head injury, as defined by the AIS head score, however, was inversely correlated with HO. We have now completed the data collection on 244 SNPs in 2869 individuals and are awaiting final analysis of this data.					
15. SUBJECT TERMS Heterotopic Ossification, Single Nucleotide Polymorphisms, Genetics, Fracture Healing					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	9	19b. TELEPHONE NUMBER (include area code)

TABLE OF CONTENTS

	<u>Page</u>
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	2
Reportable Outcomes.....	3
Conclusion.....	3
References.....	4
Supporting Data.....	5
List of Personnel.....	6

INTRODUCTION

Heterotopic ossification (HO) is a complication of spinal cord injury and traumatic limb injuries in both civilian and military populations. The incidence is greater in the military population for unknown reasons. Factors that have been studied as potential causes of HO include head injury, ventilator days, traumatized soft tissue and injury severity have not been conclusive.¹⁻⁴ Our data has demonstrated that head injury, believed to be a major risk factor for HO formation, is inversely correlated with HO in our civilian trauma population. We suggest that the cause is not entirely environmental, rather, it is a genetic predisposition that in addition to injury leads to an excessive response to musculoskeletal injury resulting in HO. The greater systemic response to injury in a subset of patients that leads to conditions such as acute respiratory distress syndrome and increased systemic inflammatory response scores may be related to the excessive healing response to musculoskeletal injury that leads to HO.⁵⁻⁸ We have obtained serum samples from 2869 patients admitted to a Level I trauma center for genetic analysis. The samples have been genotyped for 244 SNP polymorphisms that may be associated with variations in the systemic human response to trauma. Of these patients, 431 had femur fractures with a minimum of 6 weeks of radiographic followup. These radiographs were examined for an excessive heterotopic fracture healing response.

BODY

Previously we had reported on 1313 patients, 125 had evidence of HO (11%). HO was inversely correlated with traumatic brain injury. Injury severity score, ventilator days, ICU days and total hospital days positively correlated with HO formation independent of one another (Aim 2.1, Table 1). In this same group of patients, the CC allele of a polymorphism coding for a $\beta 2$ adrenergic receptor was associated with HO formation (Aim 1.4, Table 2). This supports the possibility that a differential systemic response to trauma in some individuals is also an underlying cause of heterotopic bone formation.

Genotyping has now been completed on 2869 patients for 244 SNPs. Four hundred and thirty one of these patients had femur fractures and were admitted to the Trauma ICU at Vanderbilt University Medical Center. Out of the 431 patients with femur fractures, 247 met the inclusion criteria of minimum 6 week radiographic followup, Evidence of HO on plain radiograph was seen in 57 (23%) of these patients (FIGURE 1). In addition, data has been collected from the Trauma Registry of the American College of Surgeons (TRACS) to include environmental factors. Our original proposal was to genotype for 36 SNPs in a population of 1200 patients, 200 HO and 1000 no-HO (Aim 1.2). We have been able to increase the number of polymorphisms genotyped due to changes in the technology in our core labs. However, due to the changes in our research team, we chose to limit our radiographic search to femur fractures only. While this decreases the number of patients in the sample size, it also decreases variability from fracture types and treatments which may limit confounding factors due to anatomy, differential surgical techniques and degrees of soft tissue injury. Data collection is now complete but we are still awaiting the analysis.

The changes in our research team have led to the loss of the member who could perform the multivariate analysis. Nonetheless, we are continuing to pursue this albeit much delayed and outside of the research award period. We are attempting to identify an individual and funding to take this to completion for the data analysis as we feel that there is merit to our hypothesis and the data has been collected. We do expect to retain this individual and we will send a supplemental report with our final results including any publications that result from the data as soon as this is achieved.

KEY RESEARCH ACCOMPLISHMENTS

1. Data has been compiled on 2869 individuals including:
 1. AIS Head
 2. ISS
 3. Ventilator days
 4. ICU days
 5. Hospital days
 6. Gender
 7. Age
 8. Radiographic results

2. Genotyping is complete on this population. The 244 SNPs code for the following 32 genes:
 1. Leptin
 2. Leptin receptor
 3. C-reactive protein
 4. Complement Factor H
 5. WNT 3A
 6. WNT 10B
 7. IL1A
 8. IL1B
 9. IL6
 10. ACVR1
 11. GSK3 α
 12. GSK3 β
 13. Bone sialoprotein
 14. Secreted phosphoprotein 1
 15. β 2 Adrenergic receptor
 16. β 3 Adrenergic receptor
 17. α 1 Adrenergic receptor 1A
 18. α 1 Adrenergic receptor 2A
 19. α 1 Adrenergic receptor 1B
 20. Tumor necrosis factor
 21. Vascular endothelial growth factor A
 22. Toll like receptor 4
 23. Exostosin 2
 24. LRP5
 25. LRP6
 26. Vitamin D receptor
 27. BMP2
 28. BMP4
 29. Sclerostin
 30. Frizzled 9
 31. FOXO1
 32. FOXO3

3. Radiographs for these individuals have been examined for HO around fractures of the femur
 1. 57 determined to have HO out of 247 (23%) consistent with civilian data

REPORTABLE OUTCOMES

Publications

Mitchell, E.J., Canter, J., Norris, P., Jenkins, J., Morris, J.A. *The Genetics of Heterotopic Ossification: Insight into the Bone Remodeling Pathway*. J Orthop Trauma, 24(9):530-533, 2010.

Presentations

Mitchell, E.J., Morris, J.A., Norris, P., Canter, J., Jenkins, J., *Injury Severity But Not Head Injury Associated with Heterotopic Ossification in Patients Admitted to a Level 1 Trauma Center*. (Poster) Orthopaedic Trauma Assoc., Annual Meeting 2011.

Mitchell, E.J., Morris, J.A., Norris, P., Canter, J., Jenkins, J., *Genetic Predictors of Heterotopic Ossification: Insight into the Bone Remodeling Pathway*. (Invited Presentation) War Extremity Injuries Symposium, AAOS, January 2010

Mitchell, E.J., Morris, J.A., Norris, P., Canter, J., Jenkins, J., *Genetic Predictors of Heterotopic Ossification: Insight into the Bone Remodeling Pathway*. (Podium Presentation) Orthopaedic Trauma Assoc., Annual Meeting 2009.

CONCLUSION

We have completed data collection including genotyping, radiographic examination, and environmental factors on 2869 patients admitted to the Trauma ICU at Vanderbilt University Medical Center. Our preliminary results from a subset of this data have shown a relationship between a polymorphism of the $\beta 2$ Adrenergic Receptor and the development of HO. Our environmental data has shown that a greater degree of total injury severity but *not* head injury is associated with HO formation. We use the Abbreviated Injury Score (AIS) for head injury instead of the Glasgow Coma Scale (GCS) which is a more specific indicator of traumatic brain injury. The GCS on admission has inaccuracies due to intubation and medication of a patient in the field which obscures the ability to assess mental status.

The dataset that has now been completed is more thorough and expected to reveal more relationships. If the $\beta 2$ Adrenergic Receptor again shows a correlation, this would support our hypothesis that an altered systemic sympathetic response to trauma in certain individuals is a potential factor in the formation of HO. In our preliminary data, patients with longer ICU stays and ventilator days had an increased risk of HO independent of ISS. This also supports our hypothesis. Despite similar injury profiles and demographics, some patients recover without significant complication while others suffer from acute respiratory distress syndrome, systemic inflammatory response syndrome and/or sepsis. We assert that this is a phenomenon dictated by a genetically determined response to trauma that differs across individuals. A better understanding of the underlying causes of this response not only will allow us to better prevent HO but also potentially prevent other morbid events after severe traumatic injuries.

REFERENCES

1. Giannoudis, P. V. *et al.* Accelerated bone healing and excessive callus formation in patients with femoral fracture and head injury. *Injury* **37**, S18–S24 (2006).
2. Renfree, K. J. *et al.* Evaluation of serum osteoblast mitogenic activity in spinal cord and head injury patients with acute heterotopic ossification. *Spine* **19**, 740–746 (1994).
3. Kaplan, F., Glaser, D., Hebel, N. & Shore, E. Heterotopic ossification. *J Am Acad Orthop Surg* **12**, 116–125 (2004).
4. Pape, H.-C. *et al.* Trauma-induced inflammation and fracture healing. *J Orthop Trauma* **24**, 522–525 (2010).
5. Takeda, S. *et al.* Leptin regulates bone formation via the sympathetic nervous system. *Cell* **111**, 305–317 (2002).
6. Takeda, S. Central control of bone remodeling. *Biochem.Biophys.Res Commun.* **328**, 697–699 (2005).
7. Bonnet, N., Pierroz, D. D. & Ferrari, S. L. Adrenergic control of bone remodeling and its implications for the treatment of osteoporosis. *J Musculoskelet.Neuronal.Interact.* **8**, 94–104 (2008).
8. Commins, S. P. *et al.* Norepinephrine is required for leptin effects on gene expression in brown and white adipose tissue. *Endocrinology* **140**, 4772–4778 (1999).

SUPPORTING DATA




FIGURE 1: Femur fracture with heterotopic bone formation (arrows) as demonstrated by ‘flares’ of bone away from the site of injury and heterogenous density compared to typical callus formation at 12 weeks post-injury.

	No HO	HO	p-Value
Age	40.67 ± 17.89	42.69 ± 16.38	0.087
ISS	24.84 ± 12.08	28.35 ± 11.75	0.003
AIS Head	2.13 ± 1.84	1.70 ± 1.80	0.016
Hosp Days	12.81 ± 12.84	18.42 ± 17.28	<0.001
ICU Days	5.61 ± 7.32	8.14 ± 8.75	<0.001
Vent Days	4.66 ± 7.11	6.60 ± 6.91	<0.001

TABLE 1: Age did not show any relationship with HO. AIS Head score was inversely correlated with HO formation. ISS, Hospital Days, ICU Days, and days on a ventilator were associated with the development of HO.

Polymorphysm	p - value
ADBR2 GG	0.421
ADBR2 GC	0.153
ADBR2 CC	0.039
TLR4 CC	0.210
TLR4 CT	0.142
TLR4 TT	0.220
CFH CC	0.161
CFH CT	0.826
CFH TT	0.209

TABLE 2: One of the polymorphisms for the β 2 adrenergic receptor was positively associated with the development of HO.

LIST OF PERSONNEL

Erika J Mitchell, MD (PI)

Loyola University Medical Center, Maywood, IL
Dept of Orthopedic Surgery and Rehabilitation

John A Morris, MD (co-PI)

Vanderbilt University Medical Center, Nashville, TN
Department of General Surgery

Patrick Norris, PhD

Vanderbilt University Medical Center, Nashville, TN
Department of General Surgery

Jeffery Canter, PhD (2008-2010)

Vanderbilt University Medical Center, Nashville, TN
Department of General Surgery

Judy Jenkins, RN

Vanderbilt University Medical Center, Nashville, TN
Department of General Surgery

David Taber, MD (2010)

Vanderbilt University Medical Center, Nashville, TN
Department of Radiology