

THE CLOSEST M-DWARF QUADRUPLE SYSTEM TO THE SUN

CASSY L. DAVISON¹, R. J. WHITE¹, W.-C. JAO¹, T. J. HENRY¹, J. I. BAILEY III², S. N. QUINN¹,
J. R. CANTRELL¹, A. R. RIEDEL^{3,4}, J. P. SUBASAVAGE⁵, J. G. WINTERS¹, AND C. J. CROCKETT⁵

¹ Department of Physics and Astronomy, Georgia State University, Atlanta, GA 30303, USA

² Department of Astronomy, University of Michigan, Ann Arbor, MI 48109, USA

³ Department of Physics and Astronomy, Hunter College, New York, NY 10065, USA

⁴ American Museum of Natural History, New York, NY 10024, USA

⁵ United States Naval Observatory, Flagstaff, AZ 86002, USA

Received 2013 August 23; accepted 2013 October 22; published 2013 December 24

ABSTRACT

We report new infrared radial velocity measurements obtained with CSHELL at NASA's Infrared Telescope Facility that reveal the M3.5 dwarf GJ 867B to be a single-lined spectroscopic binary with a period of 1.795 ± 0.017 days. Its velocity semi-amplitude of 21.4 ± 0.5 km s⁻¹ corresponds to a minimum mass of $61 \pm 7 M_{\text{JUP}}$; the new companion, which we call GJ 867D, could be a brown dwarf. Stable astrometric measurements of GJ 867BD obtained with CTIO's 0.9 m telescope over the last decade exclude the presence of any massive planetary companions ($7\text{--}18 M_{\text{JUP}}$) with longer orbital periods (2–10 yr) for the majority of orientations. These complementary observations are also used to determine the trigonometric distance and proper motion of GJ 867BD; the measurements are consistent with the *HIPPARCOS* measured values of the M2 dwarf GJ 867AC, which is itself a 4.1 day double-lined spectroscopic binary at a projected separation of $24''.5$ (216 AU) from GJ 867BD. These new measurements strengthen the case that GJ 867AC and GJ 867BD are physically associated, making the GJ 867 system one of only four quadruple systems within 10 pc of the Sun ($d = 8.82 \pm 0.08$ pc) and the only among these with all M-dwarf (or cooler) components.

Key word: stars: low-mass

Online-only material: color figure

1. INTRODUCTION

Although the processes involved in single star formation have a wide variety of observational and theoretical constraints (e.g., McKee & Ostriker 2007), the complex physics involved in the formation of binary and especially multiple star systems is much less understood (e.g., Halbwachs et al. 2003; Tokovinin 2008; Duchêne & Kraus 2013). Prompt fragmentation of a collapsing prestellar core is favored as setting the initial conditions for multiple star systems (e.g., Tohline 2002), though how this depends upon the initial conditions of the cloud core (e.g., metallicity, mass, etc.) is not well understood. Once initial fragments form, subsequent fragmentation (possibly within a disk), accretion from the remnant cloud core, N -body dynamics, and possibly orbital migration have been proposed to play competing roles in setting the final properties of multiple star systems (e.g., Bate 2000; Bonnell 2001; Delgado-Donate et al. 2004; Sterzik & Tokovinin 2002). While the task of identifying the relative importance of these dynamical phenomenon remains daunting, useful constraints can nevertheless be established by first determining fundamental multiple star properties, such as the overall multiplicity fraction, the distribution of mass ratios, the distribution of separations, and the mass dependence of companion properties. As an example, one of the most well-established observational trends is that the fraction of stars in binary systems decreases toward lower stellar masses; studies of nearby stellar samples reveal that 26% of M-dwarfs have companions (Delfosse et al. 2004; also see Fischer & Marcy 1992; Henry & McCarthy 1992), compared to 46% for Sun-like stars (Raghavan et al. 2010; also see Duquennoy & Mayor 1991). A likely implication of this is that lower mass cloud

cores fragment less often, despite any subsequent dynamical evolution.

Formation events that lead to triple, quadruple, and even higher order stellar systems also appear to occur less often at low stellar masses. The emerging statistics are that 2.1% of M-dwarfs occur in triple systems, while 0.03% occur in quadruple or higher order systems, compared to 5.6% of G-dwarfs occurring in triple systems and 2.2% occurring in quadruple or higher order systems (Eggleton & Tokovinin 2008). While it is tempting to interpret this as a corresponding decrease in the frequency of multiple fragmentation events with lower cloud core mass, accurate assessments of high-order multiplicity are much more subject to observational biases and incompleteness, especially at low stellar masses. As an example of the incompleteness that exists, Law et al. (2010) find a surprisingly high fraction of triple and quadruple systems among a targeted sample of 36 extremely wide M-dwarf pairs, formerly known only as wide M-dwarf binary stars; the results mimic the discovery that brown dwarfs are more often found to be a close binary system when first identified as a wide companion to a star (Burgasser et al. 2005).

The most direct way to establish a more complete assessment of the multiplicity fraction over a broad range of masses is to target volume-limited samples of nearby stars. Not only are these the closest, brightest stars with the most precisely determined kinematics, but if trigonometric distances are known, Malmquist biases can be avoided. Nevertheless, observational biases remain even for the closest stars. Of the 348 stars known within 10 pc, 21 are G-dwarfs and 239 are M-dwarfs (Henry et al. 2006). But while two of these 21 G-dwarfs are the primary stars in quadruple star systems (no higher order G-dwarf systems are

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE DEC 2013		2. REPORT TYPE		3. DATES COVERED 00-00-2013 to 00-00-2013	
4. TITLE AND SUBTITLE The Closest M-Dwarf Quadruple System To The Sun				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Naval Observatory,,Flagstaff,,AZ,86002				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES The Astronomical Journal, 147:26 (9pp), 2014 February					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			
unclassified	unclassified	unclassified	Same as Report (SAR)	10	

Table 1
Quadruple or Higher Order Systems within 10 pc

Primary Star	Spectral Type of Primary ^a	Parallax (mas) ^b	Configuration ^c
GJ 423A	G0VJ ¹	119.51 ± 0.79 ^{1, 2, 3}	AC-BD
GJ 570A	K4V ²	170.62 ± 0.67 ^{1, 2, 3}	A-BC-D
GJ 695A	G5IVJ ³	120.32 ± 0.16 ^{2, 3}	AD-BC
GJ 867A	M2VJ ⁴	115.01 ± 1.30 ^{2, 3}	AC-BD
GJ 644A	M3.5VJ ⁴	154.96 ± 0.52 ^{1, 2, 3, 4}	ABC-D-GJ 643

Notes.

^a The letter “J” indicates a spectral type based on light from a spatially unresolved pair. References: (1) Batten et al. 1989; (2) Gray et al. 2006; (3) Gliese & Jahreiß 1991; (4) This work.

^b Parallaxes are weighted parallaxes from the following: (1) Söderhjelm (1999); (2) van Altena et al. (1995); (3) van Leeuwen (2007); (4) Costa et al. (2005); (5) Martin et al. (1998); (6) Martin & Mignard (1998).

^c Widely separated pairs are indicated with a hyphen.

known), only one of the 239 M-dwarfs is the primary of a quintuple star system; no M-dwarf quadruple star systems are known within 10 pc (see Table 1).⁶

The deficiency of known higher order pairs, in part, stems from a lack of high-dispersion spectroscopy that would reveal short-period companions. Early M-dwarfs are faint in the optical and chromospherically active making spectroscopic companion searches observationally difficult. It becomes even more challenging to search for short-period companions around mid to late M-dwarfs with spectroscopy, as these stars are even fainter and tend to have higher projected rotational velocities compared with earlier M-dwarfs (Jenkins et al. 2009; Reiners & Basri 2010; Reiners et al. 2012). These biases have inhibited good assessments of high-order multiples. For example, of the 90 M-dwarfs included in the spectroscopic companion search by Endl et al. (2006), only four stars have spectral types of M4 or later. The HARPS RV sample of 102 southern M-dwarfs only included stars with $V < 14$ mag (Bonfils et al. 2013). With the ratio of early M-dwarfs (M0–M4) to mid and late M-dwarfs (M4.5–M9) being 23/28 for stars within 5 pc (Cantrell et al. 2013), about half of all M-dwarfs are left out of high-dispersion spectroscopic surveys.

To address the question of M-dwarf multiplicity more completely, we initiated a volume-limited, astrometric and infrared spectroscopic search for companions with orbital periods between 3 days and 8 yr around mid M-dwarfs within 10 pc, which includes GJ 867B as a target (C. L. Davison et al., in preparation). A first result of this work is that GJ 867B is revealed to be a short-period spectroscopic binary (SB). We discuss the known characteristics of the GJ 867 system in Section 2. We list the astrometric results in Section 3 and the photometric results in Section 4, and newly assigned spectral types in Section 5. We discuss the spectroscopic results of GJ 867B and the new companion, GJ 867D, in Section 6. In Section 7, we summarize the results and discuss the case for physical association of this hierarchical system.

⁶ LHS1070 was claimed to be a quadruple system by Henry et al. (1999), but later determined to only be a triple system (Köhler et al. 2012). GJ 896A and GJ 896B were, on initial inspection, both claimed to be spectroscopic binaries (Delfosse et al. 1999) because both stars showed very large variations in radial velocity. These variations were later thought to be caused by inhomogeneous surface features combined with a large magnetic spot (X. Delfosse 2013, private communication).

2. THE GJ 867 SYSTEM

GJ 867A ($V_{KC}^7 = 9.09$ mag; Bessel 1990) is the primary component of a widely separated visual binary. It was classified spectroscopically by Vyssotsky & Mateer (1952) as an M-type star; we more accurately classify this star as M2.0V (see Section 5). This star is a member of young disk (Kiraga & Stepień 2007). Its visual companion, GJ 867B ($V_{KC} = 11.45$ mag; Bessel 1990), is separated by 24".5 at a position angle (P.A.) of 350°.45, based on Two Micron All Sky Survey position measurements. The spectral type of GJ 867B is M3.5V (see Section 5). Coordinates, magnitudes, and stellar properties of these stars are given in Table 2.

The two stars are presumed to be physically associated from their large (~ 0.45 yr⁻¹) and very similar proper motions (Dyer 1954; van Altena et al. 1995; Röser et al. 2011; Zacharias et al. 2013). Nevertheless, the stars have slightly but steadily changed their relative position since first recorded in 1877 with a separation of 21".4 and a P.A. of 359°, which presumably represents the orbital motion of the system (Herbig & Moorhead 1965).

GJ 867A was discovered by Herbig & Moorhead (1965) to be a double-lined SB with an orbital period of 4.083 days. The two components (A and C) have an estimated mass ratio (M_C/M_A) of 0.80 based on their relative velocity amplitudes. GJ 867B was identified with high-resolution optical spectroscopy as a candidate SB by Gizis et al. (2002), and later was listed as either having a high rotational velocity or being an SB by Bonfils et al. (2013). However, until now, its status as a short-period binary remained unconfirmed.

3. ASTROMETRY

3.1. Observations

Direct images of GJ 867BD⁸ were obtained at the 0.9 m telescope at CTIO from 2003 July until 2012 September, yielding a total of 16 astrometric epochs over 9.17 yr. Images were obtained using the 2048 × 2048 Tektronix CCD camera on the central quarter of the CCD chip. The CCD chip has a plate scale of 0".401 pixel⁻¹, which gives a field of view of 6".8 × 6".8 (Jao et al. 2003). Astrometric measurements of GJ 867BD were made in the V_J filter, which provided a field of five suitable reference stars. No astrometric measurements were taken of GJ 867AC, as this star saturates before sufficient signal-to-noise ratios (S/Ns) are obtained for the reference stars. Each night that GJ 867BD was observed, we obtained 5 to 10 images, for a total of 96 images on 16 different nights. Exposure times for an individual image varied from 25 to 100 s, with an average time of 54 s. We aimed to observe all images within 30 minutes of the star crossing the meridian to minimize the effects of differential color refraction (Jao et al. 2005). In each image, the FWHM of GJ 867BD is less than 2".4; this size is typically set by the seeing. The ellipticity of the target star and all reference stars are less than 20%. These criteria are adopted to eliminate elongated reference stars (e.g., binaries), and to avoid data with tracking/guiding errors.

During the time of observations, two different V_J filters were used. The first V_J filter possessed a small crack and was replaced in 2005 February with a nearly identical V_J filter,

⁷ Subscripts: KC indicates Kron–Cousins (SAAO system), and used later in the text, J indicates Johnson.

⁸ When the combined light of the individual components of the binary pair cannot be separated, we give the object the name of the binary pair.

Table 2
Characteristics of the GJ 867 System

Star	R.A. (J2000.0)	Decl. (J2000.0)	V_{KC} (mag)	R_{KC} (mag)	I_{KC} (mag)	J (mag)	H (mag)	K_s (mag)	Spectral Class	$v \sin i$ (km s ⁻¹)
GJ 867AC	22 38 45.60 ¹	-20 37 16.1 ¹	9.09 ²	8.08 ²	6.88 ²	5.67 ¹	5.11 ¹	4.80 ¹	M2.0VJ ³	4.7 ⁴
GJ 867BD	22 38 45.31 ¹	-20 36 51.9 ¹	11.47 ⁶	10.29 ⁶	8.77 ⁶	7.34 ¹	6.82 ¹	6.49 ¹	M3.5VJ ³	8.7 ³

References. (1) Cutri et al. 2003; (2) Bessel 1990; (3) This work; (4) Houdebine 2010.

which we confidently adopted knowing that it possessed the same transmission spectrum as the original. After a few years of obtaining data, we noticed a few milliarcsecond (mas) offset in the astrometric residuals of some stars that were known to be single from other techniques (Subasavage et al. 2009); we determined that the offset first appeared when we replaced the V_J filter. The effect of using the replacement V_J filter is more clearly seen in the astrometric residuals in the right ascension axis. After a close inspection of the first V_J filter, it was realized that the small crack does not project onto the part of the chip that is regularly used (i.e., the central quarter). Therefore, in 2009 July, we replaced the second V_J filter with the original V_J filter (Riedel et al. 2010). All data acquired after 2009 July were with the original V_J filter. Using data from both filters may increase the parallax error, but does not change the parallax value. We use all points in this analysis to maximize the parallactic ellipse coverage.

3.2. Astrometric Reductions

Here, we briefly summarize the reduction technique; further details can be found in Jao et al. (2005) and Subasavage et al. (2009). We obtain photometric parallaxes using the CCD distance relations of Henry et al. (2004) from which distances are calculated for all of our reference stars. We require reference stars to be further than 100 pc from the Sun, and initially assume the sums of all reference stars' proper and parallactic motions to be zero mas yr⁻¹. We measure the centroid of our target star against the presumed fixed grid of five reference stars using the SExtractor centroiding algorithm described in Bertin & Arnouts (1996). Relative to this reference grid, we obtain the proper motion, parallax and any residual astrometric motion of the target star using the Gaussfit program from Jefferys et al. (1987). We use the photometric parallaxes of the reference stars from our *VRI* photometry and correct for the parallactic motions of our reference stars to determine the absolute parallax of GJ 867BD.

3.3. Parallax and Proper Motion of GJ 867BD

We measure a relative parallax for GJ 867BD of 109.45 ± 1.74 mas. The correction for parallactic motion is 0.93 ± 0.20 , yielding a final absolute parallax of 110.38 ± 1.75 mas. This is the first parallax measurement for this star and corresponds to a trigonometric distance of 9.06 ± 0.15 pc. The errors on the parallax measurement represent our internal statistical precision. To estimate our systematic errors, we determine the standard deviation of the differences between the parallaxes of previous RECONS measurements (e.g., Subasavage et al. 2009; Riedel et al. 2010; Jao et al. 2011) and *HIPPARCOS*, which is 4.3 mas. This error dominates our internal error and we use it here to compare our values to previously published values. Two parallax values are available for GJ 867AC, 115.10 ± 7.40 mas (van Altena et al. 1995) and 115.01 ± 1.32 mas by *HIPPARCOS* (van Leeuwen 2007). Our parallax measurement of GJ 867BD is within 1.1σ of the weighted parallax measurement from van

Altena et al. (1995) and van Leeuwen (2007) for GJ 867AC. The similar distance measurements strengthen the case that GJ 867AC and GJ 867BD are physically associated. The weighted mean parallax of the GJ 867 system is 113.37 ± 1.04 mas, which corresponds to a distance of 8.82 ± 0.08 pc.

Our measurements of the proper motion amplitude, μ , and the proper motion P.A. for GJ 867BD are 421.8 ± 0.6 mas yr⁻¹ and $97^\circ.1 \pm 0^\circ.1$, respectively. The errors on the proper motion amplitude represent our internal statistical precision. To estimate our systematic errors, we again use the standard deviation of the differences between the proper motion amplitudes of previous RECONS measurements (e.g., Subasavage et al. 2009; Riedel et al. 2010; Jao et al. 2011) and *HIPPARCOS*, which is 16.8 mas yr⁻¹. Similar to our parallax errors, the systematic error for the proper motion amplitude dominates the error budget. Salim & Gould (2003) measure the μ of GJ 867BD to be 431.2 ± 10.2 mas yr⁻¹ and the proper motion P.A. to be $98^\circ.1$, which is within 0.6σ of our measurement of the μ of GJ 867BD; they do not provide an uncertainty on their measured position angle. GJ 867AC has a μ of 455.8 ± 2.7 mas yr⁻¹ and a position angle of $99^\circ.9$ (Salim & Gould 2003). The difference in proper motion amplitudes for the AC and BD components is ~ 35 mas yr⁻¹, which is slightly larger than our systematic error of 16.8 mas yr⁻¹ and nevertheless consistent with orbital motion. Table 3 summarizes previous parallax and proper motion measurements.

3.4. Detection Limits for Astrometric Companions

The precise position measurements of GJ 867BD spanning almost 10 yr allow us to set mass limits on long-period astrometric companions. This first requires an accurate assessment of the position uncertainties which we estimate empirically. This is done by characterizing the statistical uncertainty associated with each epoch's position measurement. To do this, we calculate the mean position error per night, which is the average standard deviation of position offsets, once parallactic and proper motions are removed. The mean position errors for all nights in R.A. and decl. are 2.31 and 4.00 mas, respectively. The average residual deviation from the position predicted by the proper motion and parallax fit is calculated as the standard deviation of the nightly offsets, and in R.A. and decl. are 1.37 and 3.83 mas, respectively. As we are measuring the predicted position with better precision than the average error, this may be an indicator that we are slightly overestimating the mean position errors. See Figure 1.

In order to set limits on the presence of companions orbiting GJ 867BD with orbital periods between 2 to 10 yr, we perform 100,000 Monte Carlo simulations with a range of orbital properties. Orbital periods are restricted to 2, 4, 6, 8, and 10 yr, which are representative of our observational sampling and baseline. We estimate the mass of the primary to be $0.29 \pm 0.06 M_\odot$ based on the mass luminosity relation of Henry & McCarthy (1993). Although we adopt this mass for all of our

Table 3
Parallax and Space Motions of the GJ 867 System

Group	Targets	Abs. π (mas)	μ (mas yr ⁻¹)	P.A. (deg)	V_{tan} (km s ⁻¹)	Reference
RECONS	GJ 867AC
RECONS	GJ 867BD	110.38 ± 1.75	421.8 ± 0.6	97.1 ± 0.1	18.1	This paper
HIP	GJ 867AC	115.01 ± 1.32	455.9 ± 1.7	100.0	18.8	van Leeuwen (2007)
HIP	GJ 867BD
NLTT	GJ 867AC	...	455.8 ± 2.7	99.9	...	Salim & Gould (2003)
NLTT	GJ 867BD	...	431.2 ± 10.2	98.1	...	Salim & Gould (2003)
YPC	GJ 867AC	115.10 ± 7.40	459	97	18.9	van Altena et al. (1995)
YPC	GJ 867BD	...	449	99	...	van Altena et al. (1995)
LEP	GJ 867AC	...	458.0	100	...	Lépine & Gaidos (2011)
LEP	GJ 867BD	Lépine & Gaidos (2011)
TYCHO-2	GJ 867AC	...	455.8 ± 2.7	99.9	...	Høg et al. (2000)
TYCHO-2	GJ 867BD	Høg et al. (2000)
UCAC4	GJ 867AC	...	457.6 ± 11.3	100.0	...	Zacharias et al. (2013)
UCAC4	GJ 867BD	...	432.6 ± 15.2	100.6	...	Zacharias et al. (2013)
PPMXL	GJ 867AC	...	454.9 ± 2.7	100.2	...	Röser et al. (2011)
PPMXL	GJ 867BD	...	419.1 ± 4.6	98.4	...	Röser et al. (2011)
Weighted mean	GJ 867	113.37 ± 1.04	426.2 ± 0.4	99.2	18.6	

Notes. Several numbers related to characterizing the motions of this system are given without errors. All parallax and proper motion values given with errors of GJ 867AC and GJ 867BD are included in the calculation for the weighted mean values of GJ 867. The weighted mean position angle and weighted mean tangential velocities values are pure averages of the available numbers, as none of these numbers have listed errors.




Figure 1. Astrometric residuals of GJ 867BD after solving for the proper motion and parallactic motion. The mean position errors in R.A. and decl. are 2.31 mas and 4.00 mas, respectively. These are slightly larger than the average residual deviations about zero, which are 1.37 mas and 3.83 mas for R.A. and decl., respectively. From these data we see no indication of long term variations and exclude the presence of most brown dwarf and massive planets with long orbital periods (2–10 yr).

calculations, we note that this mass is based on the brightness of both GJ 867B and GJ 867D, as we are unable to deconvolve the binary and determine magnitude and color estimates for each component. The companion star, GJ 867D, appears to contribute little light to the system (see Section 6.3), and thus is unlikely to severely bias this estimate. This mass determination is slightly higher than the mass determined by the spectral type of GJ 867B and the spectral type mass relations in Kraus & Hillenbrand (2007) of $\sim 0.25 M_{\odot}$, but consistent within the errors. We use the weighted distance measurement of the GJ 867 system reported in Section 3.3. We allow the inclination to vary between 0° and 90° and the eccentricity to vary between 0 and 0.9. We allow the longitude of periastron, ω , and the longitude of the ascending node, Ω , to vary from 0° to 180° . This simulation suggests that

90% of the time, we would have been able to detect, with 99.7% (or 3σ) confidence, a companion with a mass of at least $18 M_{\text{JUP}}$ in a 2 yr period, $11.5 M_{\text{JUP}}$ in a 4 yr period, $9 M_{\text{JUP}}$ in a 6 yr period, $7.5 M_{\text{JUP}}$ in a 8 yr period, and $7 M_{\text{JUP}}$ in a 10 yr period. The implication is that GJ 867BD is unlikely to have any brown dwarf or massive planetary companions with periods between 2 to 10 yr. The orbital periods correspond to separations of 1.1 to 3.1 AU ($0''.07$ to $0''.22$, if face on).

4. NEW SYSTEM PHOTOMETRY

4.1. Photometric Observations

We obtained photometric measurements of GJ 867BD and its reference stars on three different nights using an identical setup as that used for the astrometric measurements at the CTIO 0.9 m telescope. As our initial goal was to obtain photometry of GJ 867BD, we allowed GJ 867AC to saturate in our images, and therefore provide no new apparent magnitude measurements for GJ 867AC. We obtained V_J , R_{KC} , and I_{KC} photometry of GJ 867BD that were calibrated using at least 10 photometric standard stars per night from Landolt (1992) and Graham (1982). All images were obtained at an airmass of less than 2.

4.2. Photometric Reductions

Observations of the photometric standards are used to determine transformation equations and to create the extinction curves, enabling us to calculate absolute photometry for GJ 867BD. Specific details are given in Jao et al. (2005) and Winters et al. (2011). The apparent magnitudes through the V_J , R_{KC} , and I_{KC} filters are 11.47 mag, 10.29 mag, and 8.77 mag, respectively, with an error of 0.01 mag each. Our measurements are within 0.02 mag of the measured photometry by Bessel (1990).

Although astrometric observations do not allow absolute photometry, relative photometry can be extracted. Consequently, all images used to search for variability are in the V_J filter. We measure the standard deviation of the instrumental magnitude of GJ 867BD, (σ_{mag}), relative to reference stars for over 16 epochs spanning 9.17 yr to assess the amount of photometric variability. For GJ 867BD, we use the same set of reference stars in the astrometric reduction. The average σ_{mag} is 32.4 mmag, which is above the mean activity level of most field dwarfs (~ 13 mmag; Jao et al. 2011), but is not surprising given its classification as a flare star (Byrne 1978).

5. NEW SPECTRAL TYPES

New spectral types are determined using the RECONS classification system, which focuses on K- and M-dwarfs. The spectral region used for this covers 6000–9500 Å at a resolution of 8.6 Å; further details are given in Jao et al. (2008) and A. R. Riedel et al. (in preparation). Observations were made using the R-C Spectrograph on the CTIO 1.5 m with the Loral 1200 × 800 CCD camera, utilizing the #32 grating in first order with tilt 15°:1, and order blocking filter OG570. We remove the H α feature and the telluric features on the spectra using the sky transmission map from Hinkle et al. (2003). We then compare our stellar spectra to our library of standards spectra from Henry et al. (2002) and determine that the best match with the lowest standard deviation to be the spectral type of our star. Spectral types are accurate to 0.5 spectral subclasses. We determine the spectral types of GJ 867AC and GJ 867BD based on the combined light of the spatially unresolved pairs to be M2.0 and M3.5, respectively. These values are consistent to within 0.5 subclasses of previously adopted values (e.g., Reid et al. 2004).

6. HIGH-DISPERSION SPECTROSCOPY

6.1. Observations

Spectrographic observations were made at NASA’s Infrared Telescope Facility, a 3 m telescope, using the high-resolution infrared echelle spectrograph, CSHELL (Tokunaga et al. 1990; Greene et al. 1993). Spectra were centered at 2.298 μm (vacuum) and span approximately 50 Å. This wavelength region was chosen to obtain precise radial velocities (RVs) by measuring the positions of the rich photospheric ^{12}CO R lines (2.3 μm) relative to a wavelength solution set by telluric methane absorption features in the Earth’s atmosphere (e.g., Blake et al. 2010; Bailey et al. 2012). Observations were made using the 0.5 slit, yielding a measured resolving power of $\sim 46,000$. Ten measurements of GJ 867B were obtained spanning a temporal baseline of 100 days between 2012 August 22 and 2012 November 29. Each measurement consists of two spectra taken in quick succession of one another along the vertical direction of the slit, separated by 10". The two different positions are referred to as nod positions. The first three RV measurements were obtained during late 2012 August, over a 4 day period. At that time, we noticed a large RV variation of almost 18 km s $^{-1}$ and flagged this variable candidate as a high priority target for our upcoming runs to confirm it as an SB. Exposure times averaging ~ 500 s each were set to obtain predicted S/Ns of ~ 100 per exposure, and resulted in pair-combined S/Ns ranging from 70 to 230. Along with observations of GJ 867B, spectra of either the A7V star Altair or the A1V star Sirius were obtained to gain an initial estimate of the wavelength solution and the instrumental profile of CSHELL, since these spectra contain only telluric features.

Finally, 20 flats and darks with integration times of 10 or 20 s were taken each night.

6.2. Spectroscopic Reductions

A brief overview highlighting the aspects of our spectral extraction and RV fitting routine will be given here. For a full explanation of the data reduction technique, the reader is referred to Bailey et al. (2012) and C. L. Davison et al. (in preparation).

To prepare the two-dimensional images for spectral extraction, we first correct for flat fielding using dark-subtracted, median-combined flat field images. Then, we remove sky emission, detector bias, and dark current from the spectral images by subtracting each individual image from its pair image taken at a different nod position that immediately preceded or followed the image. We assume that the changes in the detector and spectrograph properties are negligible over the short period in which the two spectra were taken at the different nod positions. Then, we perform an optimal extraction of the spectra as implemented for nod subtracted pairs (Horne 1986; Bailey et al. 2012; Tanner et al. 2012). Unlike a standard extraction that simply sums over all the pixels to determine the one-dimensional spectral profile, an optimal extraction sums pixels weighted by a variance image of the spectrum’s spatial profile to determine the spectral profile. It is assumed that the spectral profile varies smoothly in the direction parallel to the dispersion. Using a weighted image to determine the spectral profile gives the advantage that it minimizes the noisy contribution of the profile wings and eliminates the majority of the features on the spectral profile caused by cosmic rays or hot pixels.

6.3. Spectroscopic Analysis and Modeling as a Single-lined Spectroscopic Binary

If GJ 867BD is a double-lined SB, we would expect it to appear double-lined in at least some of our spectra. Visual inspection showed no sign of a secondary component. Likewise, if its companion is of comparable brightness, we might observe broader absorption lines when the stars in the system are at opposition. Broader absorption lines would then be translated by our analysis code into larger $v \sin i$ values. However, all 10 $v \sin i$ measurements are consistent to within 3.1 km s $^{-1}$ and show no apparent correlated variation with the changes in RV. These factors lead us to believe that this star is a single-lined spectroscopic binary (SB1). Also, there is a lack of obvious remaining residuals in the single star spectral modeling, described below.

We subsequently analyze the system as an SB1. After extracting the spectra, we fit each observation to high-resolution spectral models that are convolved to the resolution of CSHELL. The models are formed by combining synthetic spectra created from NextGen (Hauschildt et al. 1999) with empirical telluric models extracted from ultra high-resolution KPNO/FTS telluric spectra of the Sun (Livingston & Wallace 1991). Our synthetic model is composed of spectral features from a single M3.5V star. In order to make a realistic model, we assume the starting stellar parameters of GJ 867B as follows: We estimate the stellar temperature to be 3200 K based on its spectral type of M3.5V, using the relationship from Kraus & Hillenbrand (2007); we set its surface gravity ($\log(g)$) to be 4.8 (cgs), as this number is consistent with other field dwarfs of similar mass (Hillenbrand & White 2004). Our model spectrum consists of 19 parameters. The limb darkening coefficient parameter is set to 0.6, a reasonable assumption for cool stars at infrared wavelengths (Claret




Figure 2. Spectra of GJ 867BD. Spectra are modeled by combining a telluric spectrum (top spectrum), which provides an absolute wavelength reference, with a theoretical stellar spectrum (second spectrum). The CSHELL spectrum of GJ 867B is shown (black; third spectrum) in comparison with the best fit (red; third spectrum). The mean residuals of the best-fit spectrum are 1.45% (bottom spectrum). The average mean residuals for all CSHELL spectra obtained is 3.09%.

(A color version of this figure is available in the online journal.)

2000) and the remaining 18 parameters are left to vary. Of those variable parameters, 3 of those parameters are used to solve the wavelength solution and 9 of those parameters are Gaussians used to model the point-spread function of the spectrum. The remaining 6 are the depth of the telluric spectra, depth of the synthetic spectra, $v \sin i$, RV, and 2 continuum normalization constants.

To determine the 12 parameters related to the wavelength and instrumental profile, we first fit the empirical telluric spectrum to A star observations. We obtain the optimal parameters by minimizing the variance weighted reduced χ^2 . On the one night when an A star was not observed, we averaged parameters determined on nights with A star observations and during the same run together to estimate initial guesses for the wavelength solution and instrumental profile. This step is critical, as our optimization routine is sensitive to initial guesses.

After obtaining these initial parameters, we fit the combined model of the telluric and synthetic spectra to our observed CSHELL spectra. During this iterative process, we first fit for the wavelength solution, the depth of the telluric and synthetic spectra, the RV, and the two normalization constants. Then, on our second iteration, we fit for $v \sin i$, the depth of the telluric and model spectral features, and the two normalization constants. On our final iteration, we optimize the wavelength solution, the depth of the telluric and synthetic spectra, the RV, and the two normalization constants. All of our fits are optimized using AMOEBA, which uses a downhill simplex method to find the best solution for multiple variables (Nelder & Mead 1965). We restrict AMOEBA to specified ranges of physically reasonable solutions. An example of an optimally extracted spectrum of GJ 867BD, the telluric model, the stellar model, and the best-fit combination are shown in Figure 2.

6.4. Spectroscopic Results

The determined RVs and projected rotational velocities ($v \sin i$ values) are listed in Table 4. The observed RV error is a combination of theoretical photon noise error, intrinsic stellar error, and instrumental error ($\sigma_{\text{obs}}^2 = \sigma_{\text{photon}}^2 + \sigma_{\text{stellar}}^2 + \sigma_{\text{instr}}^2$). The photon error is calculated based on the prescription by Butler et al. (1996). We assume the intrinsic stellar error to




Figure 3. RV measurements of GJ 867B vs. barycentric Julian date. A Keplerian curve is fit to the RV measurements, centered on the systemic velocity of -6.1 km s^{-1} . The errors on the RV measurements are smaller than the plot symbols.

Table 4
Spectroscopic Results for GJ 867BD

JD-240000	Orbital Phase	Radial Velocity (m s^{-1})	S/N	$v \sin i$ (km s^{-1})
56161.36	1.00	7594 ± 49	177	9.3
56163.41	1.25	-10132 ± 53	168	8.4
56164.45	0.49	7645 ± 61	135	9.8
56173.32	0.40	1396 ± 53	162	7.6
56173.41	0.49	7274 ± 62	138	8.9
56174.31	1.39	-19888 ± 46	204	9.3
56174.41	1.48	-24330 ± 44	228	8.4
56259.26	0.22	-14613 ± 47	127	9.5
56260.23	1.20	-3936 ± 41	197	9.5
56261.32	0.50	6415 ± 139	71	6.7

Note. The reported RV measurements have been corrected for the barycentric motion of Earth.

be zero. We know this is an underestimate of the stellar error for GJ 867B, which is an active flare star. However, we do not have enough data to accurately assess the error caused by intrinsic stellar error for this star. We adopt the average instrumental error of 68 m s^{-1} determined for 10 stars with low rotational velocities from our survey of mid M-dwarfs (C. L. Davison et al., in preparation) as the instrumental error for our RV measurements of GJ 867BD. The average $v \sin i$ value from our fitting prescription is $8.74 \pm 0.98 \text{ km s}^{-1}$. The error on the $v \sin i$ measurement is set from the standard deviation of the independent measurements from all 10 epochs.

6.5. Periodicity Analysis

6.5.1. The Orbital Period

GJ 867B shows a large change in RV on timescales of less than 1 day (Figure 3). To determine if the changes in RV are periodic, we use a Lomb–Scargle algorithm to search for periods between 0.5 and 10 days. The highest power of the periodogram favors a period of 1.795 days. Unfortunately, because of the limited sampling, there are many other peaks of comparable power (Figure 4). The highest peak at 1.795 days has a power value of 4.37, which is closely followed by power values of 4.24 and 4.21 for the next most likely peaks at periods of 1.541 and 2.841 days, respectively. To confirm that the highest peak is representative of the best-fit period, we conduct an independent χ^2 analysis, where we fit circular orbits to the top three periods given from the periodogram fit. Knowing the orbital period is less than 3 days, we assume the orbits are circularized




Figure 4. Periodograms of the RV measurements of GJ 867BD. The left panel illustrates the periodogram for the full range of periods considered, while the right panel is the portion containing the most significant period of 1.795 ± 0.017 days.




Figure 5. Phased RV curves of GJ 867BD. Shown are the RV curves of GJ 867BD phased to 1.541, 1.795, and 2.841 days, which are the three periods with the highest significance from the periodogram analysis. The black dotted line is the best-fit Keplerian model; the best-fit residuals are shown in the bottom panels. The period of 1.795 days is strongly favored because of its much smaller residuals.

due to tidal effects (Mayor & Mermilliod 1984). We optimize the circular fitted curves to give the lowest observed-minus-calculated ($O - C$) RV value using the AMOEBA minimization routine (Section 3.4); the results are shown in Figure 5. The average ($O - C$) value for 1.795 days is 0.58 km s^{-1} and is almost five times smaller than the ($O - C$) values for the periods of 2.841 days (2.88 km s^{-1}) and 1.541 days (3.00 km s^{-1}). Similarly, the χ^2 value for 1.795 days is 27 and 38 times smaller than the χ^2 values at 1.54 days and 2.841 days, respectively. From these comparisons, we conclude that 1.795 days is the orbital period of the BD system.

6.5.2. The Uncertainties in the Orbit

To determine the orbital properties and the associated uncertainties of the orbit, we use the nonlinear least squares fitting IDL routine, mpfitfun.pro (Markwardt 2009) to model our data as a sinusoidal curve. The amplitude of the fitted curve is $21.4 \pm 0.5 \text{ km s}^{-1}$, the velocity of the system is $-6.1 \pm 0.3 \text{ km s}^{-1}$, and the epoch of conjunction is JD $2,456,160.5611 \pm 0.0268$.

The peak with the most likely period is in a cluster of aliased peaks. To test whether the neighboring peaks represent reasonable orbital solutions, we again fit circular orbits to the top three aliasing peaks in closest proximity to the favorable period. The ($O - C$) values for these peaks range from 3.16 km s^{-1} to 5.56 km s^{-1} making these values significantly larger than the ($O - C$) values from the top three likely periods given by the periodogram fit. We conclude that neither

Table 5
Characteristics of the Spectroscopic Binaries

	GJ 867AC ^a	GJ 867BD
Period (days)	4.08322 ± 0.00004	1.795 ± 0.017
γ (km s^{-1})	-8.7	-6.1 ± 0.3
K_1 (km s^{-1})	46.8 ± 0.45	21.4 ± 0.5
K_2 (km s^{-1})	58.1 ± 0.46	...
T_0 (days)	JD 2437144 ± 0.006	JD 2456160.5611 ± 0.0268
M_1 (M_\odot)	$0.271 \pm 0.006/\sin^3 i$	0.29 ± 0.59^b
M_2 (M_\odot)	$0.218 \pm 0.005/\sin^3 i$	$\geq 0.056 \pm 0.007^b$
$a_1 \sin i$ (km)	$2.63 (\pm 0.03) \times 10^6$	$0.528 (\pm 0.02) \times 10^6$
$a_2 \sin i$ (km)	$3.26 (\pm 0.03) \times 10^6$...
e	0.010 ± 0.010	0^c
ω	$356^\circ \pm 32^\circ$	0^c

Notes.

^a Binary properties for GJ 867AC are from Herbig & Moorhead (1965).

^b The mass estimate of the primary is based on a mass luminosity relation while that of the secondary is a minimum mass ($m \sin i$; see the text).

^c The eccentricity and longitude of periastron are assumed to be zero.

of these periods represent good fits to our data and determine the uncertainty in the period from the FWHM of the central peak only, which yields a final best-fit period and uncertainty of 1.795 ± 0.017 days.

The average difference between the ($O - C$) value for our best-fit circular orbit of 0.58 km s^{-1} is significantly higher than our average estimated error for this star based on the theoretical photon noise error and the instrumental error. This could be caused by a few possibilities. GJ 867B could have a slightly elliptical orbit, and therefore the deviation we see from the observed compared to calculated value is real. Alternatively, our estimated errors could be low, as we did not account for any intrinsic stellar error. GJ 867B is a flare star with a nominally higher rotational velocity, which could be an indicator of a higher amount of intrinsic stellar jitter compared to other older field dwarfs. We caution that if the RV errors are too low, then the calculated errors on the orbital properties will be underestimated as well.

6.6. Limits on the Mass of GJ 867D

Without a direct detection of GJ 867D, we can only provide a lower mass limit based on the observed reflex motion of GJ 867B using Kepler's third law. We assume the orbit is circular and that the mass of the primary star is $0.29 \pm 0.06 M_\odot$ (Section 3.4). Under these assumptions the minimum mass of GJ 867D is $61 \pm 7 M_{\text{JUP}}$ ($0.059 \pm 0.007 M_\odot$); the uncertainty in the minimum mass is determined primarily by the uncertainty in the primary star's mass. GJ 867D could be a brown dwarf. The spectral properties of GJ 867AC and GJ 867BD are summarized in Table 5.

The brightness of GJ 867BD supports the conclusion that GJ 867D is very low mass. We compare the absolute magnitude in V to the ($V - K$) color of stars within 5 pc (Cantrell et al. 2013) to GJ 867 system (see Figure 6). GJ 867BD lies within the broad band making up the main sequence. Therefore, there is no evidence that the companion contributes much light, consistent with its undetected spectral features.

7. SUMMARY

We report infrared spectroscopic observations obtained with the CSHELL spectrograph that reveal GJ 867B to be an SB1 with an orbital period of 1.795 ± 0.017 days; the components are




Figure 6. Absolute V magnitude vs. $(V-K)$ colors. This color–absolute magnitude diagram shows GJ 867AC and GJ 867BD relative to all known M-dwarfs within 5 pc. This plot shows that GJ 867BD is not significantly over-luminous relative to this empirical main sequence, consistent with GJ 867BD being a single-lined SB and GJ 867D being low mass.

labeled B and D. Assuming that GJ 867B has a mass of $0.29 M_{\odot}$, then GJ 867D has a minimum mass ($m \sin i$) of $61 \pm 7 M_{\text{JUP}}$; it could be a brown dwarf. Astrometric measurements spanning nearly 10 yr are used to determine the trigonometric parallax and proper motion of GJ 867BD. The trigonometric parallax of GJ 867BD is consistent to within 1.1σ of the measured parallax of GJ 867AC by *HIPPARCOS*, confirming that the components of this 24/5 pair are at similar distances. The measured proper motion amplitude of GJ 867BD is $421.8 \pm 0.6 \text{ mas yr}^{-1}$, which is similar to but slightly smaller than the proper motion amplitude of GJ 867AC ($455.8 \pm 2.7 \text{ mas yr}^{-1}$; Salim & Gould 2003); both stars are moving in the same direction to within 2° . The difference ($-34.0 \text{ mas yr}^{-1}$) can easily be explained by orbital motion. Adopting the minimum masses in Table 5 and assuming the projected separation (216 AU) is the semi-major axis of the AC and BD pair, these components would be moving relative to one another at 1.8 km s^{-1} , corresponding to 43.2 mas yr^{-1} (at 8.82 pc) if in a face-on circular orbit. If in a somewhat inclined orbit (and if the components are more massive than the minimum masses used here), orbital motion could also explain the difference in the systemic velocities of GJ 867BD, at $-6.1 \pm 0.3 \text{ km s}^{-1}$ (Table 5), and GJ 867AC, at -8.7 km s^{-1} (Herbig & Moorhead 1965); we consider this apparent discrepancy less relevant, however, since the systemic velocity for GJ 867AC is based on much coarser precision individual measurements (1.4 km s^{-1}) and no formal uncertainty on the systemic velocity is provided. Altogether, the similar proximity, distance, and space motion of these two high proper motion SBs establish them as very likely physically associated. As such, the four star system GJ 867 becomes one of only four quadruple systems within 10 pc, and the closest quadruple system to the Sun with an M-dwarf primary.

The discovery of this nearby quadruple system suggests that many more hierarchical low-mass systems may exist nearby, but have not been found because of observational biases. As highlighted in the introduction, the discovery and characterization of systems like this potentially offer a powerful tool for constraining the uncertain physics involved in binary and multiple star formation. We note that nearby systems like GJ 867 are especially valuable, since these stars are close enough to have the spectroscopic orbits that can be spatially resolved interferometrically, allowing accurate masses of these systems to be determined.

The authors would like to thank Guillem Anglada-Escudé, Doug Gies, John Lurie, and Robert Parks for intellectual con-

versations and Travis Barman for generating our synthetic spectral models from NextGen (Hauschildt et al. 1999). The authors would like to note that the astrometric data were obtained by the RECONS team through the NOAO Surveys Program and the SMARTS consortium. The infrared spectroscopic data were obtained from the Infrared Telescope Facility, which is operated by the University of Hawaii under Cooperative Agreement no. NNX-08AE38A with the National Aeronautics and Space Administration, Science Mission Directorate, Planetary Astronomy Program. The authors are grateful for the telescope time granted to them. This work was supported by the National Science Foundation through a Graduate Research Fellowship to Cassy Davison. This project was funded in part by NSF/AAG grant no. 0908018 and the NSF grant no. AST05-07711 and AST09-08402.

REFERENCES

- Bailey, J. I., White, R. J., Blake, C. H., et al. 2012, *ApJ*, 749, 16
- Bate, M. R. 2000, *MNRAS*, 314, 33
- Batten, A. H., Fletcher, J. M., & MacCarthy, D. G. 1989, *PDAO*, 17, 1
- Bertin, E., & Arnouts, S. 1996, *A&AS*, 117, 393
- Bessel, M. S. 1990, *A&AS*, 83, 357
- Blake, C. H., Charbonneau, D., & White, R. J. 2010, *ApJ*, 723, 684
- Bonfils, X., Delfosse, X., Udry, S., et al. 2013, *A&A*, 549, A109
- Bonnell, I. A. 2001, in *IAU Symp.* 200, *The Formation of Binary Stars*, ed. H. Zinnecker & R. D. Mathieu (Cambridge: Cambridge Univ. Press), 23
- Burgasser, A. J., Kirkpatrick, J. D., & Lowrance, P. J. 2005, *AJ*, 129, 2849
- Butler, R. P., Marcy, G. W., Williams, E., et al. 1996, *PASP*, 108, 500
- Byrne, P. B. 1978, *IBVS*, 1407, 1
- Cantrell, J. R., Henry, T. J., & White, R. J. 2013, *AJ*, 146, 99
- Claret, A. 2000, *A&A*, 363, 1081
- Costa, E., Méndez, R. A., Jao, W.-C., et al. 2005, *AJ*, 130, 337
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *yCat*, 2246, 0
- Delfosse, X., Beuzit, J.-L., Marchal, L., et al. 2004, in *ASP Conf. Ser.* 318, *Spectroscopically and Spatially Resolving the Components of the Close Binary Stars*, ed. R. W. Hilditch, H. Hensberge, & K. Pavlovski (San Francisco, CA: ASP), 166
- Delfosse, X., Forveille, T., Beuzit, J.-L., et al. 1999, *A&A*, 344, 897
- Delgado-Donate, E. J., Clarke, C. J., Bate, M. R., & Hodgkin, S. T. 2004, *MNRAS*, 351, 617
- Duchêne, G., & Kraus, A. 2013, *ARA&A*, 51, 269
- Duquenois, A., & Mayor, M. 1991, *A&A*, 248, 485
- Dyer, E. R., Jr. 1954, *AJ*, 59, 218
- Eggleton, P. P., & Tokovinin, A. A. 2008, *MNRAS*, 389, 869
- Endl, M., Cochran, W. D., Kürster, M., et al. 2006, *ApJ*, 649, 436
- Fischer, D. A., & Marcy, G. W. 1992, *ApJ*, 396, 178
- Gizis, J. E., Reid, I. N., & Hawley, S. L. 2002, *AJ*, 123, 3356
- Gliese, W., & Jahreiß, H. 1991, *On: The Astronomical Data Center CD-ROM: Selected Astronomical Catalogs, Vol. 1*, ed. L. E. Brodzmann & S. E. Gesser (Greenbelt, MD: NASA)
- Graham, J. A. 1982, *PASP*, 94, 244
- Gray, R. O., Corbally, C. J., Garrison, R. F., et al. 2006, *AJ*, 132, 161
- Greene, T. P., Tokunaga, A. T., Toomey, D. W., & Carr, J. B. 1993, *Proc. SPIE*, 1946, 313
- Halbwachs, J. L., Mayor, M., Udry, S., & Arenou, F. 2003, *A&A*, 397, 159
- Hauschildt, P. H., Allard, F., & Baron, E. 1999, *ApJ*, 512, 377
- Henry, T. J., Franz, O. G., Wasserman, L. H., et al. 1999, *ApJ*, 512, 864
- Henry, T. J., Jao, W.-C., Subasavage, J. P., et al. 2006, *AJ*, 132, 2360
- Henry, T. J., & McCarthy, D. W., Jr. 1992, in *ASP Conf. Ser.* 32, *IAU Colloq. 135: Complementary Approaches to Double and Multiple Star Research*, ed. H. A. McAlister & W. I. Hartkopf (San Francisco, CA: ASP), 10
- Henry, T. J., & McCarthy, D. W., Jr. 1993, *AJ*, 106, 773
- Henry, T. J., Subasavage, J. P., Brown, M. A., et al. 2004, *AJ*, 128, 2460
- Henry, T. J., Walkowicz, L. M., Barto, T. C., & Golimowski, D. A. 2002, *AJ*, 123, 2002
- Herbig, G. H., & Moorhead, J. M. 1965, *ApJ*, 141, 649
- Hillenbrand, L. A., & White, R. J. 2004, *ApJ*, 604, 741
- Hinkle, K. H., Wallace, L., & Livingston, W. 2003, *BAAS*, 35, 1260
- Høg, E., Fabricius, C., Makarov, V. V., et al. 2000, *A&A*, 355, L27
- Horne, K. 1986, *PASP*, 98, 609
- Houdebine, E. R. 2010, *MNRAS*, 403, 2157

- Jao, W.-C., Henry, T. J., Beaulieu, T. D., & Subasavage, J. P. 2008, *AJ*, **136**, 840
- Jao, W.-C., Henry, T. J., Subasavage, J. P., et al. 2003, *AJ*, **125**, 332
- Jao, W.-C., Henry, T. J., Subasavage, J. P., et al. 2005, *AJ*, **129**, 1954
- Jao, W.-C., Henry, T. J., Subasavage, J. P., et al. 2011, *AJ*, **141**, 117
- Jefferys, W. H., Fitzpatrick, M. J., & McArthur, B. E. 1987, *CeMec*, **41**, 39
- Jenkins, J. S., Ramsey, L. W., Jones, H. R. A., et al. 2009, *ApJ*, **704**, 975
- Kiraga, M., & Stepien, K. 2007, *AcA*, **57**, 149
- Köhler, R., Ratzka, T., & Leinert, C. 2012, *A&A*, **541**, A29
- Kraus, A. L., & Hillenbrand, L. A. 2007, *AJ*, **134**, 2340
- Landolt, A. U. 1992, *AJ*, **104**, 340
- Law, N. M., Dhital, S., Kraus, A., Stassun, K. G., & West, A. A. 2010, *ApJ*, **720**, 1727
- Lépine, S., & Gaidos, E. 2011, *AJ*, **142**, 138
- Livingston, W., & Wallace, L. 1991, NSO Technical Report (Tucson, AZ: National Solar Observatory, National Optical Astronomy Observatory)
- Markwardt, C. B. 2009, in ASP Conf. Ser. 411, *Astronomical Data Analysis Software and Systems XVIII*, ed. D. A. Bohlender, D. Durand, & P. Dowler (San Francisco, CA: ASP), 251
- Martin, C., & Mignard, F. 1998, *A&A*, **330**, 585
- Martin, C., Mignard, F., Hartkopf, W. I., & McAlister, H. A. 1998, *A&AS*, **133**, 149
- Mayor, M., & Mermilliod, J. C. 1984, in IAU Symp. 105, *Observational Tests of the Stellar Evolution Theory*, ed. A. Maeder & A. Renzini (Dordrecht: Reidel), 411
- McKee, C. F., & Ostriker, E. C. 2007, *ARA&A*, **45**, 565
- Nelder, J. A., & Mead, R. 1965, *CompJ*, **7**, 308
- Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010, *ApJS*, **190**, 1
- Reid, I. N., Cruz, K. L., Allen, P., et al. 2004, *AJ*, **128**, 463
- Reiners, A., & Basri, G. 2010, *ApJ*, **710**, 924
- Reiners, A., Joshi, N., & Goldman, B. 2012, *AJ*, **143**, 93
- Riedel, A. R., Subasavage, J. P., Finch, C. T., et al. 2010, *AJ*, **140**, 897
- Röser, S., Schilbach, E., Piskunov, A. E., Kharchenko, N. V., & Scholz, R.-D. 2011, *A&A*, **531**, A92
- Salim, S., & Gould, A. 2003, *ApJ*, **582**, 1011
- Söderhjelm, S. 1999, *A&A*, **341**, 121
- Sterzik, M. F., & Tokovinin, A. A. 2002, *A&A*, **384**, 1030
- Subasavage, J. P., Jao, W.-C., Henry, T. J., et al. 2009, *AJ*, **137**, 4547
- Tanner, A., White, R., Bailey, J., et al. 2012, *ApJS*, **203**, 10
- Tohline, J. E. 2002, *ARA&A*, **40**, 349
- Tokovinin, A. 2008, *MNRAS*, **389**, 925
- Tokunaga, A. T., Toomey, D. W., Carr, J., Hall, D. N. B., & Epps, H. W. 1990, *Proc. SPIE*, **1235**, 131
- van Altena, W. F., Lee, J. T., & Hoffleit, E. D. 1995, *The General Catalogue of Trigonometric [Stellar] Parallaxes* (4th ed.; New Haven, CT: Yale Univ. Observatory)
- van Leeuwen, F. 2007, *A&A*, **474**, 653
- Vysotsky, A. N., & Mateer, B. A. 1952, *ApJ*, **116**, 117
- Winters, J. G., Henry, T. J., Jao, W.-C., et al. 2011, *AJ*, **141**, 21
- Zacharias, N., Finch, C. T., Girard, T. M., et al. 2013, *AJ*, **145**, 44