

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 NOV 2013		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE 'Bastion classification': evolution of experience mandates caution when considering using class as predictor for method of temporary vascular control				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Lundy J. B., Hobbs C. M.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

2. Kwiatkowski TC, Hanley Jr EN, Ramp WK. Cigarette smoking and its orthopedic consequences. *American Journal of Orthopedics (Belle Mead NJ)* 1996; 25:590–7.
3. Moghaddam A, Weiss S, Wolf CG, Schmeckenbecher K, Wentzensen A, Grutzner PA, et al. Cigarette smoking decreases TGF- β 1 serum concentrations after long bone fracture. *Injury*;41:1020–5.
4. Gustilo RB, Merkow RL, Templeman D. The management of open fractures. *Journal of Bone and Joint Surgery* 1990;72:299–304.

Arash Moghaddam-Alvandi*

Department of Orthopaedics, Trauma & Paraplegiologie,
University Hospital Heidelberg, Schlierbacher Landstraße 200a,
D-69118 Heidelberg, Germany

Gerald Zimmermann

Theresienkrankenhaus Mannheim, Department of Trauma Surgery,
Bassermannstr. 1, 68165 Mannheim, Germany

Kathrin Hammer

Klinik für Unfallchirurgie und Orthopädie, Berufsgenossenschaftliche
Unfallklinik Ludwigshafen, Ludwig – Guttman – Str. 13, 67071
Ludwigshafen, Germany

Thomas Bruckner

Institute for Medical Biometry and Informatics, University Hospital
Heidelberg, Im Neuenheimer Feld 305, 69120 Heidelberg, Germany

Paul Alfred Grützner¹

Jan von Recum²

Klinik für Unfallchirurgie und Orthopädie, Berufsgenossenschaftliche
Unfallklinik Ludwigshafen, Ludwig – Guttman – Str. 13, 67071
Ludwigshafen, Germany

*Corresponding author. Tel.: +49 06221 56 35394;
fax: +49 6221 56 29123

E-mail addresses: Arash.Moghaddam@med.uni-heidelberg.de
email@arash.de (A. Moghaddam-Alvandi)

g.zimmermann@theresienkrankenhaus.de (G. Zimmermann)
kath_hammer@yahoo.de (K. Hammer)

bruckner@imbi.uni-heidelberg.de (T. Bruckner)
Grutzner@bgu-ludwigshafen.de, recum@bgu-ludwigshafen.de
(P.A. Grützner)

¹Tel.: +49 0621 6810 2311.

²Tel.: +49 0621 6810 2013; fax: +49 0621 6810 2617.

<http://dx.doi.org/10.1016/j.injury.2012.08.017>

Letter to the Editor

'Bastion classification': Evolution of experience mandates caution when considering using class as predictor for method of temporary vascular control

We have read with great interest and applaud Jacobs and colleagues on their production and validation of a "Bastion score" to classify lower extremity injuries due to improvised explosive devices.¹ As the authors indicate, the ideal method of vascular control for class 3 injuries remains unclear. Jacobs and colleagues describe the use of iliac vascular control (either extra- or intraperitoneal) in a total of 51 of 177 limbs injured (46%). After review of the authors' findings, we conducted a retrospective analysis of lower extremity injuries cared for during the past 104 days during Op Herrick 15 and 16.

Table 1

Breakdown of Bastion class during study period (2 March–13 June 2012).

Bastion class	Number of extremities	Percentage of all extremities injured
1	0	0
2	8	7%
3	91	78%
4	12	10%
5	6	5%

Identical triage, resuscitative, and surgical strategies continue to be employed at Bastion hospital. During this time, we have cared for 67 patients with 117 injured limbs as a result of dismantled blast. A total of 41% ($n = 48$) of limbs injured resulted in traumatic amputation. Consistent with Jacobs report, class 3 injuries were the most common making up 78% of injured lower extremities (see Table 1). Proximal vascular control with temporary occlusion of iliac vessels via laparotomy was utilised in only 6 injured limbs (5% of all injured limbs; no patients underwent extraperitoneal iliac control). When considering Jacobs' manuscript, we noted that of 83 class 3 injured limbs without associated abdominal injuries, iliac control was utilised in 19 (23%). We found iliac control in this setting (class 3 injury without associated abdominal injury) necessary in only one patient (see Table 2). Whilst we agree that the classification system appears to be predictive of the initial musculoskeletal treatment, it was much less useful in predicting the need for proximal vascular control. As Table 2 indicates, the 6 limbs that had control of iliac vessels performed tended to be in patients with large injury burden (all had bilateral lower extremity injuries; all had associated injuries according to the ABCD component of the system). Our current approach for determination of method of obtaining vascular control is based on a combination of physiologic burden of injury, level of injury, need for intervention due to suspected abdominal injuries, and judgement and expertise of operating general and orthopaedic surgeons. This evolution in approach has contributed to the reduction in overall use of iliac control for lower limb injury. Other reasons contributing to this observed lower rate of this method of temporary vascular occlusion may exist. Injury mechanism has dictated widespread adoption of pelvic protective devices for ground troops. Pelvic protection has been shown in recent data analysis to contribute to a significant reduction in severe genitourinary injuries and likely has resulted in a decrease in perineal/pelvic haemorrhage (Oh JS, personal communication, 24 April 2012). In addition, similar group meetings and predeployment training have taken place to allow for discussion of techniques used and the associated risks and benefits of various methods of temporary proximal vascular control. These discussions have led our current group of deployed surgeons to determine that for thigh wounds amenable to proximal tourniquet application (i.e. class 3 injuries), separate incisions for exposure and occlusion of the iliac arteries are unnecessary in most cases.

We do not wish to recommend that there be a reluctance in obtaining vascular control proximal to the zone of injury in type 2 junctional haemorrhage or when the wounding pattern/physiologic burden of injuries make less aggressive methods of vascular control ineffective.^{2,3} We simply feel that as our skill and expertise has evolved, and as wounding patterns change with the evolution of blast and protective devices, broad statements mandating such approaches when more distal control is reasonable may be inappropriate and avoid unnecessary risks of laparotomy. We again congratulate the authors on such a monumental classification system which is long overdue and will contribute to easier communication with receiving hospitals along the echelon of

Table 2

Data for limbs managed with temporary occlusion of iliac vessels.

Patient	Vessel controlled	Approach to vessel	Bastion class of injured extremity	Traumatic amputation	Associated injuries ^a	Outcome of injured extremity	Outcome of contralateral extremity
1	Right EIA	Laparotomy	5S	N	A, B, C, D	AKA	TKA
2	Right CIA	Laparotomy	3	Y	A, B, D	TKA	BKA
3	Right CIA	Laparotomy	4	Y	B, C	AKA	AKA
3	Left CIA	Laparotomy	3	Y	B, C	AKA	AKA
4	Right CIA	Laparotomy	4	Y	A	AKA (high)	BKA
5	Right EIA	Laparotomy	3S	N	A, D	No amputation	No amputation

AKA = above knee amputation; BKA = below knee amputation; CIA = common iliac artery; EIA = external iliac artery; TKA = through knee amputation.

^a Bastion classification associated injuries: A = abdominal; B = genital/perineal; C = pelvic ring; D = upper limb.¹

medical evacuation and improved outcomes. A continuation of prospective observation of the observed rate of each class, surgical management, and outcomes based on the Bastion classification and our added modification of presence of traumatic amputation at presentation is currently planned.

Conflict of interest statement

Neither author has any conflicts of interest to disclose.

References

- Jacobs N, Rourke K, Rutherford J, Hicks A, Smith SRC, Templeton P, *et al*. Lower limb injuries caused by improvised explosive devices: proposed 'Bastion classification' and prospective validation. *Injury* 2012. [Epub ahead of print].

- Tai NR, Dickson EJ. Military junctional trauma. *Journal of the Royal Army Medical Corps* 2009;**155**:285–92.

- Parker P, Limb Trauma Working Group. Consensus statement on decision making in junctional trauma care. *Journal of the Royal Army Medical Corps* 2011;**157**(3 Suppl. 1):S293–5.

Jonathan B. Lundy*

Christopher M. Hobbs

Bastion Hospital, Op Herrick 16, Helmand Province,
Afghanistan

*Corresponding author. Tel.: +93 011 210 916 3301;

fax: +93 011 210 827 5397

E-mail address: Jonathan.lundy1@amedd.army.mil (J.B. Lundy)

<http://dx.doi.org/10.1016/j.injury.2012.08.016>