

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 13 FEB 2012		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Stem cells in large animal models of retinal and neurological disease				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Klassen H., Tucker B. A., Liew C. G., La Cour M., Wang H. C.,				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 2	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

Editorial

Stem Cells in Large Animal Models of Retinal and Neurological Disease

Henry Klassen,¹ Budd A. Tucker,² Chee G. Liew,³ Morten La Cour,⁴ and Heuy-Ching Wang⁵

¹ Gavin Herbert Eye Institute and Stem Cell Research Center, University of California-Irvine, Irvine, CA 92697, USA

² Department of Ophthalmology and Visual Science, University of Iowa, Iowa City, IA 52242, USA

³ Stem Cell Core, University of California-Riverside, Riverside, CA 92521, USA

⁴ Department of Ophthalmology, Glostrup Hospital, University of Copenhagen, 2600 Glostrup, Denmark

⁵ Ocular Trauma Division, U.S. Army Institute of Surgical Research, Fort Sam Houston, TX 78234, USA

Correspondence should be addressed to Henry Klassen, hklassen@uci.edu

Received 13 February 2012; Accepted 13 February 2012

Copyright © 2012 Henry Klassen et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The mammalian central nervous system (CNS) is notable in terms of complexity and sophistication, but also for a very limited regenerative ability in the face of injury and disease. While it is not uncommon to find remarkable regenerative capabilities in the CNS tissues of fish and amphibian species, similar findings are much more limited in reptiles and quite difficult to replicate in mammals. One method used to circumvent this inherent limitation has been the use of fetal tissue transplantation which has resulted in evidence of graft survival, integration and functional repair in a number of rat [1, 2] and also mouse models. Nevertheless, the use of fetal tissue grafts is generally impractical as a source of donor material and of relatively limited utility in the adult human eye [3]. Stem cell transplantation has more recently emerged as a potential treatment modality that offers even greater potential for tissue integration, while also being more scalable for widespread therapeutic use.

Much progress has been made in the development of stem cell transplantation in rodent models of retinal disease [4–6]; however, limitations to rodent-based research include nonhuman features such as small eyes, rod-dominated vision, the absence of a macula or visual streak, and short life span. As stem cell research moves toward clinical application in ophthalmology, the value of modeling and refining transplantation procedures in larger mammals becomes increasingly evident.

This special issue includes 10 papers that focus on stem and progenitor cells from the central nervous system (both brain and retina) of nonrodent mammals, or cells modified

to resemble such cells (e.g., iPS cells), together with the development of useful allogeneic donors and recipients. Although any species is of potential interest, here there is a particular emphasis on pig and cat following on from previous work conducted in these species.

The reports included in this special issue represent an interesting range of animals, including a nonhuman primate, as well as experimental approaches, including genetic modification and tissue engineering. Collectively, they serve to extend both the scope and horizon of stem cell research in nonrodent mammalian species. Despite the challenges of such work, it can be hoped that further efforts will be made to harness the potential of this area for translation of basic research findings into the clinical domain, not only as preclinical models for human treatment, but also as bone fide veterinary applications.

Henry Klassen
Budd A. Tucker
Chee G. Liew
Morten La Cour
Heuy-Ching Wang

References

- [1] S. B. Dunnett, A. Björklund, U. Stenevi, and S. D. Iversen, "CNS transplantation: structural and functional recovery from brain damage," *Progress in Brain Research*, vol. 55, pp. 431–443, 1982.
- [2] H. Klassen and R. Lund, "Retinal transplants can drive a pupillary reflex in host rat brains," *Proceedings of the National*

Academy of Sciences of the United States of America, vol. 84, no. 19, pp. 6958–6960, 1987.

- [3] N. D. Radtke, R. B. Aramant, H. M. Petry, P. T. Green, D. J. Pidwell, and M. J. Seiler, “Vision improvement in retinal degeneration patients by implantation of retina together with retinal pigment epithelium,” *American Journal of Ophthalmology*, vol. 146, no. 2, pp. 172–182, 2008.
- [4] M. Young, J. Ray, S. Whiteley, H. Klassen, and F. Gage, “Neuronal differentiation and morphological integration of hippocampal progenitor cells transplanted to the retina of immature and mature dystrophic rats,” *Molecular and Cellular Neurosciences*, vol. 16, no. 3, pp. 197–205, 2000.
- [5] H. Klassen, T.-F. Ng, Y. Kurimoto et al., “Multipotent retinal progenitors express developmental markers, differentiate into retinal neurons, and preserve light-mediated behavior,” *Investigative Ophthalmology and Visual Science*, vol. 45, no. 11, pp. 4167–4173, 2004.
- [6] H. Klassen, D. Sakaguchi, and M. Young, “Stem cells and retinal repair,” *Progress in Retinal and Eye Research*, vol. 23, no. 2, pp. 149–181, 2004.