



ARL-TN-0680 • JUNE 2015



Theoretical Prediction of the Heat of Formation, Density, and Relative Sensitivity for 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine

by Edward FC Byrd

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.



Theoretical Prediction of the Heat of Formation, Density, and Relative Sensitivity for 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine

by Edward FC Byrd

Weapons and Materials Research Directorate, ARL

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) June 2015		2. REPORT TYPE Technical Note		3. DATES COVERED (From - To) July 2014	
4. TITLE AND SUBTITLE Theoretical Prediction of the Heat of Formation, Density, and Relative Sensitivity for 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Edward FC Byrd				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Research Laboratory ATTN: RDRL-WML-B Aberdeen Proving Ground, MD 21005-5069				8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TN-0680	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Using a series of scripts developed by the US Army Research Laboratory, written to dramatically simplify the computation of crystalline density and heat of formation, the performance properties for the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine notional energetic material were evaluated. Additionally, a qualitative estimation of the impact sensitivity has been calculated. This report outlines the procedures used to generate this information, as well as Cheetah calculations using the predicted crystalline density and heat of formation.					
15. SUBJECT TERMS computational toolbox, script, crystalline density, heat of formation, impact sensitivity, synthesis, energetic materials, 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 20	19a. NAME OF RESPONSIBLE PERSON Edward FC Byrd
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) (410) 306-0729

Contents

List of Figures	iv
List of Tables	iv
Acknowledgments	v
1. Introduction	1
1.1 Neutral Heat of Formation	1
1.2 Crystalline Density	3
1.3 Impact Sensitivity	3
2. Method of Solution	4
3. Results and Discussion	4
4. Conclusions	7
5. References	8
List of Symbols, Abbreviations, and Acronyms	10
Distribution List	11

List of Figures

Fig. 1	Optimized structure of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule	5
Fig. 2	Electrostatic potential map of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule, with molecule overlay.....	6
Fig. 3	Electrostatic potential map of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule, without molecule overlay.....	6

List of Tables

Table	Cheetah-predicted properties of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule	7
-------	---	---

Acknowledgments

Dr Betsy Rice and Jennifer J Hare are acknowledged for their efforts in the original coding of the neutral heat of formation and EDAT tools, respectively. Dr James Ianni (Applications Engineer with Lockheed-Martin and contractor to the US Army Research Laboratory Department of Defense Supercomputing Resource Center [ARL DSRC]) is acknowledged for his gsubmit script, initially written for the ARL DSRC. Dr Rice is acknowledged for running the Cheetah calculations. All computations were performed at the ARL DSRC, Aberdeen Proving Ground, MD. Dr David Chavez (Los Alamos National Laboratory) is acknowledged as originator of the molecule modeled herein.

INTENTIONALLY LEFT BLANK.

1. Introduction

US Army Research Laboratory (ARL) researchers have achieved robust theoretical models capable of predicting performance properties, such as heats of formation^{1,2} and densities^{3,4} of energetic materials. This capability, coupled with a growing ARL synthetic experimental faculty, allows synthetic and formulation chemists to safely and quickly screen candidate materials to focus efforts only on the most promising compounds. This technical note will detail theoretical predictions of a notional energetic material.

The tools are based on correlations of quantum mechanically derived properties of isolated energetic molecules with heats of formation and crystalline densities, critical elements in predicting the detonation and propulsion properties, for both neutral and ionic solid energetic materials. As the crystal structure of a novel material is usually unknown, as it has yet to be synthesized, each of the correlations requires quantum mechanical data solely from single molecules and not from knowledge of the bulk material. To date, the neutral heat of formation tool can only treat molecules containing carbon, hydrogen, nitrogen, and oxygen. Additionally, a tool to qualitatively predict the impact sensitivity for neutral molecules was developed, also based solely on quantum mechanically derived data based on a single molecule. This report details the theoretically predicted performance and vulnerability properties for notional energetic molecules; as such, we will recap the methodologies used in the predictions in this technical report.

1.1 Neutral Heat of Formation

Developed in 2006,¹ the neutral heat of formation prediction tool is based on Hess's law (Eq. 1),⁵ where an atomic or group equivalency prediction for the gas phase heat of formation (Eq. 2) is added to a prediction for the heats of vaporization (Eq. 3) or sublimation (Eq. 4). Equations 2–4 use information determined from quantum mechanical calculations.

$$\Delta H^\circ_{\text{f (liquid/solid)}} = \Delta H^\circ_{\text{f (g)}} - \Delta H_{\text{vap/sub}}. \quad (1)$$

$$\Delta H^\circ_{\text{f (g)}} = E - \sum n_j \varepsilon_j. \quad (2)$$

$$\Delta H_{\text{vap}} = a(\text{SA})^{1/2} + b(\sigma^2_{\text{totV}})^{1/2} + c. \quad (3)$$

$$\Delta H_{\text{sub}} = a(\text{SA})^2 + b(\sigma^2_{\text{totV}})^{1/2} + c. \quad (4)$$

In Eq. 2, E is the quantum mechanically determined electronic energy, n_j is either the number of atom or group types j contained in the neutral molecule, and ε_j is

either the atom or group equivalent energy of atom/group j . These equivalent energies were determined through the fitting of experiment gas phase heats of formation $[\Delta H^\circ_{f(g)}]$ to quantum mechanically computed energies E .

Equations 3 and 4 use statistics derived from the electrostatic surface potential (ESP), as demonstrated by Politzer et al.⁶⁻⁸, to correlate to heats of vaporization and sublimation. As defined in the Murray et al.⁹ and Rice and Hare¹⁰ papers, the definitions of these electrostatic properties are global properties across the 0.001 electron/bohr³ isosurface of electronic density, an isosurface calculated to contain 98%–99% of the electronic density. To compute these statistics, the electrostatic potential V is required and is defined as

$$V(\mathbf{r}) = \sum_i \frac{Z_i}{|\mathbf{R}_i - \mathbf{r}|} - \int \frac{\rho(\mathbf{r}')d\mathbf{r}'}{|\mathbf{r}' - \mathbf{r}|}, \quad (5)$$

where Z_i and $\mathbf{R}_i$ represent the charge and position of atom i , and $\rho(\mathbf{r})$ is the electronic density. The statistical tools developed by Murray et al.^{9,11} used in Eqs. 3 and 4 are the surface area (SA), the variance of electrostatic surface potential (σ^2), and the balance parameter (v). These are defined as

$$\Sigma^x = \frac{1}{n} \sum_{i=1}^n V^x(r_i), \quad (6)$$

$$\sigma_x^2 = \frac{1}{n} \sum_{i=1}^n |V^x(r_i) - \Sigma^x|^2, \quad (7)$$

$$\sigma_{tot}^2 = \sigma_+^2 + \sigma_-^2, \quad (8)$$

and

$$v = \frac{\sigma_+^2 \sigma_-^2}{[\sigma_{tot}^2]^2}, \quad (9)$$

where $V(r_i)$ is the value of the ESP at r_i and Σ is the average electrostatic surface potential (x is either the positive $[+]$ or negative $[-]$ portion of the ESP). SA is the surface area of the 0.001 electron/bohr³ isosurface of the electron density of the molecule, σ_{tot}^2 is a measure of the variability of electronic potential on the surface, and v is the degree of balance between the positive and negative charges on the isosurface. Equations 3 and 4 contain a different set of fitted a , b , and c parameters. This methodology enjoys considerable accuracy, with the root mean square errors for the solid phase heat of formation under 6 kcal/mol.

1.2 Crystalline Density

Initially developed in 2007³ and revised in 2013,⁴ this method correlates the quantum mechanically derived electronic density volume of the molecular unit to the crystalline density. This correlation exhibits significant success in predicting the crystal density, with errors under 3% for neutral materials. This correlation only requires the quantum mechanically derived molecular volume and the molecular mass for a single molecular unit, along with associated electrostatic properties (Eq. 10).

$$\rho_{\text{crystal}} = a(\mathbf{M} / \mathbf{V}_M) + b(v\sigma_{\text{tot}}^2) + c, \quad (10)$$

where a , b , and c are fitted terms, different from those in Eqs. 3 or 4, $\mathbf{M}$ is the total molecular mass, $\mathbf{V}_M$ is the total molecular volume, and σ_{tot}^2 and v are the same as in Eqs. 8 and 9.

1.3 Impact Sensitivity

Besides performance parameters, ARL researchers have also developed a visual qualitative indicator of impact sensitivity (David Chavez, Los Alamos National Laboratory, private communication, 2014; unreferenced) as a method to assess any potential safety concerns with yet to be synthesized novel materials. Rice and Hare¹² attempted to correlate Politzer-type statistical properties of the electronic density to impact sensitivity for a series of polynitroaromatic and benzofuroxan molecules. While no strong correlations were found with the statistical parameters, there was a qualitative visual correlation related to the degree of electron deficiency over covalent bonds contained within the “backbone” of the molecules. Molecules that were very sensitive exhibited large electron deficiencies over the main structure, while less-sensitive molecules had a more evenly distributed electron density. This method, while quite approximate and limited to neutral molecules (due to the inherent charge separation found in ionic materials), can still yield a relative means of estimating impact sensitivities and thus potential hazards of notional materials. The visualizer used in this study, titled EDAT, was originally coded by Dr Jennifer Hare and later revised by Dr Anthony Yau. This tool is also used to derive the Politzer statistical properties used in the heat of formation and density predictors. In this report we shall apply these methods toward the molecule 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine (David Chavez, Los Alamos National Laboratory, private communication, 2014; unreferenced) to predict the heat of formation and crystalline density of this material.

2. Method of Solution

The analysis was performed using scripts written at ARL to perform the following required computations¹²:

1. A driver script that calls the other scripts, as well as submits the required Gaussian09¹³ calculations and monitors the queues (master_densheat.s.)
2. A script to construct the Gaussian input files (gbuild.s).
3. A script to gather and compute the data needed to determine the heats of formation and/or crystalline densities and subsequently calculate said properties (arl_densheat.s).

Given a properly formatted input file, the scripts will analyze the .dat input file, design the proper Gaussian input files and submit them, monitor the supercomputer queues for job completion, postprocess the necessary files, and, finally, compute the desired information.

The scripts also perform a limited series of error checking, such as checking that structures are optimized (i.e., the structure is not a transition state structure but rather a minimum), that the Gaussian calculations ran successfully to completion, and that, for neutral calculations, the number of atom types listed correspond to the number of atoms in the Cartesian coordinates.

The initial, user-designed structure has its geometry optimized using the B3LYP spin-restricted Kohn-Sham density functional theory (KS-DFT)¹⁴ with the 6-31G* Pople Gaussian basis set,¹⁵ currently done using the Gaussian09 program package.¹³ For the computation of the electronic energy required in Eq. 2, the B3LYP density functional theory energy¹⁵ with the 6-311++G(2df,2p) Pople basis set¹⁶ is obtained on the optimized geometry. For the crystalline density, the initial structure is optimized using the B3LYP KS-DFT with the 6-31G** Pople basis set,¹⁶ with the associated electronic density used in obtaining the molecular volume and electronic statistical properties required in Eq. 10. The visualization of the electrostatic potential surface was performed with the EDAT tool.

3. Results and Discussion

The 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule (David Chavez, Los Alamos National Laboratory, private communication, 2014; unreferenced) (Fig. 1) was constructed and submitted with the ARL scripts.¹²




Fig. 1 Optimized structure of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule

The resultant crystal density was computed to be 1.856 g/cc. The gas and solid phase heats of formation from the group additivity methodology were 120.12 and 95.22 kcal/mol, respectively.

Using the ARL-developed EDAT tool, the electrostatic map on the 0.001 isosurface was generated and is illustrated in Figs. 2 and 3 (with and without the molecule overlaid). For this visualization, the scalar range of the ESP ranges from -0.05 to 0.075 . For this visualization methodology, regions of large positive charge (red) over the backbone of the structure tend to indicate increased sensitivity. For this molecule, this ESP distribution would indicate moderate impact sensitivity. Using this predicted data, we then performed Cheetah 7.0 calculations¹⁸ to predict the performance parameters (Table).




Fig. 2 Electrostatic potential map of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule, with molecule overlay




Fig. 3 Electrostatic potential map of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule, without molecule overlay

Table Cheetah-predicted properties of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule

Property	Value
Pressure	31.084 GPa
Shock velocity	8.679 km/s
Temperature	3,395.8 K
Total energy of detonation	1.284 TNT equivalence (per cc)
	1.145 TNT equivalence (per gram)

4. Conclusions

The ARL-developed software tools were used to predict the heat of formation and crystalline density of the 3,7-dinitro-[1,2,4]triazolo[5,1-c][1,2,4]triazin-4-amine molecule. Using this predicted data, we then ran Cheetah calculations to predict the performance of this novel material. Additionally, we predicted the qualitative impact sensitivity of this compound using electrostatic potential maps. This information has been transferred back to the requesting synthetic chemist, Dr David Chavez of Los Alamos National Laboratory.

5. References

1. Byrd EFC, Rice BM. Improved prediction of heats of formation of energetic materials using quantum mechanical calculations. *J Phys Chem A*. 2006;110(3):1005–1013; *ibid* (errata) 2009;113:5813.
2. Byrd EFC, Rice BM. A comparison of methods to predict solid phase heats of formation of molecular energetic salts. *J Phys Chem A*. 2009;113(1):345–352.
3. Rice BM, Hare JJ, Byrd EFC. Accurate predictions of crystal densities using quantum mechanical molecular volumes. *J Phys Chem A*. 2007;111(42):10874–10879.
4. Rice BM, Byrd EFC. Evaluation of electrostatic descriptors for predicting crystalline density. *J Comp Chem*. 2013;34(25):2146–2151.
5. Atkins PW. *Physical chemistry*. Oxford (England): Oxford University Press; 1982.
6. Politzer P, Murray JS, Brinck T, Lane P. In: Nelson JO, Karu AE, Wong RB, editors. *Immunoanalysis of agrochemicals*. ACS Symposium Series; 1994. Washington (DC): American Chemical Society. p. 586.
7. Murray JS, Politzer P. In: Politzer P, Murray JS, editors. *Quantitative treatment of solute/solvent interactions: theoretical and computational chemistry*, 1. Amsterdam (Netherlands): Elsevier Scientific; 1994. p. 243–289.
8. Politzer P, Murray JS. Relationships between lattice energies and surface electrostatic potentials and areas of anions. *J Phys Chem A*. 1998;102:1018.
9. Murray JS, Lane P, Politzer P. Effects of strongly electron-attracting components on molecular surface electrostatic potentials: application to predicting impact sensitivities of energetic molecules. *Mol Phys*. 1998;93(2):187–194.
10. Rice BM, Hare JJ. A quantum mechanical investigation of the relation between impact sensitivity and the charge distribution in energetic molecules. *J Phys Chem A*. 2002;106:1770.
11. Murray JS, Lane P, Politzer P. Relationships between impact sensitivities and molecular-surface electrostatic potentials of nitroaromatic and nitroheterocyclic molecules. *Mol Phys* 1995;85(1):1–8.
12. Rice BM, Hare JJ. A quantum mechanical investigation of the relation between impact sensitivity and the charge distribution in energetic molecules. *J Phys Chem A*. 2002;106(9):1770–1783.

13. Byrd EFC. Simple computation of the heat of formation and density from theoretically predicted values. Aberdeen Proving Ground (MD): Army Research Laboratory (US); 2012 Sep. Report No.: ARL-TR-6206. Also available at: <http://www.arl.army.mil/arlreports/2012/ARL-TR-6206.pdf>.
14. Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Mennucci B, Petersson GA, et al. Gaussian 09, Rev. A.1. Wallingford (CT): Gaussian, Inc.; 2009.
15. Becke AD. Density-functional thermochemistry. III. The role of exact exchange. *J Chem Phys.* 1993;98:5648–5652; Lee C, Yang W, Parr RG. Development of the Colle-Salvetti correlation-energy formula into a functional of the electron density. *Phys Rev B.* 1988;37:785–789; Vosko SH, Wilk L, Nusair M. Accurate spin-dependent electron liquid correlation energies for local spin density calculations: a critical analysis. *Can J Phys.* 1980;58:1200–1211; Stephens PJ, Devlin FJ, Chabalowski CF, Frisch MJ. Ab Initio Calculation of Vibrational Absorption and Circular Dichroism Spectra Using Density Functional Force Fields. *J Phys Chem.* 1994;98:11623–11627.
16. Krishnan R, Binkley JS, Seeger R, Pople JA. Self-consistent molecular orbital methods. XX. A basis set for correlated wave functions. *J Chem Phys.* 1980;72:650; Frisch MJ, Pople JA, Binkley JS. Self-Consistent Molecular Orbital Methods 25: Supplementary Functions for Gaussian Basis Sets. *J Chem Phys.* 1984;80:3265; Clark T, Chandrasekhar J, Schleyer PVR. Efficient diffuse function-augmented basis sets for anion calculations. III. The 3-21+G basis set for first-row elements, Li–F. *J Comp Chem.* 1983;4:294.
17. Bastea S, Fried LE, Glaeseman KR, Howard WM, Kuo I-FW, Souers PC, Vitello PA. Cheetah 7.0 thermochemical code. Livermore (CA): Energetic Materials Center, Lawrence Livermore National Laboratory; 2012.

List of Symbols, Abbreviations, and Acronyms

6-31G*	Pople's Gaussian double-zeta polarized basis set with <i>d</i> polarization functions on each of the atoms Li through Ca
6-31G**	Pople's Gaussian double-zeta polarized basis set with <i>d</i> polarization functions on each of the atoms Li through Ca and <i>p</i> polarization functions on H and He
6-311++G(2df,2p)	Pople's Gaussian triple-zeta polarized basis set with 2 <i>d</i> and <i>f</i> polarization functions on each of the atoms Li through Ca and 2 <i>p</i> polarization functions on H and He along with <i>s</i> and <i>p</i> diffuse functions on each of the atoms Li through F and <i>s</i> diffuse functions on H and He
v	balance parameter
Σ	average electrostatic surface potential
σ^2	variance of electrostatic surface potential
ρ	electronic density
$\rho_{crystal}$	crystalline density
$\Delta H^\circ_f (g)$	gas phase heat of formation
$\Delta H^\circ_f (l)$	liquid phase heat of formation
$\Delta H^\circ_f (s)$	solid phase heat of formation
V	electrostatic potential
ARL	US Army Research Laboratory
B3LYP	Becke 3 parameter exchange with Lee-Yang-Parr correlation DFT functional
DFT	density functional theory
ESP	electrostatic surface potential
g/cc	grams per cubic centimeter
kcal/mol	kilocalories per mole (unit of energy)
KS-DFT	Kohn-Sham density functional theory

1 (PDF)	DEFENSE TECHNICAL INFORMATION CTR DTIC OCA	RDRL WMM J ZABINSKI RDRL WMP G R EHLERS
2 (PDF)	RDRL CIO LL IMAL HRA MAIL & RECORDS MGMT	
1 (PDF)	GOVT PRINTG OFC A MALHOTRA	
1 (PDF)	OFFICE OF NAVAL RSRCH C BEDFORD	
1 (PDF)	US ARMY ARDEC A DISTASIO S NICOLICH	
1 (PDF)	NAVAL SURF WARFARE CNTR H HAYDEN	
2 (PDF)	LOS ALAMOS NATL LAB DYN ENERG MATS DIV D CHAVEZ P LEONARD	
19 (PDF)	US ARMY RESEARCH LAB DIR USARL RDRL WM P BAKER B FORCH RDRL WML M ZOLTOSKI RDRL WML A W OBERLE RDRL WML B N TRIVEDI J MORRIS B RICE E BYRD RDRL WML C S AUBERT J SABATINI W SHERRILL J BANNING RDRL WML D R BEYER RDRL WML E P WEINACHT RDRL WML G W DRYSDALE RDRL WML H J NEWILL RDRL WML P D LYON	

INTENTIONALLY LEFT BLANK.