



---

**Hybridized Multiscale Discontinuous Galerkin Methods for Multiphysics**

**Jaime Peraire**  
**MASSACHUSETTS INSTITUTE OF TECHNOLOGY**

---

**09/14/2015**  
**Final Report**

DISTRIBUTION A: Distribution approved for public release.

Air Force Research Laboratory  
AF Office Of Scientific Research (AFOSR)/ RTA2  
Arlington, Virginia 22203  
Air Force Materiel Command

**REPORT DOCUMENTATION PAGE**

Form Approved  
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.  
**PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. REPORT DATE (DD-MM-YYYY)</b> 08/09/2015	<b>2. REPORT TYPE</b> Final Report	<b>3. DATES COVERED (From - To)</b> Jul 2012 – Jun 2015
--	---------------------------------------	--

<b>4. TITLE AND SUBTITLE</b> Hybridized Multiscale Discontinuous Galerkin Methods for Multiphysics	<b>5a. CONTRACT NUMBER</b>
	<b>5b. GRANT NUMBER</b> FA9550-12-1-0357
	<b>5c. PROGRAM ELEMENT NUMBER</b>

<b>6. AUTHOR(S)</b> Peraire, Jaime Nguyen, Ngoc C.	<b>5d. PROJECT NUMBER</b>
	<b>5e. TASK NUMBER</b>
	<b>5f. WORK UNIT NUMBER</b>

<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Massachusetts Institute of Technology 77 Massachusetts Avenue Cambridge, MA 02139	<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>
--	---

<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> Air Force Office of Scientific Research 875 N. Randolph St. Arlington, VA 22203	<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b> AF/OSR
	<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>

**12. DISTRIBUTION/AVAILABILITY STATEMENT**  
Public Release DISTRIBUTION A

**13. SUPPLEMENTARY NOTES**

**14. ABSTRACT**  
We have continued the development of the Hybridized Discontinuous Galerkin (HDG) method for the solution of systems of conservation laws. The focus has been on the development of robust, accurate, and efficient methods capable of handling complex physics in realistic geometries. In particular, we have: 1) proposed a Hybridized Multiscale Discontinuous Galerkin method (HMDG), 2) extended the HDG approximation to second-order elliptic eigenvalue problems, 3) presented a new phased-based method for high frequency wave propagation, 4) presented an embedded discontinuous Galerkin (EDG) method, 5) introduced a novel shock capturing technique, and 6) presented a novel approach for UQ

**15. SUBJECT TERMS**  
Discontinuous Galerkin, Hybridized Discontinuous Galerkin, Finite Elements, Fluid Mechanics, Uncertainty Quantification.

<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b>
<b>a. REPORT</b>	<b>b. ABSTRACT</b>	<b>c. THIS PAGE</b>			Jaime Peraire
U	U	U	U	5	<b>19b. TELEPHONE NUMBER (Include area code)</b> 617-253-1981

## INSTRUCTIONS FOR COMPLETING SF 298

**1. REPORT DATE.** Full publication date, including day, month, if available. Must cite at least the year and be Year 2000 compliant, e.g. 30-06-1998; xx-06-1998; xx-xx-1998.

**2. REPORT TYPE.** State the type of report, such as final, technical, interim, memorandum, master's thesis, progress, quarterly, research, special, group study, etc.

**3. DATE COVERED.** Indicate the time during which the work was performed and the report was written, e.g., Jun 1997 - Jun 1998; 1-10 Jun 1996; May - Nov 1998; Nov 1998.

**4. TITLE.** Enter title and subtitle with volume number and part number, if applicable. On classified documents, enter the title classification in parentheses.

**5a. CONTRACT NUMBER.** Enter all contract numbers as they appear in the report, e.g. F33315-86-C-5169.

**5b. GRANT NUMBER.** Enter all grant numbers as they appear in the report. e.g. AFOSR-82-1234.

**5c. PROGRAM ELEMENT NUMBER.** Enter all program element numbers as they appear in the report, e.g. 61101A.

**5e. TASK NUMBER.** Enter all task numbers as they appear in the report, e.g. 05; RF0330201; T4112.

**5f. WORK UNIT NUMBER.** Enter all work unit numbers as they appear in the report, e.g. 001; AFAPL30480105.

**6. AUTHOR(S).** Enter name(s) of person(s) responsible for writing the report, performing the research, or credited with the content of the report. The form of entry is the last name, first name, middle initial, and additional qualifiers separated by commas, e.g. Smith, Richard, J, Jr.

**7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES).** Self-explanatory.

**8. PERFORMING ORGANIZATION REPORT NUMBER.** Enter all unique alphanumeric report numbers assigned by the performing organization, e.g. BRL-1234; AFWL-TR-85-4017-Vol-21-PT-2.

**9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES).** Enter the name and address of the organization(s) financially responsible for and monitoring the work.

**10. SPONSOR/MONITOR'S ACRONYM(S).** Enter, if available, e.g. BRL, ARDEC, NADC.

**11. SPONSOR/MONITOR'S REPORT NUMBER(S).** Enter report number as assigned by the sponsoring/monitoring agency, if available, e.g. BRL-TR-829; -215.

**12. DISTRIBUTION/AVAILABILITY STATEMENT.** Use agency-mandated availability statements to indicate the public availability or distribution limitations of the report. If additional limitations/ restrictions or special markings are indicated, follow agency authorization procedures, e.g. RD/FRD, PROPIN, ITAR, etc. Include copyright information.

**13. SUPPLEMENTARY NOTES.** Enter information not included elsewhere such as: prepared in cooperation with; translation of; report supersedes; old edition number, etc.

**14. ABSTRACT.** A brief (approximately 200 words) factual summary of the most significant information.

**15. SUBJECT TERMS.** Key words or phrases identifying major concepts in the report.

**16. SECURITY CLASSIFICATION.** Enter security classification in accordance with security classification regulations, e.g. U, C, S, etc. If this form contains classified information, stamp classification level on the top and bottom of this page.

**17. LIMITATION OF ABSTRACT.** This block must be completed to assign a distribution limitation to the abstract. Enter UU (Unclassified Unlimited) or SAR (Same as Report). An entry in this block is necessary if the abstract is to be limited.

# HYBRIDIZED DISCONTINUOUS GALERKIN METHODS FOR MULTIPHYSICS

J. Peraire and C. N. Nguyen

Department of Aeronautics and Astronautics, Massachusetts Institute of Technology

## **Final Performance Report**

AFOSR Grant: FA9550-12-1-0357

Period covered: 7/1/2012-6/30/2015

**Title:** Hybridizable Discontinuous Galerkin (HDG) Methods for Systems of Conservation Laws

In recent years, discontinuous Galerkin (DG) methods have attracted considerable attention because they possess a number of desirable properties for solving hyperbolic systems of conservation laws. In particular, the DG methods work well on arbitrary meshes, result in stable high-order discretization of the convective and diffusive operators, allow for a simple and unambiguous imposition of boundary conditions, and are well-suited to adaptive strategies. However, the DG methods have been criticized for providing sub-optimally convergent approximations for the flux, as well as for producing a substantially larger amount of globally-coupled degrees of freedom (for the same mesh and polynomial degree of the approximation) in comparison to the well-established finite element methods for diffusion problems. In response to these criticisms, the hybridizable discontinuous Galerkin (HDG) methods were first introduced in [1] and later extended in [2], [3], [4], [5], [6], [7], [8], [9]. The essential ingredients are a local Galerkin projection of the underlying PDEs at the element level onto spaces of polynomials of degree  $k$  to parametrize the numerical solution in terms of the numerical trace; a judicious choice of the numerical flux to provide stability and consistency; and a global jump condition that enforces the continuity of the numerical flux to arrive at a global weak formulation in terms of the numerical trace. The HDG method is fully implicit, high-order accurate and endowed with several unique features which distinguish themselves from other discontinuous Galerkin methods. First, they reduce the globally coupled unknowns to the approximate trace of the solution on element boundaries, thereby leading to a significant reduction in the degrees of freedom. Second, they provide, for smooth viscous-dominated problems, approximations of all the variables which converge with the optimal order of  $k + 1$  in the  $L_2$ -norm. Third, they possess some superconvergence properties that allow us to define inexpensive element-by-element postprocessing procedures to compute a new approximate solution which may converge with higher order than the original solution. And fourth, they allow for a novel and systematic way for imposing boundary conditions for the total stress, viscous stress, vorticity, and pressure which are not naturally associated with the weak formulation of the methods. In addition, they possess other interesting properties for specific problems. Their approximate solution can be postprocessed to yield an exactly divergence-free and  $H(\text{div})$ -conforming velocity field for incompressible flows. They do not exhibit volumetric locking for nearly incompressible solids.

The first HDG method for the compressible Euler and Navier-Stokes equations was introduced in [10]. This method possesses a number of desirable properties. First, in implicit formulations, it reduces the globally coupled unknowns to the numerical trace of the solution on element boundaries, thereby leading to a significant reduction in the degrees of freedom. Second, it provides, for smooth (e.g., viscous-dominated) problems, approximations for all the variables which converge with the optimal order of  $k + 1$  in the  $L^2$ -norm. Third, it possesses superconvergence properties that allow us to compute a new approximate velocity which converges with order  $k + 2$  for  $k \geq 1$ . And fourth, it allows for a novel and systematic way

for imposing general (external or internal) boundary conditions.

In this grant, we have continued the development of Hybridized Discontinuous Galerkin (HDG) methods for the solution of systems of conservation laws. In particular, the focus has been the development of robust, accurate, and efficient methods capable of handling complex physics in realistic geometries. Below, we describe the main research results.

## I. RESEARCH RESULTS

### A. Multiscale HDG Method

We have introduced a hybridized multiscale discontinuous Galerkin (HMDG) method for the numerical solution of compressible flows [11]. The HMDG method is developed upon extending the hybridizable discontinuous Galerkin (HDG) by modifying the local approximation spaces on elements. The essential ingredients are (1) a local Galerkin projection of the underlying PDEs at the element level onto suitable local approximation spaces, (2) a judicious choice of the numerical flux to provide stability and consistency, and (3) a global jump condition that enforces the continuity of the numerical flux to arrive at a global system in terms of the numerical trace. The local approximation spaces are characterized by two integers  $(n^*, k^*)$ , where  $n^* \in [1, n]$  is the number of subcells within an element and  $k^* \in [0, k]$  is the polynomial degree of shape functions defined on the subcells. The selection of the value of  $(n^*, k^*)$  on a particular element depends on the smoothness of the solution on that element. More specifically, for elements on which the solution is smooth, we choose the smallest value  $n^* = 1$  and the highest degree  $k^* = k$ . For elements containing shocks in the solution, we use the largest value  $n^* = n$  and the lowest degree  $k^* = 0$ . The proposed method thus combines the accuracy and efficiency of high-order approximations with the robustness of low-order approximations. A key advantage of the HMDG method is that it can capture shocks without using artificial viscosity and limiting slopes/fluxes.

### B. Eigenvalue Problems

We have introduced hybridization and postprocessing techniques for the HDG approximation of second-order elliptic eigenvalue problems [12]. Hybridization reduces the HDG approximation to a condensed eigenproblem. The condensed eigenproblem is nonlinear, but smaller than the original mixed approximation. We derived iterative algorithms for solving the condensed eigenproblem and examined their interrelationships and convergence rates. For smooth eigenfunctions, the approximate eigenvalues and eigenfunctions converge at the rate  $2k + 1$  and  $k + 1$ , respectively. Here  $k$  is the degree of the polynomials used to approximate the solution, its flux, and the numerical traces. An element-by-element postprocessing technique to improve accuracy of computed eigenfunctions is also presented. We prove that a projection of the error in the eigenspace approximation superconverges and that the postprocessed eigenfunction approximations converge faster for smooth eigenfunctions. Our numerical studies showed that a Rayleigh quotient-like formula applied to the locally postprocessed approximations can yield eigenvalues that converge faster at the rate  $2k + 2$ .

### C. Phased-based Methods for Wave Propagation

For wave propagation [13], we have presented a new class of HDG methods capable of generating high-quality approximations of the Helmholtz equation for a very wide range of wave frequencies. Our approach combines the hybridizable discontinuous Galerkin methodology with the geometrical optics in an elegant fashion that allows us to take advantage of the strengths of these two methodologies. First, we enrich the local approximation spaces of the hybridizable discontinuous Galerkin methods with precomputed phases which are solutions of the eikonal equation in geometrical optics. Second, we propose a systematic procedure for computing multiple solutions of the eikonal equation. Third, we utilize the eigenvalue decomposition to remove redundant modes and obtain orthogonal basis functions. And fourth, these locally

orthogonal approximation spaces are used to define the global approximation spaces of the phase-based HDG methods. This class of methods possesses a number of critical advantages over the current state-of-the-art low-frequency and high-frequency schemes. Unlike other numerical methods which require the degrees of freedom to increase drastically with the wave number, the proposed methods can provide full-wave solutions with a fixed number of degrees of freedom for a wide range of frequencies. Unlike asymptotic high-frequency methods which may not be rigorous and accurate for moderate frequencies, they provide accurate and reliable approximations for moderate frequencies. We present several numerical examples to demonstrate the performance of the proposed approach.

#### *D. Embedded Discontinuous Galerkin Methods (EDG)*

We have presented an embedded discontinuous Galerkin (EDG) method for numerically solving the Euler equations and the Navier-Stokes equations [14]. The essential ingredients are a local Galerkin projection of the underlying governing equations at the element level onto spaces of polynomials of degree  $k$  to parametrize the numerical solution in terms of the approximate trace, a judicious choice of the numerical flux to provide stability and consistency, and a global jump condition that weakly enforces the single-valuedness of the numerical flux to arrive at a global formulation in terms of the numerical trace. The EDG method is thus obtained from a Hybridizable Discontinuous Galerkin (HDG) method by requiring the approximate trace to belong to a smaller space than the one in the HDG method. In the EDG method, the numerical trace is taken to be continuous, thus resulting in an even smaller number of globally coupled degrees of freedom than in the HDG method. On the other hand, the EDG method is no longer locally conservative. In the framework of convection-diffusion problems, this lack of local conservativity is reflected in the fact that the EDG method does not provide the optimal convergence of the approximate gradient or the superconvergence for the scalar variable for diffusion-dominated problems as the HDG method does. However, since the HDG method does not display these properties in the convection-dominated regime, the associated EDG method becomes a reasonable alternative since it produces smaller algebraic systems than the HDG method. In fact, the resulting stiffness matrix has the same sparsity structure as that of the statically condensed continuous Galerkin (CG) method. The main advantage of the EDG method is that it is more stable and robust than the CG method for solving convection-dominated problems. Numerical results indicate that, even though the EDG method is not locally conservative, it is a viable alternative to the HDG method in the convection-dominated regime.

#### *E. Shock Capturing*

We have introduced a novel shock capturing artificial viscosity technique [15] for high-order unstructured mesh methods. This artificial viscosity model is based on a non-dimensional form of the divergence of the velocity. The approach presented has a number of attractive properties, over previously presented methods, including non-dimensional analytical form, sub-cell resolution, and robustness for complex shock flows on highly anisotropic meshes. We presented extensive numerical results to demonstrate the performance of the proposed approach.

#### *F. Explicit HDG*

We have presented an explicit hybridizable discontinuous Galerkin (HDG) method for the acoustic wave equation [16]. The method provides optimal convergence of order  $k + 1$  for all the approximate variables including the gradient of the solution, and, when the time-stepping method is of order  $k + 2$ , it displays a superconvergence property which allow us, by means of local post-processing, to obtain new improved approximations of the scalar field variables at any time level. In particular, the new approximations converge with order  $k + 2$  in the  $L_2$ -norm for  $k$  equal to 1 or greater. These properties do not hold for all numerical fluxes. Indeed, our results show that, when the HDG numerical flux is replaced by the Lax-Friedrichs flux, the above-mentioned super-convergence properties are lost.

### G. Electromagnetics: An Application in Nanophotonics

We have used our HDG method to solve Maxwell's equations for a nanophotonics application [17]. First, we experimentally show that terahertz (THz) waves confined in sub-10 nm metallic gaps can detect refractive index changes caused by only a 1 nm thick ( $\sim \lambda/106$ ) dielectric overlayer. We use atomic layer lithography to fabricate a wafer-scale array of annular nanogaps. Using THz time-domain spectroscopy in conjunction with atomic layer deposition, we measure spectral shifts of a THz resonance peak with increasing  $\text{Al}_2\text{O}_3$  film thickness in 1 nm intervals. Because of the enormous mismatch in length scales between THz waves and sub-10 nm gaps, conventional modeling techniques cannot readily be used to analyze our results. We employ our Hybridizable Discontinuous Galerkin (HDG) scheme, for full three-dimensional modeling of the resonant transmission of THz waves through an annular gap that is 2 nm in width and 32  $\mu\text{m}$  in diameter. Our multiscale 3D FEM technique and atomic layer lithography will enable a series of new investigations in THz nanophotonics that has not been possible before.

### H. Uncertainty Quantification

In the last year of the grant, we have presented a new statistical approach to the problem of incorporating experimental observations into a mathematical model described by linear partial differential equations (PDEs) to improve the prediction of the state of a physical system [18]. We augment the linear PDE with a functional that accounts for the uncertainty in the mathematical model and is modeled as a Gaussian process. We develop a Gaussian functional regression method to determine the posterior mean and covariance of the Gaussian functional, thereby solving the stochastic PDE to obtain the posterior distribution for our prediction of the physical state. Our method has the following features which distinguish itself from other regression methods. First, it incorporates both the mathematical model and the observations into the regression procedure. Second, it can handle the observations given in the form of linear functionals of the field variable. Third, the method is non-parametric in the sense that it provides a systematic way to optimally determine the prior covariance operator of the Gaussian functional based on the observations. Fourth, it provides the posterior distribution quantifying the magnitude of uncertainty in our prediction of the physical state. In [19], we built on the previous work but considered the problem of experimental design. We developed an algorithm for designing experiments to efficiently reduce the variance of our state estimate. We provide several examples from heat conduction, the convection-diffusion equation, and the reduced wave equation, all of which demonstrate the performance of the proposed methodology. Finally, in [20] we introduced a Gaussian functional regression (GFR) technique that integrates multi-fidelity models with model reduction to efficiently predict the input-output relationship of a high-fidelity model. We developed a greedy sampling algorithm to select the training inputs. Our approach results in an output prediction model that inherits the fidelity of the high-fidelity model and has the computational complexity of the reduced basis approximation.

We expect to continue this line of research in the future.

### REFERENCES

- [1] B. COCKBURN, J. GOPALAKRISHNAN, AND R. LAZAROV. Unified hybridization of discontinuous Galerkin, mixed and continuous Galerkin methods for second order elliptic problems. *SIAM J. Numer. Anal.*, 47, 1319–1365, 2009.
- [2] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *An implicit high-order hybridizable discontinuous Galerkin method for linear convection-diffusion equations*, *J. Comput. Phys.*, 228 (2009), pp. 3232–3254.
- [3] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *An implicit high-order hybridizable discontinuous Galerkin method for nonlinear convection-diffusion equations*, *J. Comput. Phys.*, 228 (2009), pp. 8841–8855.
- [4] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *A comparison of HDG methods for Stokes flow*, *J. Sci. Comput.*, 45 (2010), pp. 215–237.
- [5] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *A hybridizable discontinuous Galerkin method for Stokes flow*, *Comput. Methods Appl. Mech. Engrg.*, 199 (2010), pp. 582–597.
- [6] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *High-order implicit hybridizable discontinuous Galerkin methods for acoustics and elastodynamics*, *J. Comput. Phys.*, 230 (2011), pp. 3695–3718.

- [7] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *Hybridizable discontinuous Galerkin methods for the time-harmonic Maxwell's equations*, J. Comput. Phys., 230 (2011), pp. 7151–7175.
- [8] N.-C. NGUYEN, J. PERAIRE, AND B. COCKBURN, *An implicit high-order hybridizable discontinuous Galerkin method for the incompressible Navier-Stokes equations*, J. Comput. Phys., 230 (2011), pp. 1147–1170.
- [9] J. PERAIRE, N. C. NGUYEN, AND B. COCKBURN. *An Embedded Discontinuous Galerkin Method for the compressible Euler and Navier-Stokes equations* (AIAA Paper 2011-3228). In *Proceedings of the 20th AIAA Computational Fluid Dynamics Conference*, Honolulu, Hawaii, June 2011.
- [10] J. Peraire, N. C. Nguyen, and B. Cockburn. *A hybridizable discontinuous Galerkin method for the compressible Euler and Navier-Stokes equations* (AIAA Paper 2010-363). In *Proceedings of the 48th AIAA Aerospace Sciences Meeting and Exhibit*, Orlando, Florida, January 2010.

PUBLICATIONS DEVELOPED UNDER THE GRANT

- [11] N.C. NGUYEN, X. ROCA, D. MORO AND J. PERAIRE, *A hybridized multiscale discontinuous Galerkin method for compressible flows*, AIAA-2013-0689 Paper, 2013.
- [12] J. GOPALAKRISHNAN, F. LI, N.-C. NGUYEN, AND J. PERAIRE, *Spectral Approximations by the HDG Method*, Math. Comp. 84, 1037–1059, 2015.
- [13] N.C. NGUYEN, J. PERAIRE, B. COCKBURN AND F. REITICH, *A phase-based hybridizable discontinuous Galerkin method for the numerical solution of the Helmholtz equation*, J. Comp. Phys., 290, 318–335, 2015.
- [14] N.C. NGUYEN, J. PERAIRE AND B. COCKBURN, *A Class of Embedded Discontinuous Galerkin Methods for Computational Fluid Dynamics*, submitted to J. Comp. Phys. 2015.
- [15] D. MORO, N.-C. NGUYEN, AND J. PERAIRE, *Dilatation-based shock capturing for high-order methods*, submitted to Int. J. for Num. Meth. in Fluids, 2015.
- [16] M. STANGLMEIER, N.-C. NGUYEN, J. PERAIRE, B. COCKBURN, *An explicit hybridizable discontinuous Galerkin method for the acoustic wave Equation*, submitted Comp. Meth. Appl. Mech. and Engrg. 2015.
- [17] H.-R. PARK, X. CHEN, N.-C. NGUYEN, J. PERAIRE AND S.-H. OH, *Nanogap-Enhanced Terahertz Sensing of 1 nm Thick ( $\lambda/10^6$ ) Dielectric Film*, ACS Photonics, 2, 417–424, 2015
- [18] N.C. NGUYEN AND J. PERAIRE, *Gaussian functional regression for linear partial differential Equations*, Comp. Meth. Appl. Mech. Engrg., 287, 69–89, 2015.
- [19] N.C. NGUYEN, H. MEN, R. M. FREUND, AND J. PERAIRE, *Gaussian functional regression for state prediction using linear PDE models and observations*, Submitted to J. Comp. Phys., 2015
- [20] N.C. NGUYEN AND J. PERAIRE, *Gaussian functional regression for output prediction: Model assimilation and experimental design*, Manuscript in Draft form 2015.

# AFOSR Deliverables Submission Survey

Thank You

Your report has been submitted. You should receive an email confirmation soon that it is being processed by AFOSR. Please print this page as proof of submission. Thank you.

## Principal

Investigator Name: Jaime Peraire

## Primary

Contact E-mail: peraire@mit.edu

## Primary

Contact Phone Number: 617-253-1981

Grant/Contract Title: HYBRIDIZED DISCONTINUOUS GALERKIN METHODS FOR MULTIPHYSICS

Grant/Contract Number: FA9550-12-1-0357

Program Manager: Jean-Luc Cambier

Report Type: Final Report

Reporting Period Start Date: 07/01/2012

Reporting Period End Date: 06/30/2015

## Abstract:

We have continued the development of the Hybridized Discontinuous Galerkin (HDG) method for the solution of systems of conservation laws. The focus has been on the development of robust, accurate, and efficient methods capable of handling complex physics in realistic geometries. In particular, we have: 1) proposed a Hybridized Multiscale Discontinuous Galerkin method (HMDG) which allow us to freely modify the local approximations spaces on each subdomain, 2) introduced hybridization and postprocessing techniques for the HDG approximation of second-order elliptic eigenvalue problems, 3) presented a new class of HDG methods capable of generating high-quality approximations of the Helmholtz equation for a very wide range of wave frequencies, 4) presented an embedded discontinuous Galerkin (EDG) method for numerically solving the Euler equations and the Navier-Stokes equations, 5) introduced a novel shock

capturing artificial viscosity technique for high-order unstructured mesh methods, 6) introduced an explicit hybridizable discontinuous Galerkin (HDG) method for the acoustic wave equation, and finally, 7) presented a new statistical approach to the problem of incorporating experimental observations into a mathematical model described by linear partial differential equations (PDEs) to improve the prediction of the state of a physical system. We have also illustrated our methods on number of applications in fluid mechanics and electromagnetics.

Distribution Statement: Distribution A - Approved for Public Release

SF298 Form: 184-ee57cf3760b82c8573b34f370403d58e\_Peraire\_SF298.pdf

Report Document 36-e60b249ab1d323f9241e3e08d9d910c2\_AFOSR-HDG2015-Report.pdf

Archival Publications: Listed in Report

Changes in Research objectives: None

Change in AFOSR Program Manager, if any: Original Program Manager: Dr. Fariba Fahroo  
Current Program Manager: Jean-Luc Cambier

Extensions granted or milestones slipped, if any: None

100%