

Award Number: W81XWH-12-1-0313

TITLE: Divergent Effects of Dendritic Cells on Pancreatitis

PRINCIPAL INVESTIGATOR: Dr. George Miller

CONTRACTING ORGANIZATION: New York University
New York, NY 10016-6402

REPORT DATE: September 2015

TYPE OF REPORT: Final report

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			<i>Form Approved</i> <i>OMB No. 0704-0188</i>		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE September 2015		2. REPORT TYPE Final		3. DATES COVERED 15 Aug 2012 – 14 Aug. 2015	
4. TITLE AND SUBTITLE Divergent Effects of Dendritic Cells on Pancreatitis			5a. CONTRACT NUMBER W81XWH-12-1-0313		
			5b. GRANT NUMBER 10911379		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) George Miller E-Mail: george.miller@nyumc.org			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) New York University New York, NY 10016-6481			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Numbers dendritic cells (DCs) increased 100-fold in pancreata of mice with acute pancreatitis to account for nearly 15% of intrapancreatic leukocytes. Intrapaneatic DCs acquired a distinct immune phenotype in mice with acute pancreatitis; they expressed higher levels of major histocompatibility complex II and CD86 and increased production of interleukin-6, MCP-1, and TNFa. DC over-expansion exacerbated disease. However, DCs were also required for pancreatic viability; the exocrine pancreas died in mice that were depleted of DCs and challenged with caerulein or L-arginine. All mice with pancreatitis that were depleted of DCs died from acinar cell death within 4 days. These data suggest that DC have simultaneous paradoxical pro-inflammatory and protective effects in pancreatitis. Further investigations are warranted.					
15. SUBJECT TERMS Dendritic cells, pancreatitis					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)
			UU	37	

Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	2
Key Research Accomplishments.....	12
Reportable Outcomes.....	13
Conclusion.....	14
References.....	15
Appendices.....	16


Introduction

Pancreatitis is a significant public health concern and also is the most common precursor for pancreatic cancer. Our work investigates the role of dendritic cells in pancreatitis. Dendritic cells are professional antigen presenting cells which initiate innate and adaptive immune responses. Our work utilizes murine models and human tissues. Dendritic cells in mice express MHC II and the integrin CD11c. They are proficient in activating T cells and NK cells yet their role in pancreatitis has not previously been investigated.

Body


Task 1.1.

Studies of DC expansion in pancreatitis: To determine the prominence of dendritic cells (DC) in the context of pancreatic inflammation, we induced pancreatitis in mice using caerulein and characterized the leukocytes in the pancreas and spleen using flow cytometry. We found that DC increase substantially in number in acute and chronic pancreatitis (**Figure 1**).




Studies of DC phenotype in pancreatitis: We tested DC expression of MHC molecules, integrins (CD40, CD54), and co-stimulatory molecules (CD80, CD86) in the pancreas and spleen in control mice and in models of pancreatitis. We showed that DC matured in acute pancreatitis (**Figure 2**).

Figure 2. Intra-pancreatic DC mature in pancreatitis. DC from the pancreata of saline (dotted lines) and caerulein-treated (solid lines) mice were gated and analyzed for surface marker expression by flow cytometry. DC mature and undergo a CD11b+CD8 α - myeloid shift in AP. Shaded histograms represent Isotype controls.




Studies of DC immune function in pancreatitis: (i) We measured DC production of immunogenic and regulatory cytokines and chemokines using a cytometric bead array and found that DC express high levels of inflammatory mediators in pancreatitis (**Figure 3a**). (ii) We measured DC expression of Toll-like



receptors (TLRs) using flow cytometry and showed that pancreatic DC express increased TLRs in pancreatitis (**Figure 3b**).

Studies of human DC in pancreatitis: To investigate whether DC expand in human pancreatitis, we obtained specimens from human pancreatitis from the NYU Tissue Bank. We analyzed these for the presence of a DC infiltrate by immunohistochemistry using antibodies directed against DC-SIGN (CD209), CD1a, and CD123 which are diverse human DC markers (**Figure 4**).


Figure 4. DC are prominent in the human pancreas. Immune cells in human pancreatitis expressed high DC-SIGN (CD209), and lower CD1a, and CD123.




Task 1.2. To determine effects of DC expansion in pancreatitis

Bone Marrow-derived Dendritic Cell (BMDC) adoptive transfer exacerbates pancreatitis - We generated BMDC *in vitro* from BM progenitors using GM-CSF (20 ng/ml) in 8 day cultures. Mice were adoptively transferred with 1×10^6 BMDC after daily caerulein injections in models of chronic pancreatitis and severity of disease was determined by histological analysis. We showed that DC adoptive transfer exacerbated pancreatic inflammation and organ destruction (**Figure 5a-f**).

Figure 5. DC exacerbate pancreatitis. Adoptive transfer of DC during pancreatitis induction resulted in worsened pancreatic inflammation (H&E staining), increased fibrosis (Sirius red staining), and increased organ destruction (evidenced by amylase and insulin staining. "C" represents pancreata of mice treated with caerulein pancreatitis only and "C+DC" represents pancreata of mice treated with caerulein + DC (n=8-10 mice/group).



FMS-like tyrosine kinase 3 ligand (Flt3L) mediated DC expansion exacerbates pancreatitis –To further elucidate the effects of DC on pancreatic injury, we administered Flt3L *in vivo*. Flt3L is a powerful DC growth factor that results in DC expansion in lymphoid and non-lymphoid organs. This recruited DC to the pancreas and exacerbated the severity of disease (**Figure 6**).





Task 1.3. To determine the effects of DC ablation in pancreatitis


Examination of DC Depletion in Pancreatitis: We used CD11c.DTR mice, in which transient DC depletion can be effected for 48h during induction of pancreatitis by treatment with diphtheria toxin (DT; 4 ng/g). We tested the effects of DC depletion on severity of pancreatitis by histological analysis and survival showed that DC depletion results in exacerbated disease (**Figure 7a, b**).

Figure 7. DC depletion exacerbates pancreatitis. (a) Pancreatitis was induced using L-arginine in controls or in the context of DC depletion. DC depletion resulted in exacerbated disease. (b) DC depletion also resulted in animal death in a survival experiment.

a



b




Task 2: To determine the role of DC in modulating both CD4⁺ effector T cell differentiation and sterile inflammation in pancreatitis.


Task 2.1. To examine the primacy of the DC-Th2 axis in exacerbating pancreatitis

(i) To elucidate whether intra-pancreatic DC differentially induce Th2 differentiation, we harvested intra-pancreatic CD4⁺ T cells from controls, pancreatitis mice (denoted "C" for caerulein), mice treated with DC alone (denoted "DC"), and pancreatitis mice adoptively transferred with DC ("C+DC"), and tested CD4⁺ T cells for Th1, Th2, and Th17 differentiation in cell culture supernatant. We found that DC transfer increased Th2 differentiation in pancreatitis by enhancing production of IL-4 and IL-10 which are Th2 cytokines (**Figure 8**).


Further, use of DC from MyD88^{-/-} mice further enhanced Th2 differentiation. 25 mice were used for these experiments.




(ii) To test whether DC are required for Th-differentiation of pancreatic T cells, we harvested intra-pancreatic CD4⁺ T cells from controls (Ctl), pancreatitis mice (C), and pancreatitis mice depleted of DC ("C-DC") and tested CD4⁺ T cells for Th1, Th2, and Th17 differentiation using a cytometric bead array. These experiments showed that depleting DC enhanced Th1 deviation as IL-6 and TNF- α were increased (**Figure 9**). 25 mice were used for these experiments.




(iii) To test whether DC in pancreatitis induce Th2 effector cells specific for pancreatic antigen, FACS-sorted pancreas-infiltrating CD4⁺ T cells from pancreatic mice were cultured with pancreatic lysate- or mock-pulsed DCs. CD4⁺ T cell activation was then determined by their expression of CD69, FAS ligand, and CD11b. Gray histograms represent isotype control. Median fluorescent indexes are shown for each group. These data show the antigen specificity of the T cells as T cells became activated by DC loaded with pancreatic lysate suggesting that the CD4 T cells respond to the presentation of pancreatic antigen (**Figure 10**). 90 mice were used for these experiments.




(iv) **MHC II Knockout mice** To further test whether DC induction of Th2-deviated CD4⁺ T cells is the critical cellular mechanism behind DC exacerbation of pancreatitis, we employed MHC II^{-/-} mice and β 2-microglobulin^{-/-} controls. MHC II^{-/-} and β 2-microglobulin^{-/-} mice treated with caerulein were recipients of DC adoptive transfer (1×10^6 cells) to enhance pancreatic inflammation. We found that MHC II^{-/-} mice (which lack Th2 signaling) were protected but not β 2-microglobulin^{-/-} mice (which lack Th1 signaling) confirming the primacy of the DC-Th2 axis (**Figure 11**). 50 Mice were used for these experiments.



(v) **DC inactivation:** We administered VAG539 which blocks Th2-deviation twice daily via oral gavage for the duration of the course of pancreatitis induction. We found reduced pancreatic injury on H&E in mice treated with caerulein + VAG539 with reduced evidence of intra-pancreatic inflammation and diminished Th2-deviation as evidenced by lower levels of IL-10 compared with mice treated with caerulein alone (**Figure 12**). 50 mice were used for these experiments.




(vi) *MyD88 Blockade*: Based on the known literature, MyD88 signaling is associated with strong Th1 polarization. Conversely, MyD88 inhibition or deletion in DC is associated with their Th2 deviation. Therefore, we postulated that MyD88 inhibition will worsen pancreatitis by favoring DC-Th2 deviation within the pancreas. We assessed the effects of MYD88 blockade on pancreatitis by histological analysis. We found that MyD88 inhibition resulted in exacerbated pancreatitis (**Figure 13**). Th2-deviation was also enhanced by MyD88 blockade (see Figure 1 above). 50 mice were used for these experiments.




Task 2.2. To determine whether DC clearance of necrotic cellular debris and apoptotic targets limits expansion of sterile inflammation in pancreatitis

(i) *Intra-pancreatic DC capacity for clearance of apoptotic and necrotic cells*: We tested the relative capacity of pancreatic DC to capture antigen as well as necrotic and apoptotic cells in pancreatitis. CD11c⁺MHCII⁺ DC harvested from pancreata of mice treated with caerulein captured FITC-Dextran at a higher rate than CD11c⁺MHCII⁺ antigen presenting cells from the same pancreata (**Figure 14A**). In addition, we found that pancreatic DC captured 7AAD⁺ necrotic cells at a far greater rate than other antigen presenting cells in pancreatitis (**Figure 14B**). Moreover, the uptake of 7AAD⁺ necrotic cellular debris (**Figure 14C**) and Annexin⁺ apoptotic bodies (**Figure 14D**) by MHC II⁺ cells was severely deficient in pancreata depleted of DC compared with pancreata with a normal complement of DC. Taken together, these data suggest that DC are primary in clearance of necrotic and apoptotic cells thereby limiting sterile inflammation in pancreatitis. 50 mice were used for these experiments.




(ii) *Changes in sterile inflammation in pancreatitis upon DC depletion* - DC have recently been assigned an important role in the clearance of cellular debris in inflammatory disease. Since we had previously showed that DC expand most sharply during the regeneration phase of pancreatitis, and DC depletion results in an marked increase in apoptotic cells and necrotic elements, we postulated that DC may have a primary role in clearance of cellular debris in pancreatitis, and, consequently, DC depletion may result in expansion of sterile inflammation in AP. In support of this notion, we found that C-DC treated pancreata have markedly higher levels of HMGB-1 (**Figure 15A**). We next determined the effects of DC depletion in pancreatitis ("C-DC") on systemic inflammation. Mice were challenged for two days with caerulein after DC depletion or mock depletion. Serum levels of MCP-1 were measured at 48 hours. There was increased evidence of systemic inflammation when DC were depleted in pancreatitis (**Figure 15B**). Taken together, these data suggest that DC depletion increases inflammation in pancreatitis. Assays were performed in triplicate and repeated three times. 50 mice were used for these experiments.



Task 3: To determine the role of DC-PSC cross-talk in initiating fibrosis in chronic pancreatitis.

Task 3.1. To determine whether pancreatic DC can active PSC in vitro

(i) *Pancreatic stellate cell (PSC) Surface Phenotype*: Using flow cytometry, we found that DC induce PSC expression of ICAM-1 (**Figure 16a**). DC also increased PSC expression of PDGF (**Figure 16b**). PSC morphology was unchanged (**Figure 16c**).



(ii) *PSC Proliferation and Migration* We found that DC reduce PSC proliferation (**Figure 17a**) but increase PSC migrational capacity in vitro using a transwell co-culture system (**Figure 17b**).

Fig. 17a







Fig. 17b




(iii) *PSC Production of Inflammatory Mediators* To test whether intra-pancreatic DC from mice with pancreatitis induce PSC to adopt a pro-inflammatory phenotype, PSC production of an array cytokines and chemokines specifically linked to progression of pancreatitis were measured after co-culture with DC. PSC production of inflammatory mediators were increased after DC co-culture (**Figure 18**).



(iv) *PSC Production of Extracellular Matrix Proteins* The ultimate effect of PSC activation in chronic pancreatitis is the deposition of extracellular matrix (ECM) proteins and modulators of the ECM such as matrix metalloproteinases (MMPs) and tissue inhibitors of metalloproteinases (TIMPs). We measured whether DC induce PSC production of ECM related genes but found no significant changes. Results of our PCR array comparing PSC + DC to PSC alone are shown (**Figure 19**).



(v) *Human Correlative DC-PSC Studies* We extended our analysis to human systems. For human DC stimulators, 2x2 cm sections of non-tumorous pancreas were digested using Collagenase and Lin⁺HLA-DR⁺ bulk pancreatic DC were isolated by FACS. As in our murine experiments, DC will be added to PSC for 48h in a 1:1 ratio before washing off the non-adherent DC. PSC were then tested 24h later for Collagen I production by western blotting. We found that DC do not induce PSC to make collagen I (**Figure 20**).




Task 3.2. To determine whether intra-pancreatic DC expansion activates PSC *in vivo*


DC PSC experiments in vivo: To test DC capacity to activate PSC *in vivo* during pancreatitis, we employed our experimental strategy of DC over-expansion in pancreatitis and, using immunohistochemistry, we tested PSC expression of Desmin and α -SMA. To further test the ability of DC to activate PSC *in vivo*, we isolated PSC from normal controls, mice with pancreatitis, and mice with pancreatitis supplemented with DC over-expansion and measure both the time course to α -SMA expression *in vitro* after PSC isolation as well as levels of PDGF and TGF- β in day 6 PSC cultures. However, all our *in vivo* data was negative and we did not appreciate DC activation of PSC *in vivo*.

Task 3.3. To determine the biochemical mechanism of DC induction of PSC

(i) *TNF- α and IL-6:* We postulated that intra-pancreatic DC production of TNF- α and IL-6 are necessary for maximal DC-induced PSC activation. We found that CpG treatment increases PSC expression of TNF- α and other cytokines shown in **Figure 21**. However, IL-6 was not upregulated.



(ii) *Cell Signaling Pathways* : Determining which PSC signaling pathway are induced by DC in pancreatitis can have therapeutic implications for targeting in experimental therapeutics. We found that ICAM expression in DC is necessary for their capacity to induce cytokine production in PSC (**Figure 22a**). Similarly, MyD88 expression in DC is necessary for their capacity to induce cytokine production in PSC (**Figure 22b**).



Key Research Accomplishments

- We showed DC expand and activate in mouse pancreatitis
- We showed that DC are prominent in human pancreatitis
- We showed that a balance of DC is needed in the pancreas as DC expansion results in worsened disease but complete DC ablation leads to organ destruction and death
- We showed DC induce Th2 differentiation in the pancreas
- We showed that DC depletion in pancreatitis results in upregulation of Th1 cytokines
- We showed that DC generate CD4 T cells that are specific to pancreatic antigen in pancreatitis
- We showed that blockade of the DC-Th2 axis ameliorates pancreatitis
- We showed that enhancement of the DC-Th2 axis exacerbates pancreatitis
- We showed that DC are primary in the clearance of necrotic debris in pancreatic cancer
- We showed that DC depletion in pancreatitis exacerbates sterile inflammation
- We showed that DC induce PSC expression of cytokines and chemokines
- We showed that DC do not induce PSC expression of ECM proteins
- We showed that DC induce PSC expression of cytokines and chemokines in a ICAM and MyD88 dependent manner

Reportable Outcomes

Publications:

1. Ochi A, Nguyen AH, Bedrosian AS, Mushlin HM, Zerbakhsh S, Barilla R, Zambirinis CP, Fallon NC, Rehman A, Pylayeva-Gupta Y, Badar S, Hajdu CH, Frey AB, Bar-Sagi D, Miller G. MyD88 inhibition amplifies dendritic cell capacity to promote pancreatic carcinogenesis via Th2 cells. *J Exp Med.* 209(9):1671-87.

2. Zambirinis CP, Levie E, Nguy S, Avanzi A, Barilla R, Xu Y, Siefert L, Daley D, Greco S, Deutsch M, Jonnadula S, Torres-Hernandez A, Tippens D, Pushalkar S, Eisenthal A, Saxena D, Ahn J, Hajdu C, Engle DD, Tuveson D, Miller G. TLR9 Ligation in Pancreatic Stellate Cells Promotes Tumorigenesis *Journal of Experimental Medicine.* 2015, **In Press**

Presentation at National Meeting:

1. George Miller: Innate immune Signaling in the Pancreatic Tumor Microenvironment. AACR Special Conference on Pancreatic Cancer. May 2014, New Orleans, LA

This was an invited oral presentation at a national meeting. There was no published abstract.

2. Dendritic cells contribute to pancreatic fibro-inflammatory disease and the transition to neoplasia. Saman Zerbakhsh, Harry Mushlin, Raghavendra Rao, Sanna Badar, Mohsin Jamal, Dafna Bar-Sagi, George Miller, Adeel Rehman, Christopher Graffeo, Atsuo Ochi, Rocky Barilla, Constantinos Zambirinis, Nina Fallon, Cristina Hajdu, Yuliya Pylayeva-Gupta. AACR pancreatic cancer meeting in May 2013.

There was no published abstracts

Conclusion

DC are an important component of the immune infiltrate in pancreatitis by induction of Th2 differentiation and in the clearance of necrotic debris. DC also activate PSC to produce inflammatory mediators.

References

None

Appendices

Our publications follows (our in press publication is not published yet)

MyD88 inhibition amplifies dendritic cell capacity to promote pancreatic carcinogenesis via Th2 cells

Atsuo Ochi,¹ Andrew H. Nguyen,² Andrea S. Bedrosian,¹ Harry M. Mushlin,² Saman Zarbakhsh,¹ Rocky Barilla,¹ Constantinos P. Zambirinis,¹ Nina C. Fallon,¹ Adeel Rehman,¹ Yuliya Pylayeva-Gupta,³ Sana Badar,¹ Cristina H. Hajdu,⁴ Alan B. Frey,² Dafna Bar-Sagi,³ and George Miller^{1,2}

¹Department of Surgery, ²Department of Cell Biology, ³Department of Biochemistry and Molecular Pharmacology, and ⁴Department of Pathology, New York University School of Medicine, New York, NY 10016

The transition of chronic pancreatic fibroinflammatory disease to neoplasia is a primary example of the paradigm linking inflammation to carcinogenesis. However, the cellular and molecular mediators bridging these entities are not well understood. Because TLR4 ligation can exacerbate pancreatic inflammation, we postulated that TLR4 activation drives pancreatic carcinogenesis. In this study, we show that lipopolysaccharide accelerates pancreatic tumorigenesis, whereas TLR4 inhibition is protective. Furthermore, blockade of the MyD88-independent TRIF pathway is protective against pancreatic cancer, whereas blockade of the MyD88-dependent pathway surprisingly exacerbates pancreatic inflammation and malignant progression. The protumorigenic and fibroinflammatory effects of MyD88 inhibition are mediated by dendritic cells (DCs), which induce pancreatic antigen-restricted Th2-deviated CD4⁺ T cells and promote the transition from pancreatitis to carcinoma. Our data implicate a primary role for DCs in pancreatic carcinogenesis and illustrate divergent pathways in which blockade of TLR4 signaling via TRIF is protective against pancreatic cancer and, conversely, MyD88 inhibition exacerbates pancreatic inflammation and neoplastic transformation by augmenting the DC-Th2 axis.

CORRESPONDENCE

George Miller:
george.miller@nyumc.org

Abbreviations used: HPF, high-powered field; MAP, mitogen-activated protein; PanIN, pancreatic intraepithelial neoplasia; PDEC, pancreatic ductal epithelial cell; Poly I:C, polyinosinic:polycytidylic acid.

Chronic pancreatitis is a fibroinflammatory disorder characterized by irreversible pancreatic destruction leading to both exocrine and endocrine organ failure. The hallmark of chronic pancreatitis is repeated cyclical fibroinflammatory responses to a primary insult such as alcoholic injury (Braganza et al., 2011). Intrapaneatic inflammation requires the activation of transcription factors, including NF- κ B and AP-1 which can be induced by the RANK ligand and TNF (Vaquero et al., 2001). However, the precise cellular and biochemical mediators in chronic pancreatitis are uncertain.

An even more interesting feature of fibroinflammatory pancreatic disease is the observation that chronic pancreatitis is the leading risk factor for the development of pancreatic cancer. Epidemiological studies suggest that the incidence of pancreatic carcinoma in individuals suffering

from chronic pancreatitis ranges between 1.4 and 2.7% (Ammann et al., 1984; Lankisch et al., 1993). Overall, individuals with chronic pancreatitis have a 16-fold greater lifetime risk of developing pancreatic adenocarcinoma than the general population (Jura et al., 2005). As such, the transition of chronic pancreatitis to pancreatic carcinoma represents an exemplary clinical example of the paradigm of chronic inflammation degenerating into neoplasia. Although the cellular and biochemical links between benign and malignant pancreatic disease remain poorly understood, a distinct feature of pancreatic cancer, and a commonality with chronic pancreatitis, is the extensive fibroinflammatory stroma surrounding the transformed ductal epithelial cells (Ricci et al., 2005). This suggests that the

© 2012 Ochi et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons License (Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>).

A. Ochi and A.H. Nguyen contributed equally to this paper.

inflammatory stroma may have a causative role in the neoplastic transformation of chronic pancreatitis.

DCs have recently emerged as important mediators of organ-specific fibroinflammatory disease. DCs are reported to be critical to toxin-induced pulmonary inflammation and the resulting interstitial pulmonary fibrosis (Bantsimba-Malanda et al., 2010). Our recent work in the study of liver disease identified DCs as important in modulating acute and chronic inflammation after hepatic insult (Connolly et al., 2009, 2011). TLR4 activation can also drive inflammation, and recent work in pancreatitis demonstrated that TLR4 ligation stimulates pancreatic inflammation (Sharif et al., 2009; Ding et al., 2010; Zhou et al., 2010). Based on this, we postulated a role for both DCs and TLR4 in neoplastic transformation of the pancreas. Our data show that TLR4 ligation is required for neoplastic progression of pancreatic cancer as blockade of either TLR4 or the MyD88-independent TRIF pathway is protective by regulating fibroinflammatory stromal expansion. Conversely, blockade of MyD88 surprisingly accelerates pancreatic tumor progression by augmenting DC capacity to generate intrapancreatic inflammation via induction of Th2-deviated CD4⁺ T cells.

RESULTS

TLR4 regulates pancreatic carcinogenesis

To determine the effects of TLR4 ligation on pancreatic tumor progression, we treated 4-wk-old p48Cre;Kras^{G12D} mice with LPS and harvested pancreata 4 wk later. LPS greatly accelerated tumorigenesis as treated mice developed advanced pancreatic intraepithelial neoplasia (PanIN) lesions encased in a dense bed of fibroinflammatory stroma (Fig. 1, A–E). Conversely, saline-treated mice had grossly normal pancreata. Pancreata from LPS-treated p48Cre;Kras^{G12D} mice also weighed roughly three times as much as controls (Fig. 1 B). Moreover, blockade of TLR4 *in vivo* protected p48Cre;Kras^{G12D} mice against pancreatic tumorigenesis (Fig. 1, F and G). We postulated that the regulatory effects of TLR4 on pancreatic carcinogenesis result from modulation of inflammatory cell activation within the desmoplastic tumor stroma. To test this, we measured TLR4 expression on stromal leukocytes in normal pancreata and in pancreatic cancer. B cells, DCs, macrophages, and granulocytes each markedly increased their expression of TLR4 within the pancreatic cancer microenvironment (Fig. 1 H). Similarly, in human pancreatic cancer, there was robust expression of TLR4 on stromal leukocytes (Fig. 1 I). Moreover, we found elevated levels of TLR4 agonists (Fig. 1 J) and specific well-characterized DAMPs that bind TLR4 (Fig. 1 K) in human pancreatic duct fluid from cancer patients, suggesting there is ample substrate for TLR4 activation within the pancreatic cancer tumor microenvironment.

To definitively implicate peritumoral inflammation in mediating the effects of TLR4 activation on tumor progression, we made p48Cre;Kras^{G12D} mice chimeric with TLR4^{-/-} mice such that inflammatory cells within the tumor microenvironment would be deficient in TLR4 signaling, whereas neoplastic epithelial cells would have intact TLR4 signaling.

For controls, we made age-matched p48Cre;Kras^{G12D} mice chimeric using bone marrow derived from WT mice. Selected cohorts of chimeric mice were treated with caerulein to accelerate carcinogenesis as described previously (Carrière et al., 2009). WT chimeric p48Cre;Kras^{G12D} mice treated with caerulein developed diffuse PanIN lesions, as expected. However, TLR4^{-/-} chimeric p48Cre;Kras^{G12D} mice were protected, exhibiting residual metaplastic ducts with cuboidal epithelia, suggesting that inflammatory cell TLR4 signaling can regulate pancreatic tumor progression (Fig. 2, A and B). Our extent of chimerism was >95% (not depicted; Bedrosian et al., 2011), and we confirmed deficient responses to LPS stimulation in TLR4^{-/-} chimeric mice (Fig. 2 C). To further test whether the absence of TLR4 signaling in inflammatory cells is sufficient to retard tumor progression, we grafted WT and TLR4^{-/-} mice with primary pancreatic ductal epithelial cells (PDECs) harboring oncogenic KRas^{G12D}, which we have shown form well-differentiated pancreatic lesions encased in a large stromal component mimicking early human pancreatic adenocarcinoma (Agbunag et al., 2006; Pylayeva-Gupta et al., 2012). Tumor growth was markedly slower in TLR4^{-/-} hosts, again indicating that inflammatory cell expression of TLR4 can modulate cancer progression (Fig. 2 D). Furthermore, because TLR4 generates inflammation via NF-κB and mitogen-activated protein (MAP) kinase, as anticipated, blockade of NF-κB and MAP kinase signaling partially protected LPS-treated p48Cre;Kras^{G12D} mice from accelerated carcinogenesis (not depicted).

Dichotomous effects of TRIF and MyD88 signaling interruption

To test whether effects of TLR4 activation in pancreatic cancer are primarily mediated via the MyD88-independent TRIF pathway or the MyD88-dependent pathway, we selectively blocked each mechanism in p48Cre;Kras^{G12D} mice using cell-permeable inhibitors. TRIF blockade (Fig. 2 E) prevented p48Cre;Kras^{G12D} mice from developing accelerated carcinogenesis after treatment with caerulein (Fig. 2 F) or LPS (not depicted). Similarly, adoptive transfer of Kras^{G12D} PDECs to TRIF^{-/-} mice resulted in retarded tumor growth compared with Kras^{G12D} PDEC transfer to WT mice (Fig. 2 G). Consistent with its regulatory effects in the tumor microenvironment, TRIF^{-/-} mice were also protected against pancreatitis (Fig. 2 H). Notably, TLR3 ligand (polyinosinic:polycytidylic acid [Poly I:C]), which also signals via TRIF, similarly accelerated carcinogenesis in p48Cre;Kras^{G12D} mice (Fig. 2 I).

In contrast to the protective effects of TRIF blockade, MyD88 inhibition (Fig. 3 A) markedly accelerated pancreatic carcinogenesis (Fig. 3, B–D). In particular, whereas control p48Cre;Kras^{G12D} mice treated with caerulein developed metaplastic ducts and diffuse low-grade PanIN lesions, animals treated with MyD88 inhibitory peptide developed pancreatic tumors three to four times the size of controls (Fig. 3 B). On histological examination, MyD88 blockade resulted in invasive cribriforming pancreatic adenocarcinoma which exhibited a markedly high proliferation rate and increased expression of p53 (Fig. 3 C) as well as foci of invasion as confirmed by




Figure 1. TLR4 signaling modulates pancreatic carcinogenesis. (A–E) 4-wk-old p48Cre;Kras^{G12D} mice were treated with saline or LPS and sacrificed at 4 wk. (A and B) Pancreata were stained with H&E, Trichrome, and CD45 (A) and weighed (B). (C–E) The presence of graded PanIN lesions (C), fibrotic area (D), and leukocytic infiltrate (E) were quantified by examining 10 high-powered fields (HPFs) per pancreas ($n = 6$ mice/group; ***, $P < 0.001$). (F and G) 4-wk-old p48Cre;Kras^{G12D} mice were treated with saline, caerulein, or caerulein + TLR4 inhibitor. Representative H&E-stained sections are shown, and the number of dysplastic ducts per HPF was calculated ($n = 5$ mice/group; ***, $P < 0.001$). (H) Live pancreatic mononuclear cells from 6-mo-old p48Cre;Kras^{G12D} or WT mice were gated and costained for CD45, CD11c, CD3, F480, B220, Gr1, and TLR4. Median fluorescence for TLR4 is shown for specific cellular subsets. Data are representative of experiments repeated three times. (I) Sections of normal human pancreas ($n = 3$) and human pancreatic cancer ($n = 19$) were stained for TLR4. Representative images are shown, and data were quantified (***, $P < 0.001$). (J) Pancreatic duct fluid was harvested at the time of surgical resection from four patients with pancreatic cancer and two patients with benign endocrine tumors and tested for TLR4 ligand levels on HEK-Blue reporter cells (***, $P < 0.001$). (K) Pancreatic ductal fluid from two patients with pancreatic carcinoma was harvested at the time of operative duct transection and analyzed for HMGB-1 and S100A9 expression by Western blotting. Error bars indicate standard error of the mean.




Figure 2. TLR4 regulation of pancreatic tumorigenesis and pancreatitis is mediated by stromal inflammatory cells and requires TRIF.

(A and B) p48Cre;Kras^{G12D} mice were irradiated and made chimeric by bone marrow transfer from WT or TLR4^{-/-} mice. 7 wk later, mice were treated with either saline or two doses of caerulein (C). Three weeks afterward, mice were sacrificed and pancreata were assessed by H&E. The fraction of dysplastic ducts was measured ($n = 5$ /group; ***, $P < 0.001$). Insets show higher magnification. (C) WT chimeric and TLR4^{-/-} chimeric mice were treated with 5 μ g LPS. Serum cytokine levels were measured at 6 h (***, $P < 0.001$). (D) WT or TLR4^{-/-} mice were adoptively transferred with intrapancreatic Kras^{G12D} PDECs. Pancreata were harvested and weighted at 6 wk ($n = 5$ mice/group; ***, $P < 0.001$). (E) Raji cells were stimulated for 90 s with 1 μ g/ml LPS in the presence of TRIF inhibitor (Pepinh-TRIF) or control peptide (Pepinh-Ctl). Expression of pIRF3 and β -actin was measured by Western blotting. (F) 4-wk-old p48Cre;Kras^{G12D} mice were treated with saline, caerulein + control peptide, or caerulein + TRIF inhibitor. Representative H&E-stained sections are shown, and the number of dysplastic ducts per HPF was calculated ($n = 5$ mice/group; ***, $P < 0.001$). (G) WT or TRIF^{-/-} mice were adoptively transferred with intrapancreatic Kras^{G12D} PDECs. Pancreata were harvested and weighted at 6 wk ($n = 4-5$ mice/group; ***, $P < 0.001$). (H) Acute pancreatitis was induced using caerulein in WT or TRIF^{-/-} mice. The fraction of viable acini was calculated ($n = 4$ mice/group; ***, $P < 0.001$). (I) 4-wk-old p48Cre;Kras^{G12D} mice were treated with saline or Poly I:C before sacrifice 4 wk later. Representative H&E-stained sections are shown, and the number of PanIN lesions per HPF was quantified ($n = 4$ mice/group; ***, $P < 0.001$). Error bars indicate standard error of the mean.

positive staining for CK19 outside of ductal structures (Fig. 3 D). Gene sequencing revealed no somatic mutation in p53 (not depicted). To confirm that the tumor-promoting effects of MyD88 inhibition occur within inflammatory cells, we made

p48Cre;Kras^{G12D} mice chimeric with MyD88^{-/-} bone marrow before caerulein treatment. Controls were made chimeric using WT bone marrow. MyD88^{-/-} chimeric mice developed more extensive fibroinflammatory replacement of their




Figure 3. MyD88 blockade accelerates malignant transformation and stromal inflammation. (A) Mice were treated with LPS and MyD88 inhibitory peptide (MyD88i) or control peptide. At 6 h, pancreata were assayed for expression of IRAK and p-IRAK. (B–D) p48Cre;Kras^{G12D} mice were treated with caerulein for 2 d to accelerate carcinogenesis before sacrifice 3 wk later. In addition, mice were administered either MyD88 inhibitory peptide or control peptide. (B) Tumor size was recorded. (C and D) Paraffin-embedded pancreatic sections were stained using H&E and Ki67 and using mAbs directed against p53 (C) and CK19 (D) an epithelial cell marker ($n = 6$ mice/group). (E and F) WT chimeric and MyD88^{-/-} chimeric p48Cre;Kras^{G12D} mice were treated with caerulein and sacrificed at 3 wk. The fraction of metaplastic ducts and PanIN lesions was quantified ($n = 4–6$ mice/group; ***, $P < 0.001$). (G) Kras^{G12D} PDECs were cultured with MyD88 inhibitory peptide or control peptide. Cells were pulsed with [³H]thymidine for 20 h, and its incorporation was measured. (H and I) MyD88^{-/-} and WT mice were challenged with caerulein alone or caerulein + TRIF inhibitor or control peptide for 3 wk. Fibro-inflammatory changes were quantified ($n = 4–6$ mice/group; ***, $P < 0.001$). Error bars indicate standard error of the mean.

acinar architecture and advanced PanIN lesions compared with WT chimerics (Fig. 3, E and F). Furthermore, MyD88 inhibition did not appear to have direct effects on epithelial cells as *Kras*^{G12D} PDECs did not exhibit altered viability (not depicted) or an accelerated proliferative rate in vitro in response to MyD88 inhibitor (Fig. 3 G). In consort with these findings, which suggest that MyD88 inhibition activates peritumoral inflammatory cells compared with WT mice subject to pancreatitis, MyD88^{-/-} mice developed exacerbated pancreatic fibroinflammation, even in the context of TRIF inhibition (Fig. 3, H and I). Collectively, these data show that blockade of the MyD88-dependent pathway and the MyD88-independent TRIF pathway have distinctly opposite effects in benign and malignant pancreatic disease.

DCs and CD4⁺ T cells mediate effects of MyD88 inhibition in pancreatic disease

Because Th2-deviated CD4⁺ T cells have been implicated in pancreatic inflammation (Demols et al., 2000; Oiva et al., 2010) and MyD88 inhibition skews DC-mediated T cell differentiation toward the Th2 phenotype (Kaisho et al., 2002; Kapsenberg, 2003; Chen et al., 2010), we postulated that DCs may be the central cellular entity responsible for the exacerbated pancreatic inflammation and tumorigenesis associated with MyD88 inhibition by inducing effector Th2-deviated CD4⁺ T cells. To directly test whether DCs are responsible for exacerbated pancreatic disease in MyD88-deficient mice, we generated CD11c-Cre MyD88 Floxed^{+/+} mice, which are deficient in MyD88 signaling in CD11c⁺ DCs (Hou et al., 2008). We found minimal CD11c expression in other intrapancreatic leukocytes subsets (not depicted). CD11c-Cre MyD88 Floxed^{+/+} mice were then challenged with caerulein to induce pancreatitis. In consort with our hypothesis, CD11c-Cre MyD88 Floxed^{+/+} mice experienced exacerbated pancreatic fibroinflammatory disease compared with controls (Figs. 4 A) and to a similar extent as pan-MyD88-deficient animals (not depicted). To determine whether MyD88 inhibition within DCs is also associated with accelerated pancreatic carcinogenesis, we made p48Cre;*Kras*^{G12D} mice chimeric using bone marrow from either WT or CD11c-Cre MyD88 Floxed^{+/+} mice followed by administration of caerulein 7 wk later. CD11c-Cre MyD88 Floxed^{+/+} chimeric p48Cre;*Kras*^{G12D} mice experienced accelerated cancer progression when compared with WT chimeric controls (Fig. 4 B).

Consistent with our hypothesis, we found that CD4⁺ T cell recruitment was increased >20-fold in pancreata of caerulein-treated MyD88^{-/-} mice compared with WT controls (not depicted) and MyD88 inhibition resulted in a significant Th2 deviation in both benign and malignant pancreatic disease (Fig. 4 C). Moreover, CD4⁺ T cell depletion (Miller et al., 2003) rescued mice from the exacerbated fibroinflammation (not depicted) and accelerated cancer progression (Fig. 4 D) associated with MyD88 inhibition. However, CD4⁺ T cell depletion did not rescue p48Cre;*Kras*^{G12D} mice with intact MyD88 signaling capacity (Fig. 4 D). Collectively, the aforementioned observations indicate that (a) MyD88 inhibition

exclusively within DCs is sufficient to induce exacerbated pancreatic inflammation and carcinogenesis (Fig. 4, A and B) and (b) MyD88 blockade exacerbates pancreatic disease in a CD4⁺ T cell-dependent manner (Fig. 4 D).

To directly test whether the DC–Th2 axis is central to pancreatic inflammation, before beginning caerulein challenge in WT mice, we treated mice with VAG539, which binds the DC aryl hydrocarbon receptor in vivo and prevents DC induction of antigen-restricted Th2 cells as we and others have recently shown (Hauben et al., 2008; Connolly et al., 2011). We confirmed that intrapancreatic CD4⁺ T cells are prevented from Th2 deviation after treatment with VAG539 (Fig. 4 E). Consistent with our expectation, VAG539 markedly protected against pancreatic inflammation (Fig. 4 F).

DCs expand in chronic pancreatitis and pancreatic adenocarcinoma

To further investigate the role of DCs in chronic pancreatic inflammation and malignant transformation, we examined the prevalence of DCs in pancreatic cancer and chronic pancreatitis. The fraction and total number of DCs were markedly increased in the pancreata of p48Cre;*Kras*^{G12D} mice compared with age-matched WT controls (Fig. 5, A–C). The surface phenotype of DCs infiltrating p48Cre;*Kras*^{G12D} pancreata differed from controls in that they contained a large B220⁺ plasmacytoid subset (Fig. 5 D) and an increased fraction of CD8⁺CD11⁻ lymphoid DCs (Fig. 5 E) and were highly mature, expressing elevated CD40 and CD86 (Fig. 5 E). To determine whether an intense DC infiltrate was similarly associated with human pancreatic cancer, we tested 19 human pancreatic carcinoma specimens for an array of DC markers. Consistent with our murine findings, there was a robust infiltrate of CD123⁺ cells, characteristic of plasmacytoid DCs, in all specimens tested (Fig. 5 F). DC-SIGN⁺ and CD1a⁺ cells were also present but in smaller numbers (Fig. 5 F).

To investigate whether DC expansion within the pancreas was a function of neoplasia or similarly associated with benign chronic pancreatic inflammation, we examined the prevalence of DCs, outside of the context of oncogenic *Kras*, in caerulein-induced chronic pancreatitis in WT mice. We found that DCs accounted for 1–3% of CD45⁺ leukocytes in normal pancreata but increased to 10–15% during chronic pancreatitis induction (Fig. 5 G). Moreover, the absolute number of pancreatic DCs increased 100-fold (Fig. 5 H). Conversely, the fraction of DCs in the spleen remained unchanged, implying a pancreas-specific expansion in chronic pancreatitis (Fig. 5 G). Collectively, these data show that a large DC infiltrate is a common feature in neoplastic and inflammatory pancreatic disease.

DCs exacerbate pancreatic fibroinflammation and ductal transformation

To directly test DC capacity to potentiate intrapancreatic inflammation via Th2 cells, we further expanded the pancreatic DC population by i.p. adoptive transfer of bone marrow-derived DCs to mice developing chronic pancreatitis. Creusot et al. (2009) reported that the plurality of bone marrow-derived




Figure 4. MyD88 blockade within DCs exacerbates pancreatic disease in a CD4⁺ T cell-dependent manner. (A) 4-wk-old CD11c-Cre MyD88 Floxed^{+/+} mice and control animals were treated with caerulein (C) for 3 wk to induce chronic pancreatitis. Pancreata were examined by Trichrome staining, and the fibroinflammatory area was quantified ($n = 4$ mice/group; ***, $P < 0.001$). (B) p48Cre;Kras^{G12D} mice were made chimeric using bone marrow derived from WT or CD11c-Cre MyD88 Floxed^{+/+} mice. Chimeric mice were treated with caerulein and sacrificed at 3 wk ($n = 4$ –5/group). Representative images are shown, and the fraction of graded PanIN lesions was quantified (***, $P < 0.001$). (C) CD4⁺ T cells were harvested from the pancreata of p48Cre;Kras^{G12D} mice and caerulein-treated WT mice that were administered MyD88 inhibitory peptide or control peptide. IL-2 and IL-4 levels were measured in cell culture supernatant (***, $P < 0.001$). (D) 4-wk-old p48Cre;Kras^{G12D} mice were treated with C, C + MyD88 inhibitor, C + GK1.5 to deplete CD4 T cells, C + MyD88 inhibitor + GK1.5, or additional controls. Mice were sacrificed at 3 wk, and the foci of invasive cancer were quantified by examining 10 HPFs per mouse ($n = 5$ mice/group; ***, $P < 0.001$). (E) IL-10 was measured in cell culture supernatant from purified pancreatic CD4⁺ T cells from mice treated for 2 d with saline, VAG539, caerulein, or caerulein + VAG539 (***, $P < 0.001$). (F) WT mice were treated with caerulein alone or caerulein + VAG539 for 3 wk. Representative H&E-stained sections are shown ($n = 4$ mice/group), and fibroinflammatory changes were quantified (***, $P < 0.001$). Error bars indicate standard error of the mean.





Figure 5. DCs expand in pancreatic carcinoma and chronic pancreatitis. (A) Pancreata from p48Cre;Kras^{G12D} and WT mice were examined by immunofluorescence using anti-CD11c. (B and C) In addition, the fraction of both CD45⁺ and CD11c⁺ leukocytes (B) and the total number of DCs (C) in pancreata of 9-mo-old p48-Cre;Kras^{G12D} and WT mice were determined by flow cytometry (**, $P < 0.01$). (D and E) DCs from the pancreata of WT and p48-Cre;Kras^{G12D} mice were further examined for B220 (D) and additional surface marker expression (E). Median fluorescence is shown below respective histograms. Experiments are representative of those repeated more than three times using a mean of three mice per group. (F) Frozen sections of human pancreatic cancer specimens were examined by immunofluorescence (CD123) and immunohistochemistry (DC-SIGN and CD1a). Insets show higher magnification. (G) The fraction of DCs among all CD45⁺ leukocytes in both the pancreas and the spleen was measured at timed intervals during the 3-wk course of caerulein administration in WT mice (mean of four mice per data point). (H) The total number of DCs per pancreas was measured on day 21 of saline or caerulein administration in WT mice (**, $P < 0.01$). Error bars indicate standard error of the mean.

DCs migrate to the peripancreatic tissues within 24 h after adoptive transfer and remain there for at least 12 d. After 3 wk of treatment with saline (Ctl), caerulein (C), DCs, or caerulein and DCs (C+DCs), pancreata were harvested and assessed for extent of fibrosis, inflammation, and endocrine and exocrine destruction.

Mice treated with C+DCs developed distortion of their pancreatic architecture including the formation of thick fibrous bands between lobules and effacement of the intra-lobular space with inflammatory fibroplasia (Fig. 6, A–C).

On average, >50% of the pancreatic surface area was replaced by fibroplasia in C+DC-treated mice (Fig. 6, A–C). This resulted in destruction of both the exocrine and endocrine pancreas, including a >60% reduction in acinar cell volume (Fig. 6, D and E), and a marked reduction in islet cell mass (Fig. 6 F). In consort with the latter findings, mice developed overt diabetes by 3 wk of C+DC challenge (Fig. 6 G). The severity of pancreatic exocrine destruction was exemplified by a roughly fourfold diminution in pancreatic-specific serum lipase levels in C+DC-treated mice (not depicted). Furthermore, DC



transfer in chronic pancreatitis resulted in pancreatic stellate cell activation *in vivo* as indicated by increased expression of peri-acinar desmin (Fig. 6, H and I) and α -SMA (Fig. 6, J and K) in pancreata of C+DC-treated mice. Pancreatic stellate cell activation is necessary for pancreatic fibrosis and neoplastic progression (Omary et al., 2007; Masamune et al., 2009). In addition, overexpansion of DCs in the context of pancreatitis resulted in the recruitment of a robust immune infiltrate.

Figure 6. DCs exacerbate pancreatic fibroinflammation. WT mice were treated with saline, C alone, DCs alone, or C+DCs. (A–C) Paraffin-embedded sections of pancreata were stained with H&E (A) and Picric acid–Sirius red (B), and the fibroinflammatory area per mouse pancreas was calculated for all treatment groups (C). Insets show higher magnification. (D–G) Pancreata were also stained using mAbs directed against amylase (D and E) and insulin (F), and serum glucose (G) levels were calculated. (H–L) Similarly, pancreata were stained using mAbs directed against desmin (H and I), α -SMA (J and K), and CD3 (L). (M) Using flow cytometry, the relative number of intra-pancreatic B cells, macrophages, neutrophils, and NK cells in chronic pancreatitis in the context of exogenous DCs expansion was calculated ($n = 8$ –10 mice/group; *, $P < 0.05$; ***, $P < 0.001$). Error bars indicate standard error of the mean.

In particular, there was an increased T cell infiltrate (Fig. 6 L) as well as a substantial increase in B cell, macrophage, neutrophil, and NK cell populations (Fig. 6 M). The effects of C+DC treatment were pancreas specific as there was no effacement of the kidney or liver parenchyma or associated immune infiltrate in these organs (not depicted).

To confirm the consistency of our findings, we tested whether DCs also exacerbated pancreatitis in a model of endogenous DC expansion using Flt3L (Fms-like tyrosine kinase-3 ligand; Maraskovsky et al., 1996). Mice treated with Flt3L for 10 d developed expansion of pancreatic DC populations between days 4 and 15 (Fig. 7 A). Inflammatory monocyte populations were not significantly expanded (Fig. 7 B). Moreover, treatment with Flt3L and simultaneous challenge with caerulein resulted in exacerbated acinar destruction, fibrosis, and inflammation (Fig. 7, C and D), suggesting that the effects of DC expansion in pancreatitis are not model specific.

In addition to the fibroinflammatory changes, a conspicuous phenotypic finding observed in pancreatitis associated with DC transfer is the widespread development of early PanIN lesions. Overall, $\sim 50\%$ of ducts were classified as PanINs in C+DC pancreata (Fig. 8, A and B). Affected ducts in C+DC-treated mice stained positively for Alcian blue (Fig. 8 C). Pancreata of control animals, including mice receiving DCs alone or caerulein alone, did not exhibit PanIN lesions after 3 wk of treatment. Furthermore, pancreata from C+DC-treated animals had a markedly high fraction of Ki67⁺ proliferating epithelial cells compared with controls (Fig. 8 D).

DCs accelerate the growth of pancreatic tumors

Prolonged duration of C+DC administration to WT mice for 3 mo resulted in greater fibrodysplastic obliteration of the pancreatic parenchyma and more diffuse PanIN lesions but not invasive carcinoma (Fig. 8 E). However, DCs markedly accelerated malignant transformation when transferred for




Figure 7. Fit3L treatment exacerbates pancreatitis.

(A) The time course of DC expansion in the pancreas of Fit3L-treated WT mice is shown. Both the total number of pancreatic DCs and their fraction among pancreatic leukocytes are indicated. Data are based on a mean of four mice per time point. (B) The fraction of inflammatory monocytes in the pancreata of Fit3L-treated mice and controls was calculated. Day 15 data are shown. (C and D) Mice treated with Fit3L alone, caerulein alone, or caerulein + Fit3L were stained with H&E, Gomori's Trichrome, and anti-CD45. (D) Fibroinflammatory area was quantified by examining 10 HPFs per mouse (five mice per group; ***, $P < 0.001$). Error bars indicate standard error of the mean.

transferred DCs from MyD88^{-/-} mice to WT mice undergoing caerulein pancreatitis. As predicted, MyD88^{-/-} DC transfer worsened pancreatitis (Fig. 9 A) and PanIN formation (Fig. 9 B) to a greater extent than WT DC transfer. Endocrine and exocrine pancreata were almost entirely replaced by collagen after MyD88^{-/-} DC transfer, and >70% of ducts were PanINs. In addition, transfer of MyD88^{-/-} DCs resulted in an approximately twofold increase in CD4⁺ T cell recruitment compared with transfer of WT DCs (Fig. 9 C). Collectively, these data support our findings using CD11c-Cre MyD88 Floxed^{+/+} mice (Fig. 4, A–D) and together suggest that DCs are sufficient to induce the exacerbated inflammation and dysplasia observed in the MyD88-deficient pancreata.

The DC–Th2 axis mediates exacerbated pancreatitis and transformation

Consistent with our hypothesis that suggests that the DC–CD4 axis can mediate pancreatic inflammation and transformation, we found that CD4⁺ T cell-deficient mice were protected from the effects of DC transfer in chronic pancreatitis (Fig. 9, D and E). Conversely, the absence of CD8⁺ T cells, B cells, neutrophils, and monocyte depletion or TNF blockade did not protect C+DC-treated mice (Fig. 9, D and E). Furthermore, in our pancreatic cancer model, CD4⁺ T cell depletion protected p48Cre;Kras^{G12D} mice adoptively transferred with DCs from developing accelerated carcinogenesis (Fig. 9 F). CD4⁺ T cell depletion had similarly protective effects after transfer of MyD88^{-/-} DCs (not depicted).

In line with the observations that CD4⁺ T cells are the critical effectors in DC-mediated pancreatic inflammation and transformation, we found that the number of intrapancreatic CD4⁺ T cells and CD4/CD8 T cell ratio were markedly increased in mice adoptively transferred with DCs (Fig. 10, A and B). Furthermore, intrapancreatic CD4⁺ T cells in DC-treated mice exhibited a strong Th2 differentiation (Fig. 10 C). These effects were accentuated by transfer of MyD88^{-/-} DCs (Fig. 10, A–C). In addition to the significant lack of intrapancreatic Th1 and Th17 differentiation, the fraction of intrapancreatic CD4⁺CD25⁺Foxp3⁺ regulatory T cells (T_{reg} cells) was

4 wk to p48Cre;Kras^{G12D} mice (Fig. 8, F and G). Furthermore, protein analysis revealed that DC transfer resulted in altered pancreatic expression of numerous cell cycle regulatory and tumor suppressor genes including up-regulated expression of p21, p27, p-p27, p53, and Rb (Fig. 8 H). Moreover, consistent with our previous data (Fig. 4 B), adoptive transfer of DCs derived from MyD88^{-/-} mice resulted in significantly more accelerated tumorigenesis (Fig. 8, F and G). We further tested the protumorigenic capacity of DCs in an orthotopic model. Cohorts of WT mice were challenged with intrapancreatic Kras^{G12D} PDECs and then sacrificed 6 wk later. In parallel, selected mice received thrice weekly DC adoptive transfer beginning in week 2. DCs again induced a threefold accelerated growth rate of pancreatic lesions (Fig. 8 I). The relative distribution of epithelia and stroma was similar in both the control and DC adoptive transfer groups (Fig. 8 J).

DCs mediate effects of MyD88 inhibition in pancreatic disease

To further investigate the augmented proinflammatory and transformative effects of MyD88 inhibition in DCs, we




Figure 8. DCs promote the transition from chronic pancreatitis to ductal dysplasia and accelerate the growth rate of pancreatic tumors.

(A and B) 4-wk-old WT mice were treated with C or C+DCs for 3 wk. Representative H&E-stained section from C+DC-treated pancreata is shown. The fraction of pancreatic ducts in C or C+DC mice exhibiting PanIN lesions was quantified by examining pancreata from >15 mice per group. (C) Representative paraffin-embedded section of pancreata from mice treated with C+DCs and stained with Alcian blue is shown. (D) Pancreata from mice treated with C or C+DCs were stained for CK19 (pink) and Ki67 (brown), and the fraction of Ki67⁺ ductal cells was quantified for each treatment group. (E) Representative H&E-stained image of pancreas from a mouse treated for 12 wk with C+DCs. (F and G) 4-wk-old p48Cre;Kras^{G12D} mice were adoptively transferred with DCs derived from WT or MyD88^{-/-} mice thrice weekly for 5 wk before sacrifice ($n = 5$ /group). Representative H&E-stained sections are shown, pancreata were weighed, and the fraction of PanIN 2/3 ducts per HPF was determined (G). (H) Lysate from pancreata of mice adoptively transferred with WT DCs or saline was tested for expression of p21, p27, p-p27, p53, and Rb by Western blotting. (I) Mice were challenged with 3×10^6 (experiment 1) or 10^6 (experiment 2) KRas^{G12D} PDECs. In each experiment, selected mice were adoptively transferred with DCs (10^6) thrice weekly from weeks 2–6. Mice were sacrificed after 6 wk, and pancreatic lesions were weighed. (J) Representative Trichrome-stained paraffin-embedded sections are shown, and the fraction of fibrotic pancreatic area was quantified (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$). Error bars indicate standard error of the mean.

also markedly reduced in C+DC-treated animals (Fig. 10 D). Similarly, MHC II^{-/-} mice, which are deficient in Th2 signaling, were protected from the effects of DC expansion in pancreatitis, whereas β 2-microglobulin^{-/-} mice deficient in CD8⁺ T cell signaling were not protected (Fig. 10, E and F). Collectively, our data suggest that DC expansion in the injured

pancreas increases CD4⁺ T cell Th2 differentiation, which is required for the organ destructive, fibrotic, inflammatory, and dysplastic effects, and decreases the fraction of other T cell subsets, including T_{reg} cells and CD8⁺ cells.

We postulated that DCs induce Th2-deviated CD4⁺ T cells by capturing antigen from the injured pancreas, which




Figure 9. Fibroinflammatory and dysplastic effects of DCs are exaggerated upon MyD88 inhibition and are contingent on CD4⁺ T cells.

(A–C) MyD88^{-/-} or WT DCs were adoptively transferred to WT mice undergoing caerulein-induced chronic pancreatitis. The extent of fibrosis and acinar destruction (A), as well as PanIN formation (B) and size of the CD4⁺ T cell infiltrate (C) in pancreata of treated animals were determined. Experiments were performed using a mean of five mice per group (**, $P < 0.01$; ***, $P < 0.001$). (D and E) Mice deficient in CD4⁺ T cells, CD8⁺ T cells, B cells, Gr1⁺ inflammatory monocytes and neutrophils, or TNF production were induced to develop chronic pancreatitis in the context of DC overexpansion. Pancreatic effacement and fibroinflammatory changes were quantified for each group by examining 10 HPFs per slide ($n = 4$ –6 mice/experimental group; ***, $P < 0.001$). (F) 4-wk-old p48Cre;Kras^{G12D} mice were adoptively transferred with DCs or administered saline for 5 wk. Selected cohorts were additionally treated with GK1.5 to deplete CD4⁺ T cells. Pancreas weights were measured ($n = 4$ mice/experimental group; ***, $P < 0.001$). Error bars indicate standard error of the mean.

they present to naive T cells, thereby generating antigen-restricted effector cells. To test whether these effectors are specific to pancreatic antigen, CD4⁺ T cells from the pancreata of C+DC-treated WT mice were purified and cultured for 5 d with either mock-loaded DCs or DCs loaded with lysate from digested pancreata. CD4⁺ T cells co-cultured with pancreatic lysate-pulsed DCs exhibited increased activation compared with controls, implying pancreas-directed specificity (Fig. 10 G). Furthermore, transfer of CD4⁺ T cells from C+DC-treated WT mice to p48Cre;Kras^{G12D} mice markedly

accelerated pancreatic cancer development, whereas CD4⁺ T cell transfer from control mice did not have tumor-promoting effects (Fig. 10, H and I).

DISCUSSION

Pancreatic cancer is a devastating disease with very few effective treatment options as <5% of patients achieve cure (Vincent et al., 2011). Recent evidence suggests that rather than being a dormant component, pancreatic fibroinflammatory stroma impacts malignant transformation and disease progression.




Figure 10. DCs exacerbate pancreatic disease by inducing antigen-restricted Th2 cells. (A and B) The total number of CD4⁺ cells in the pancreata of mice treated with C+DC^{WT}, C+DC^{MyD88-/-}, and controls (A) and the fraction of intrapancreatic CD4⁺ or CD8⁺ T cells among all CD3⁺ cells (B) were measured by flow cytometry (***, $P < 0.001$). (C) CD4⁺ T cell differentiation in the pancreas of mice treated with C+DC^{WT}, C+DC^{MyD88-/-}, and controls was determined by measuring their production of Th1, Th2, and Th17 cytokines after FACS sorting. (D) The fraction of intrapancreatic T_{reg} cells was determined by gating on CD45⁺CD25⁺ leukocytes and cross staining for CD4 and Foxp3. In vitro assays were repeated three times with similar results. (E and F) MHC II^{-/-} and β 2-microglobulin^{-/-} mice treated with caerulein were recipients of DC transfer. A mean of four mice per group was used in these experiments (***, $P < 0.001$). (G) To determine the pancreatic antigen specificity of effector T cells, FACS-sorted pancreas-infiltrating CD4⁺ T cells from WT C+DC-treated mice were cultured with pancreatic lysate- or mock-pulsed DCs. CD4⁺ T cell activation was then determined by their expression of CD69, FAS ligand, and CD11b. Gray histograms represent isotype control. Median fluorescent indexes are shown for each group. Experiments were repeated three times with similar results. (H and I) p48Cre;Kras^{G12D} mice were adoptively transferred for 5 wk with CD4⁺ T cells derived from control and C+DC-treated mice ($n = 3$ /group). Representative H&E-stained sections and images of pancreata are shown (H), and the fraction of pancreatic area with preserved acinar architecture was measured (I; ***, $P < 0.001$). Error bars indicate standard error of the mean.

In particular, by releasing nutrient growth factors into the neoscient tumor microenvironment, such as insulin-like growth factor and PDGF, the stromal component of pancreatic cancer has been closely linked to carcinogenesis as well as tumor growth and invasiveness (Wang et al., 2007; Mantovani et al., 2010). We postulated that TLR4 may regulate stromal inflammation as it has previously been implicated in benign pancreatic disease (Sharif et al., 2009). Our related findings that TLR4 is increased in expression on leukocytes within the tumor microenvironment in humans and mice, that TLR4 ligands such as HMGB-1 and S100A9 are rife within the tumor microenvironment, and that TLR4 ligation accelerates carcinogenesis in mice lends insight into the mechanism of regulation of pancreatic tumor progression by the inflammatory stroma. We further show that TLR4 signaling via the MyD88-independent TRIF pathway modulates pancreatic carcinogenesis as targeting TLR4 or TRIF prevents cancer progression in at risk pancreata. These data suggest that testing clinical-grade TLR4 or TRIF inhibitors in pancreatic cancer would be appropriate (Barrat and Coffman, 2008). Our findings also suggest that there may possibly be a role for endogenous LPS derived from gut bacteria in modulating pancreatic carcinogenesis. Indeed, a role for the microbiome in carcinogenesis is rapidly emerging in colon cancer and other malignancies (Ahn et al., 2012; Dapito et al., 2012), and our data suggest mechanistic plausibility for its possible influence on pancreatic oncogenesis. However, definitive linkage of pancreatic cancer development to the host microbiome via TLR4 requires more exact investigation.

An equally compelling discovery in this study is that MyD88 blockade vigorously accelerates pancreatic fibroinflammation as well as carcinogenesis. Most notably, MyD88 blockade resulted in invasive pancreatic adenocarcinoma within 3 wk in p48Cre;Kras^{G12D} mice treated with caerulein compared with the development of early PanIN lesions in age-matched mice with intact MyD88 signaling. Moreover, malignant epithelial cells in MyD88-deficient mice exhibited altered expression of p53 in the absence of an endogenous p53 mutation. Overexpression of p53 in pancreatic cancer has been linked to peritumoral inflammation and is associated with an aggressive oncogenic phenotype (Carrière et al., 2011). Our findings contrast with studies linking MyD88 to inflammation and carcinogenesis (Gasse et al., 2007; Rakoff-Nahoum and Medzhitov, 2007; Swann et al., 2008; Coste et al., 2010). For example, Rakoff-Nahoum and Medzhitov (2007) crossed *APC*^{min/+} mice, which carry a germline mutation in the *Apc* tumor suppressor gene, with MyD88-deficient mice and found that the absence of MyD88 signaling resulted in a decreased number of intestinal polyps and substantially improved survival. Recently, Coste et al. (2010) also showed that MyD88 plays a cell-autonomous role in Ras-mediated transformation in mouse and human tissues. The proinflammatory and carcinogenic effects of MyD88 inhibition in the pancreas are unlikely related to unremitting TRIF signaling as combined TRIF and MyD88 blockade nonetheless resulted in severe pancreatic disease. Rather, the resolution to our ostensibly paradoxical finding lies in the fact that in the presence of MyD88 blockade,

DCs preferentially induce Th2 polarization (Kaisho et al., 2002; Kapsenberg, 2003; Chen et al., 2010), which, in pancreatic disease, compensates for the inability to produce generalized inflammation via NF- κ B (Fig. S1). We demonstrate that DCs are the causal cellular component responsible for exacerbated chronic pancreatitis and neoplasia upon MyD88 blockade by using DC adoptive transfer experiments using MyD88^{-/-} DCs and by examining CD11c-Cre MyD88 Floxed^{+/+} mice, which are deficient in MyD88 signaling exclusively within CD11c⁺ cells.

The observation that DCs exacerbate pancreatic pathology through the activation and expansion of pancreas-restricted Th2-deviated CD4⁺ T cell population was supported by the protection offered in MHC II^{-/-} mice compared with β 2-microglobulin^{-/-} mice, our finding that CD4⁺ T cells derived from C+DC pancreata could accelerate tumorigenesis when transferred to p48Cre;Kras^{G12D} animals, and that depletion of Th2-deviated CD4⁺ T cells protects against the proinflammatory and tumorigenic effects of DC adoptive transfer or MyD88 blockade. Furthermore, given the intrapancreatic expansion of Th2 cells upon DC transfer, and suppression of CD8⁺, Th1, Th17, and T_{reg} cell differentiation, and our observation that these effector T cells recognize pancreatic antigen, a logical hypothesis governing the DC–Th2 axis in pancreatic carcinogenesis is that infiltrating DCs capture pancreatic antigen and induce antigen-restricted Th2 CD4⁺ T cell differentiation, which then serve as the effector cell mediating cellular injury and inflammation required for stromal activation and neoplastic transition. Our discovery of the importance of Th2 polarization in pancreatic disease corresponds with recent studies that have shown an association between Th2-mediated inflammation and neoplastic development in patients with melanoma and in experimental models of colon carcinogenesis (Osawa et al., 2006; Nevala et al., 2009). Human pancreatic cancer is also marked by an extensive peritumoral Th2 infiltration, which is associated with reduced survival (De Monte et al., 2011). The current data suggest that DCs are sufficient to induce accelerated pancreatic tumorigenesis via induction of pancreatic antigen-restricted Th2-deviated CD4⁺ T cells. However, it is notable that CD4⁺ T cell depletion did not rescue p48Cre;Kras^{G12D} mice outside of the context of MyD88 blockade (Fig. 4 D), suggesting that alternate protumorigenic mechanisms, rather than DC–CD4⁺ T cell activity, may be primary in pancreata of p48Cre;Kras^{G12D} mice with intact MyD88 signaling pathways.

The paradigm linking inflammation and inflammatory cells to carcinogenesis has been gaining consistent momentum in the scientific literature (Colotta et al., 2009; Mantovani et al., 2010). A heightened inflammatory milieu provides a continuous supply of cytokines and growth factors that can affect biological processes responsible for maintaining cellular homeostasis, leading to genetic instability and an increased risk of carcinogenesis (Greer and Whitcomb, 2009). Our experiments revealed a robust, pancreas-specific expansion of DC populations in animals subject to both benign and malignant pancreatic disease and in human pancreatic cancer tissue.

We show that DC expansion within the inflamed pancreas leads to the formation of early PanIN lesions with high proliferative rates. Furthermore, DCs accelerate pancreatic tumor growth and are sufficient to induce altered expression of numerous cell cycle regulatory genes and tumor suppressor genes. DCs have been considered a potent tool in nature's defense against carcinogenesis and tumor progression as DCs avidly capture tumor antigen *in vivo* and initiate both adaptive and innate antitumor immune responses (O'Neill et al., 2004; Petersen et al., 2010). Conversely, we describe a deleterious role for DCs as potent protumorigenic agents in the pancreatic stroma. The concept of DCs as protumorigenic is consistent with recent studies showing that thymic stromal lymphopoietin engenders the growth of human breast cancer by promoting Th2 inflammation via OX40L⁺ DCs (Ito et al., 2005; Liu et al., 2011). This protumorigenic capacity of DCs might explain, in part, the reason DC immunotherapy and vaccine regimens have shown limited clinical efficacy in the treatment of human cancers (Palucka et al., 2009; Robson et al., 2010). Moreover, this work could provide a rationale for experimental therapeutic strategies aimed at targeting DCs in individuals with chronic pancreatitis who are at greatest risk for transformation.

The current study must be viewed in the context of our very recent study, which found that DCs have a parallel protective role in acute pancreatic inflammation (Bedrosian et al., 2011). That is, we found that DC depletion in acute pancreatitis, rather than mitigating injury, results in complete exocrine organ necrosis. We showed that the mechanism for the pancreatic destruction upon DC depletion may be related to their critical role in the clearance of apoptotic cells and necrotic debris during the regeneration phase of pancreatitis. Conversely, in the context of DC depletion, there is a severe deficiency in the clearance of cellular debris and a parallel rise in sterile inflammation within the pancreas, resulting in acinar organ necrosis with sparing of the islets (Bedrosian et al., 2011). Thus, within the same organ and in one disease, our studies show that DCs can have dual opposing effects. These are the first findings of parallel pro-fibroinflammatory and dysplastic as well as protective roles for DCs within the context of organ-specific inflammation and may have implications to understanding the complex role of DCs in other immune-based diseases.

MATERIALS AND METHODS

Animals and procedures. Male C57BL/6 (H-2Kb) were purchased from Taconic and bred in-house. Transgenic mice deficient in CD4 T cells (B6.129S2-*Cd4^{m1Mak}/J*), CD8 T cells (B6.129S2-*Cd8a^{m1Mak}/J*), B cells (B6.129S2-*Igh-6^{m1Cgn}/J*), MHC II (B6.129S2-*C2ta^{m1Cam}/J*), β 2-microglobulin (B6.129P2-B2m^{tm1Unc}/J), TNF (B6.129S-*Tnf^{tm1Ckl}/J*), TRIF (C57BL/6J-*Ticam1^{ly2}/J*), and TLR4 (B6.B10ScN-*Tlr4^{lys-del}/Jth*) were purchased from the Jackson Laboratory. p48Cre;Kras^{G12D} mice, which develop pancreatic neoplasia endogenously by expressing a single mutant *Kras* allele in progenitor cells of the pancreas (gift of D. Tuveson, Cambridge Research Institute, Cambridge, England, UK), were generated by crossing LSL-Kras^{G12D} mice with p48-Cre mice, which express Cre recombinase from a pancreatic progenitor-specific promoter (Hingorani et al., 2003). MyD88^{-/-} mice were a gift of D. Levy (New York University School of Medicine, New York, NY). CD11c-Cre

MyD88 Floxed^{+/+} mice, which are deficient in MyD88 exclusively within CD11c⁺ cells, were generated by breeding B6.129P2(SJL)-*Myd88^{tm1Dof}/J* mice with B6.Cg-Tg(Itgax-cre)1-1Reiz/J (The Jackson Laboratory) as reported previously (Hou et al., 2008). Bone marrow chimeric animals were created by irradiating 2-mo-old mice (100 Gy) followed by *i.v.* bone marrow transfer (10⁷ cells) from nonirradiated donors as we have described previously (Bedrosian et al., 2011). Chimeric mice were used in experiments 7 wk later. To accelerate carcinogenesis in p48Cre;Kras^{G12D} mice, two doses of caerulein (50 μ g/kg; Sigma-Aldrich) were administered over 48 h as described previously with modifications, and mice were sacrificed at 3 wk (Carrière et al., 2009). Animals were housed in a clean vivarium and fed standard mouse chow. Animal procedures were approved by the New York University School of Medicine Institutional Animal Care and Use Committee.

Cellular isolation, culture, and analysis. Bone marrow-derived DCs were generated as described previously (Miller et al., 2002). In brief, bone marrow aspirates were cultured for 8 d in complete RPMI supplemented with 20 ng/ml murine GM-CSF. Pancreatic and splenic mononuclear cells were isolated by mechanical and chemical digestion using Collagenase IV (Sigma-Aldrich). For selected experiments, CD4⁺ T cells were purified from suspensions of pancreatic mononuclear cells using the MoFlo Cell Sorter (Beckman Coulter) and cultured with plate-bound anti-CD3. Cell culture supernatant was assayed in a cytometric bead array (BD).

In vivo models. Chronic pancreatitis was induced using seven consecutive hourly *i.p.* injections of caerulein (50 μ g/kg; Sigma-Aldrich) thrice weekly for 3–12 wk. In animals adoptively transferred with DCs, 10⁶ bone marrow-derived DCs were administered *i.p.* thrice weekly. In selected animals, 10 μ g Flt3L was administered *i.p.* for 10 d. In parallel, caerulein was administered thrice weekly for 2 wk starting on day 4 of Flt3L administration. For CD4⁺ T cell adoptive transfer experiments, 2 \times 10⁵ pancreatic CD4⁺ T cells were purified and transferred administered *i.p.* thrice weekly for 5 wk. To establish orthotopic pancreatic lesions, we grafted PDECs harboring oncogenic KRas^{G12D} by direct intrapancreatic injection via laparotomy as we have described previously (Agbunag et al., 2006; Pylayeva-Gupta et al., 2012). Mice were sacrificed 6 wk after PDEC transfer. To deplete Gr1⁺ cells or CD4⁺ T cells, respectively, RB6-8C5 or GK1.5 was used as described previously (Connolly et al., 2010; Monoclonal Antibody Core Facility, Memorial Sloan-Kettering Cancer Center). MyD88 inhibitory peptide or control peptide (200 μ g; InvivoGen) was administered *i.p.* thrice weekly in selected experiments. NF- κ B blockade was accomplished using the cell-permeable NEMO binding domain inhibitor (1 mg/kg/day; EMD Millipore). MAP kinase blockade was accomplished using PD98059 (2.5 mg/kg/day; InvivoGen). TLR4 and TRIF blockade was accomplished using a TLR4 inhibitory peptide (IMG-2011A; 100 μ g thrice weekly; Imgenex) or an inhibitor of the TRIF adaptor protein (Pepinh-TRIF; 150 μ g thrice weekly; InvivoGen), respectively. In selected experiments, mice were treated with 5 μ g TLR4 ligand LPS or 80 μ g TLR3 ligand Poly I:C thrice weekly (both InvivoGen). The novel immune modulator VAG539 (30 mg/kg/day; Novartis) was administered via oral gavage (Hellerbrand et al., 1996; Creusot et al., 2009). Serum levels of glucose and lipase were measured using an AU400 Chemistry Analyzer (Olympus).

Western blotting. For immunoblotting assays, cell lysates were prepared from whole pancreata or Raji lymphoma cells. For our human experiments, proteins were also isolated from human pancreatic duct fluid harvested at surgery from patients undergoing pancreatic resection. Total protein was determined by the Lowry assay, and lysates were equilibrated onto a 10% polyacrylamide gel (30% acrylamide/Bis solution, 37.5:1, 2.6% C). Proteins were then electrotransferred to a polyvinylidene difluoride membrane at 90 V for 90 min. Subsequently, the polyvinylidene difluoride was immunoblotted using antibodies directed against pIRF3 (Cell Signaling Technology), HMGB-1, S100A9, IRAK, pIRAK, p21, p27, p-p27, p53, Rb, and β -actin (Santa Cruz Biotechnology, Inc.). Levels of TLR4 agonists in pancreatic duct fluid were quantified on HEK-Blue cells (InvivoGen) using fixed quantities of TLR4 ligand (LPS) as controls (Guo et al., 2009).

Flow cytometry. Cell surface marker analysis was performed by flow cytometry using the FACSCalibur (Beckman Coulter) after incubating 5×10^5 cells with 1 μ g anti-Fc γ RIII/II antibody (2.4G2, Fc block; Monoclonal Antibody Core Facility, Memorial Sloan-Kettering Cancer Center) and then labeling with 1 μ g FITC-, PE-, PerCP-, or APC-conjugated antibodies directed against MHC II (I-A^b), B220 (RA3-6B2), CD3 ϵ (17A2), CD4 (RM4-5) CD8 α (53-6.7), CD11b (M1/70), CD11c (HL3), CD25 (PC61.5), CD40 (HM40-3), CD45.2 (104), CD69 (H1.2F3), CD86 (GL1), F/480 (BM8), Foxp3 (FJK-16s), FAS Ligand (MFL3), NK1.1 (PK136), and Gr1 (RB6-8C5; all BD, eBioscience, or BioLegend). Pancreatic mononuclear cells were also stained for TLR4 (Imgenex). Dead cells were excluded by staining with 7-amino-actinomycin D (BD).

Histology, immunohistochemistry, and microscopy. For histological analysis, paraffin-embedded or frozen sections were stained with hematoxylin and eosin (H&E), Gomori's Trichrome, Alcian blue, or Picric acid–Sirius red. In addition, immunohistochemistry was performed using antibodies directed against CD3 (Invitrogen), CD4 (Abbtotec), p53 (Novocastra), CK19 (Developmental Studies Hybridoma Bank), Desmin (Sigma–Aldrich), α -SMA (Novus Biologicals), TLR4 (Imgenex), amylase (Sigma–Aldrich), CD11c, DC-SIGN, CD1a (BD), CD123, and insulin (Abcam). Photographs were taken using a DM2M microscope (Leica) and a digital camera (Optronics). Fluorescence images were captured on an Axiovert 200M fluorescence microscope (Carl Zeiss). Ductal dysplasia was identified and graded in a blinded manner according to established criteria (Hingorani et al., 2003; Hruban et al., 2004). In brief, in PanIN I lesions, the normal cuboidal PDECs transition to columnar architecture. PanIN II lesions are associated with additional nuclear abnormalities such as loss of polarity. PanIN III lesions, or in situ carcinoma, show cribriforming, budding off of cells, and luminal necrosis with marked cytological abnormalities, without invasion beyond the basement membrane.

Statistics. Data are presented as mean \pm standard error of mean. Statistical significance was determined by the Student's *t* test or the log-rank test using Prism 4 (GraphPad Software). *P*-values < 0.05 were considered significant.

Online supplemental material. Fig. S1 shows divergent effects of modulating MyD88 and TRIF signaling on pancreatic carcinogenesis. Online supplemental material is available at <http://www.jem.org/cgi/content/full/jem.20111706/DC1>.

This work was supported in part by grants from the Society of University Surgeons (to G. Miller), the National Pancreas Foundation (to G. Miller), the Irvington Institute Postdoctoral Fellowship Program of the Cancer Research Institute (to Y. Pylayev-Gupta), and National Institutes of Health Awards CA108573 (to A.B. Frey), CA055360 (to D. Bar-Sagi), DK085278 (to G. Miller), and CA155649 (to G. Miller). The Flow Cytometry and Histology and Immunohistochemistry Core Facilities of the New York University School of Medicine are partially supported by the National Institutes of Health (grant 5 P30CA016087-31).

The authors have no conflicting financial interests.

Submitted: 12 August 2011

Accepted: 27 July 2012

REFERENCES

- Agbunag, C., K.E. Lee, S. Buontempo, and D. Bar-Sagi. 2006. Pancreatic duct epithelial cell isolation and cultivation in two-dimensional and three-dimensional culture systems. *Methods Enzymol.* 407:703–710. [http://dx.doi.org/10.1016/S0076-6879\(05\)07055-2](http://dx.doi.org/10.1016/S0076-6879(05)07055-2)
- Ahn, J., C.Y. Chen, and R.B. Hayes. 2012. Oral microbiome and oral and gastrointestinal cancer risk. *Cancer Causes Control.* 23:399–404. <http://dx.doi.org/10.1007/s10552-011-9892-7>
- Ammann, R.W., A. Akovbiantz, F. Largiader, and G. Schueler. 1984. Course and outcome of chronic pancreatitis. Longitudinal study of a mixed medical-surgical series of 245 patients. *Gastroenterology.* 86:820–828.
- Bantsimba-Malanda, C., J. Marchal-Sommé, D. Goven, O. Freynet, L. Michel, B. Crestani, and P. Soler. 2010. A role for dendritic cells in bleomycin-induced pulmonary fibrosis in mice? *Am. J. Respir. Crit. Care Med.* 182:385–395. <http://dx.doi.org/10.1164/rccm.200907-1164OC>
- Barrat, F.J., and R.L. Coffman. 2008. Development of TLR inhibitors for the treatment of autoimmune diseases. *Immunol. Rev.* 223:271–283. <http://dx.doi.org/10.1111/j.1600-065X.2008.00630.x>
- Bedrosian, A.S., A.H. Nguyen, M. Hackman, M.K. Connolly, A. Malhotra, J. Ibrahim, N.E. Cieza-Rubio, J.R. Henning, R. Barilla, A. Rehman, et al. 2011. Dendritic cells promote pancreatic viability in mice with acute pancreatitis. *Gastroenterology.* 141:1915–1926; e1–e14. <http://dx.doi.org/10.1053/j.gastro.2011.07.033>
- Braganza, J.M., S.H. Lee, R.F. McCloy, and M.J. McMahon. 2011. Chronic pancreatitis. *Lancet.* 377:1184–1197. [http://dx.doi.org/10.1016/S0140-6736\(10\)61852-1](http://dx.doi.org/10.1016/S0140-6736(10)61852-1)
- Carrière, C., A.L. Young, J.R. Gunn, D.S. Longnecker, and M. Korc. 2009. Acute pancreatitis markedly accelerates pancreatic cancer progression in mice expressing oncogenic Kras. *Biochem. Biophys. Res. Commun.* 382:561–565. <http://dx.doi.org/10.1016/j.bbrc.2009.03.068>
- Carrière, C., A.J. Gore, A.M. Norris, J.R. Gunn, A.L. Young, D.S. Longnecker, and M. Korc. 2011. Deletion of Rb accelerates pancreatic carcinogenesis by oncogenic Kras and impairs senescence in premalignant lesions. *Gastroenterology.* 141:1091–1101. <http://dx.doi.org/10.1053/j.gastro.2011.05.041>
- Chen, L., L. Lei, X. Chang, Z. Li, C. Lu, X. Zhang, Y. Wu, I.T. Yeh, and G. Zhong. 2010. Mice deficient in MyD88 Develop a Th2-dominant response and severe pathology in the upper genital tract following *Chlamydia muridarum* infection. *J. Immunol.* 184:2602–2610. <http://dx.doi.org/10.4049/jimmunol.0901593>
- Colotta, F., P. Allavena, A. Sica, C. Garlanda, and A. Mantovani. 2009. Cancer-related inflammation, the seventh hallmark of cancer: links to genetic instability. *Carcinogenesis.* 30:1073–1081. <http://dx.doi.org/10.1093/carcin/bgp127>
- Connolly, M.K., A.S. Bedrosian, J. Mallen-St Clair, A.P. Mitchell, J. Ibrahim, A. Stroud, H.L. Pachter, D. Bar-Sagi, A.B. Frey, and G. Miller. 2009. In liver fibrosis, dendritic cells govern hepatic inflammation in mice via TNF- α . *J. Clin. Invest.* 119:3213–3225.
- Connolly, M.K., J. Mallen-St Clair, A.S. Bedrosian, A. Malhotra, V. Vera, J. Ibrahim, J. Henning, H.L. Pachter, D. Bar-Sagi, A.B. Frey, and G. Miller. 2010. Distinct populations of metastases-enabling myeloid cells expand in the liver of mice harboring invasive and preinvasive intra-abdominal tumor. *J. Leukoc. Biol.* 87:713–725. <http://dx.doi.org/10.1189/jlb.0909607>
- Connolly, M.K., D. Ayo, A. Malhotra, M. Hackman, A.S. Bedrosian, J. Ibrahim, N.E. Cieza-Rubio, A.H. Nguyen, J.R. Henning, M. Dorvil-Castro, et al. 2011. Dendritic cell depletion exacerbates acetaminophen hepatotoxicity. *Hepatology.* 54:959–968. <http://dx.doi.org/10.1002/hep.24429>
- Coste, I., K. Le Corf, A. Kfoury, I. Hmitou, S. Druilennec, P. Hainaut, A. Eychene, S. Lebecque, and T. Renno. 2010. Dual function of MyD88 in RAS signaling and inflammation, leading to mouse and human cell transformation. *J. Clin. Invest.* 120:3663–3667. <http://dx.doi.org/10.1172/JCI42771>
- Creusot, R.J., S.S. Yaghoubi, P. Chang, J. Chia, C.H. Contag, S.S. Gambhir, and C.G. Fathman. 2009. Lymphoid-tissue-specific homing of bone-marrow-derived dendritic cells. *Blood.* 113:6638–6647. <http://dx.doi.org/10.1182/blood-2009-02-204321>
- Dapito, D.H., A. Mencin, G.Y. Gwak, J.P. Pradere, M.K. Jang, I. Mederacke, J.M. Caviglia, H. Khiabani, A. Adeyemi, R. Bataller, et al. 2012. Promotion of hepatocellular carcinoma by the intestinal microbiota and TLR4. *Cancer Cell.* 21:504–516. <http://dx.doi.org/10.1016/j.ccr.2012.02.007>
- De Monte, L., M. Reni, E. Tassi, D. Clavenna, I. Papa, H. Recalde, M. Braga, V. Di Carlo, C. Dogliani, and M.P. Protti. 2011. Intratumor T helper type 2 cell infiltrate correlates with cancer-associated fibroblast thymic stromal lymphopoietin production and reduced survival in pancreatic cancer. *J. Exp. Med.* 208:469–478. <http://dx.doi.org/10.1084/jem.20101876>
- Demols, A., O. Le Moine, F. Desalle, E. Quertinmont, J.L. Van Laethem, and J. Devière. 2000. CD4(+) T cells play an important role in acute experimental pancreatitis in mice. *Gastroenterology.* 118:582–590. [http://dx.doi.org/10.1016/S0016-5085\(00\)70265-4](http://dx.doi.org/10.1016/S0016-5085(00)70265-4)
- Ding, S.Q., Y. Li, Z.G. Zhou, C. Wang, L. Zhan, and B. Zhou. 2010. Toll-like receptor 4-mediated apoptosis of pancreatic cells in cerulein-induced acute pancreatitis in mice. *HBPD INT.* 9:645–650.

- Gasse, P., C. Mary, I. Guenon, N. Noulin, S. Charron, S. Schnyder-Candrian, B. Schnyder, S. Akira, V.F. Quesniaux, V. Lagente, et al. 2007. IL-1R1/MyD88 signaling and the inflammasome are essential in pulmonary inflammation and fibrosis in mice. *J. Clin. Invest.* 117:3786–3799.
- Greer, J.B., and D.C. Whitcomb. 2009. Inflammation and pancreatic cancer: an evidence-based review. *Curr. Opin. Pharmacol.* 9:411–418. <http://dx.doi.org/10.1016/j.coph.2009.06.011>
- Guo, J., J. Loke, F. Zheng, F. Hong, S. Yea, M. Fukata, M. Tarocchi, O.T. Abar, H. Huang, J.J. Sninsky, and S.L. Friedman. 2009. Functional linkage of cirrhosis-predictive single nucleotide polymorphisms of Toll-like receptor 4 to hepatic stellate cell responses. *Hepatology.* 49:960–968. <http://dx.doi.org/10.1002/hep.22697>
- Hauben, E., S. Gregori, E. Draghici, B. Migliavacca, S. Olivieri, M. Woisetschlager, and M.G. Roncarolo. 2008. Activation of the aryl hydrocarbon receptor promotes allograft-specific tolerance through direct and dendritic cell-mediated effects on regulatory T cells. *Blood.* 112:1214–1222. <http://dx.doi.org/10.1182/blood-2007-08-109843>
- Hellerbrand, S.C., S.C. Wang, H. Tsukamoto, D.A. Brenner, and R.A. Rippe. 1996. Expression of intracellular adhesion molecule 1 by activated hepatic stellate cells. *Hepatology.* 24:670–676. <http://dx.doi.org/10.1002/hep.510240333>
- Hingorani, S.R., E.F. Petricoin, A. Maitra, V. Rajapakse, C. King, M.A. Jacobetz, S. Ross, T.P. Conrads, T.D. Veenstra, B.A. Hitt, et al. 2003. Preinvasive and invasive ductal pancreatic cancer and its early detection in the mouse. *Cancer Cell.* 4:437–450. [http://dx.doi.org/10.1016/S1535-6108\(03\)00309-X](http://dx.doi.org/10.1016/S1535-6108(03)00309-X)
- Hou, B., B. Reizis, and A.L. DeFranco. 2008. Toll-like receptors activate innate and adaptive immunity by using dendritic cell-intrinsic and -extrinsic mechanisms. *Immunity.* 29:272–282. <http://dx.doi.org/10.1016/j.immuni.2008.05.016>
- Hruban, R.H., K. Takaori, D.S. Klimstra, N.V. Adsay, J. Albores-Saavedra, A.V. Biankin, S.A. Biankin, C. Compton, N. Fukushima, T. Furukawa, et al. 2004. An illustrated consensus on the classification of pancreatic intraepithelial neoplasia and intraductal papillary mucinous neoplasms. *Am. J. Surg. Pathol.* 28:977–987. <http://dx.doi.org/10.1097/01.pas.0000126675.59108.80>
- Ito, T., Y.H. Wang, O. Duramad, T. Hori, G.J. Delespese, N. Watanabe, F.X. Qin, Z. Yao, W. Cao, and Y.J. Liu. 2005. TSLP-activated dendritic cells induce an inflammatory T helper type 2 cell response through OX40 ligand. *J. Exp. Med.* 202:1213–1223. <http://dx.doi.org/10.1084/jem.20051135>
- Jura, N., H. Archer, and D. Bar-Sagi. 2005. Chronic pancreatitis, pancreatic adenocarcinoma and the black box in-between. *Cell Res.* 15:72–77. <http://dx.doi.org/10.1038/sj.cr.7290269>
- Kaisho, T., K. Hoshino, T. Iwabe, O. Takeuchi, T. Yasui, and S. Akira. 2002. Endotoxin can induce MyD88-deficient dendritic cells to support T(h)2 cell differentiation. *Int. Immunol.* 14:695–700. <http://dx.doi.org/10.1093/intimm/14.6.695>
- Kapsenberg, M.L. 2003. Dendritic-cell control of pathogen-driven T-cell polarization. *Nat. Rev. Immunol.* 3:984–993. <http://dx.doi.org/10.1038/nri1246>
- Lankisch, P.G., A. Löhr-Happe, J. Otto, and W. Creutzfeldt. 1993. Natural course in chronic pancreatitis. Pain, exocrine and endocrine pancreatic insufficiency and prognosis of the disease. *Digestion.* 54:148–155. <http://dx.doi.org/10.1159/000201029>
- Liu, T., T.L. Li, F. Zhao, C. Xie, A.M. Liu, X. Chen, C. Song, L. Cheng, and P.C. Yang. 2011. Role of thymic stromal lymphopoietin in the pathogenesis of nasal polyposis. *Am. J. Med. Sci.* 341:40–47. <http://dx.doi.org/10.1097/MAJ.0b013e3181f20489>
- Mantovani, A., C. Garlanda, and P. Allavena. 2010. Molecular pathways and targets in cancer-related inflammation. *Ann. Med.* 42:161–170. <http://dx.doi.org/10.3109/07853890903405753>
- Maraskovsky, E., K. Brasel, M. Teepe, E.R. Roux, S.D. Lyman, K. Shortman, and H.J. McKenna. 1996. Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J. Exp. Med.* 184:1953–1962. <http://dx.doi.org/10.1084/jem.184.5.1953>
- Masamune, A., T. Watanabe, K. Kikuta, and T. Shimosegawa. 2009. Roles of pancreatic stellate cells in pancreatic inflammation and fibrosis. *Clin. Gastroenterol. Hepatol.* 7:S48–S54. <http://dx.doi.org/10.1016/j.cgh.2009.07.038>
- Miller, G., S. Lahrs, V.G. Pillarisetty, A.B. Shah, and R.P. DeMatteo. 2002. Adenovirus infection enhances dendritic cell immunostimulatory properties and induces natural killer and T-cell-mediated tumor protection. *Cancer Res.* 62:5260–5266.
- Miller, G., S. Lahrs, and R.P. DeMatteo. 2003. Overexpression of interleukin-12 enables dendritic cells to activate NK cells and confer systemic antitumor immunity. *FASEB J.* 17:728–730.
- Nevala, W.K., C.M. Vachon, A.A. Leontovich, C.G. Scott, M.A. Thompson, and S.N. Markovic; Melanoma Study Group of the Mayo Clinic Cancer Center. 2009. Evidence of systemic Th2-driven chronic inflammation in patients with metastatic melanoma. *Clin. Cancer Res.* 15:1931–1939. <http://dx.doi.org/10.1158/1078-0432.CCR-08-1980>
- O'Neill, D.W., S. Adams, and N. Bhardwaj. 2004. Manipulating dendritic cell biology for the active immunotherapy of cancer. *Blood.* 104:2235–2246. <http://dx.doi.org/10.1182/blood-2003-12-4392>
- Oiva, J., H. Mustonen, M.L. Kylänpää, L. Kyhälä, K. Kuuliala, S. Siitonen, E. Kempainen, P. Puolakkainen, and H. Repo. 2010. Acute pancreatitis with organ dysfunction associates with abnormal blood lymphocyte signaling: controlled laboratory study. *Crit. Care.* 14:R207. <http://dx.doi.org/10.1186/cc9329>
- Omary, M.B., A. Lugea, A.W. Lowe, and S.J. Pandol. 2007. The pancreatic stellate cell: a star on the rise in pancreatic diseases. *J. Clin. Invest.* 117:50–59. <http://dx.doi.org/10.1172/JCI30082>
- Osawa, E., A. Nakajima, T. Fujisawa, Y.I. Kawamura, N. Toyama-Sorimachi, H. Nakagama, and T. Dohi. 2006. Predominant T helper type 2-inflammatory responses promote murine colon cancers. *Int. J. Cancer.* 118:2232–2236. <http://dx.doi.org/10.1002/ijc.21639>
- Palucka, K., H. Ueno, J. Fay, and J. Banchereau. 2009. Harnessing dendritic cells to generate cancer vaccines. *Ann. N. Y. Acad. Sci.* 1174:88–98. <http://dx.doi.org/10.1111/j.1749-6632.2009.05000.x>
- Petersen, T.R., N. Dickgreber, and I.F. Hermans. 2010. Tumor antigen presentation by dendritic cells. *Crit. Rev. Immunol.* 30:345–386. <http://dx.doi.org/10.1615/CritRevImmunol.v30.i4.30>
- Pylayeva-Gupta, Y., K.E. Lee, C.H. Hajdu, G. Miller, and D. Bar-Sagi. 2012. Oncogenic Kras-Induced GM-CSF Production Promotes the Development of Pancreatic Neoplasia. *Cancer Cell.* 21:836–847. <http://dx.doi.org/10.1016/j.ccr.2012.04.024>
- Rakoff-Nahoum, S., and R. Medzhitov. 2007. Regulation of spontaneous intestinal tumorigenesis through the adaptor protein MyD88. *Science.* 317:124–127. <http://dx.doi.org/10.1126/science.1140488>
- Ricci, F., S.E. Kern, R.H. Hruban, and C.A. Iacobuzio-Donahue. 2005. Stromal responses to carcinomas of the pancreas: juxtatumoral gene expression conforms to the infiltrating pattern and not the biologic subtype. *Cancer Biol. Ther.* 4:302–307. <http://dx.doi.org/10.4161/cbt.4.3.1501>
- Robson, N.C., S. Hoves, E. Maraskovsky, and M. Schnurr. 2010. Presentation of tumour antigens by dendritic cells and challenges faced. *Curr. Opin. Immunol.* 22:137–144. <http://dx.doi.org/10.1016/j.coi.2010.01.002>
- Sharif, R., R. Dawra, K. Wasiluk, P. Phillips, V. Dudeja, E. Kurt-Jones, R. Finberg, and A. Saluja. 2009. Impact of toll-like receptor 4 on the severity of acute pancreatitis and pancreatitis-associated lung injury in mice. *Gut.* 58:813–819. <http://dx.doi.org/10.1136/gut.2008.170423>
- Swann, J.B., M.D. Vesely, A. Silva, J. Sharkey, S. Akira, R.D. Schreiber, and M.J. Smyth. 2008. Demonstration of inflammation-induced cancer and cancer immunoediting during primary tumorigenesis. *Proc. Natl. Acad. Sci. USA.* 105:652–656. <http://dx.doi.org/10.1073/pnas.0708594105>
- Vaquero, E., I. Gukovsky, V. Zaninovic, A.S. Gukovskaya, and S.J. Pandol. 2001. Localized pancreatic NF-kappaB activation and inflammatory response in taurocholate-induced pancreatitis. *Am. J. Physiol. Gastrointest. Liver Physiol.* 280:G1197–G1208.
- Vincent, A., J. Herman, R. Schulick, R.H. Hruban, and M. Goggins. 2011. Pancreatic cancer. *Lancet.* 378:607–620. [http://dx.doi.org/10.1016/S0140-6736\(10\)62307-0](http://dx.doi.org/10.1016/S0140-6736(10)62307-0)
- Wang, Z., D. Kong, S. Banerjee, Y. Li, N.V. Adsay, J. Abbruzzese, and F.H. Sarkar. 2007. Down-regulation of platelet-derived growth factor-D inhibits cell growth and angiogenesis through inactivation of Notch-1 and nuclear factor-kappaB signaling. *Cancer Res.* 67:11377–11385. <http://dx.doi.org/10.1158/0008-5472.CAN-07-2803>
- Zhou, X.Y., Z.G. Zhou, J.L. Ding, L. Wang, R. Wang, B. Zhou, J. Gu, X.F. Sun, and Y. Li. 2010. TRAF6 as the key adaptor of TLR4 signaling pathway is involved in acute pancreatitis. *Pancreas.* 39:359–366. <http://dx.doi.org/10.1097/MPA.0b013e3181bb9073>

SUPPLEMENTAL MATERIAL

Ochi et al., <http://www.jem.org/cgi/content/full/jem.20111706/DC1>




Figure S1. Divergent effects of modulating MyD88 and TRIF signaling on pancreatic carcinogenesis.