

Shelfbreak Acoustics: The ASIAEX Volume Interaction Experiment

James F. Lynch
MS #11

Woods Hole Oceanographic Institution
Woods Hole, MA 02543

Phone: (508)289-2230 fax:(508) 457-2194 email: jlynch@whoi.edu

Grant Number: N00014-98-1-0413

<http://www.oal.whoi.edu/AO/topics/Primer/Primer.html>

<http://www.oal.whoi.edu/ASIAEX/ASIAEX.html>

LONG TERM GOALS

The long term goal of our work in the ASIAEX program is to understand the nature of low frequency (50-600 Hz) acoustic propagation and scattering in shallow water when strong oceanic variability in the form of fronts, eddies, boundary layers, and internal waves is present. We are also greatly interested in the effects of geological variability, which is as important to acoustics as ocean variability in shallow water regions.

OBJECTIVES

In the ASIAEX experiment, there were a large number of specific acoustics objectives that were pursued - we will not list them all here. Some of these objectives are being pursued by WHOI and some by other ASIAEX P.I.'s. Over the last year, we have specifically focused on the issues of: 1) horizontal array coherence in shallow water, 2) the travel time and intensity fluctuations of the acoustic field in the 50-600 Hz band, and 3) the ambient noise field in the 50-1400 Hz band. These efforts both reinforced and extended the goals of the two predecessor experiments to ASIAEX, the 1995 SWARM internal wave experiment and the 1996-97 PRIMER shelfbreak front experiment. We also are pursuing studies of the physical oceanography and geology of the South China Sea (SCS) volume experiment region, in an interdisciplinary approach together with other ASIAEX investigators.

APPROACH

Similar to the SWARM and PRIMER experiments, the ASIAEX SCS effort combined high-resolution physical oceanographic and geologic measurements with precise measurements of the acoustic field (see Figure 1.) Acoustically, the heart of our experiment was a combined vertical/horizontal aperture array, which listened to both moored and towed sources with frequencies from 50 Hz to 600 Hz. The moored sources enabled us to examine acoustic propagation time series both for a cross slope and an along slope geometry. The towed sources allowed us to study propagation loss versus range, a simple study that had been intended in previous experiments, but had never been accomplished due to time constraints. The receiver array was also able to support ambient noise studies in the 50-1400 Hz band. The oceanographic support for the acoustics was quite extensive, consisting of twenty-nine separate oceanographic moorings, a Sea Soar hydrography survey, a high frequency acoustic backscatter survey, satellite imagery (SAR and AVHRR), and numerous CTD casts. The geological support was also good, consisting of two high-resolution chirp sonar surveys along the stationary propagation

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2002		2. REPORT TYPE		3. DATES COVERED 00-00-2002 to 00-00-2002	
4. TITLE AND SUBTITLE Shelfbreak Acoustics: The ASIAEX Volume Interaction Experiment				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Woods Hole Oceanographic Institution,,Woods Hole,,MA, 02543				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The long term goal of our work in the ASIAEX program is to understand the nature of low frequency (50-600 Hz) acoustic propagation and scattering in shallow water when strong oceanic variability in the form of fronts, eddies, boundary layers, and internal waves is present. We are also greatly interested in the effects of geological variability, which is as important to acoustics as ocean variability in shallow water regions.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




Figure 1. Upper panel shows geographical location of ASIAEX SCS volume interaction experiment. Lower panel shows locations of moorings, as well as fixed acoustic paths and Sea Soar tracks.

paths, as well as the propagation loss tows. (We had hoped for somewhat more geological data, but area permission problems in the FY 2000 preliminary experiment precluded our obtaining all the data we desired. We are hoping that a further chirp sonar survey by L. Bartek, planned for spring, 1993, can remedy this.) Our experiment extended temporally over a three-week period, so that we were able to examine the effects of a full spring-neap tidal cycle on the acoustics transmissions.

WORK COMPLETED

Our main accomplishment this year was the initiation of full scale data analysis efforts, concentrating on (as mentioned above) horizontal array coherence, travel time and amplitude fluctuations, and the ambient noise field. This work was performed in close collaboration with other ASIAEX PI's, specifically: M. Orr (and his group) from the Naval Research Laboratory (horizontal array coherence), Lixin Wu of the National Laboratory of Acoustics in Beijing and T. Duda of WHOI (amplitude and travel time fluctuations), and R.C. Wei of the National Sun Yat Sen University in Kaohsiung (ambient

noise). Each of these studies has progressed well, and we are planning to publish a paper on each topic in the upcoming IEEE Journal of Oceanic Engineering special issue devoted to the Asian Marginal Seas. The current results in each area will also be presented at an ASIAEX workshop in Chengdu, China this coming October. It should be noted that Drs. Wu and Wei both came to WHOI as Guest Investigators during the summer of 2002, enabling us to work closely. Several visits to NRL also help facilitate the interaction on the analysis work. Additionally, we (Newhall and Lynch) were able to visit both the Beijing and Kaohsiung laboratories of our Guest Investigators this past year as guests of their institutions.

RESULTS

We are at the “intermediate” stage in all the analyses, which means that we now have some interesting results to report, though not the complete story.

In the horizontal array coherence analysis, we have performed the critical array element location studies using three different methods: 1) long-baseline high frequency navigation, 2) dropped light-bulb sources, and 3) distant source low frequency navigation. All these methods showed basic agreement, with the distant, low-frequency source localization showing probably the best accuracy. Having these localizations in hand, we have embarked on obtaining the horizontal coherence length via two methods: 1) phone-to-phone cross correlation of the pulses and 2) studies of the beamformed array gain. The correlation calculations gave an $\sim 20\lambda$ value for the array horizontal coherence, which is probably a lower bound, as the mode temporal dispersion due to the non-perpendicular array geometry probably decreased this horizontal coherence estimate. The beamformed outputs have been obtained for most of the acoustic frequencies used, but we are just at the stage of converting these into coherence length estimates.

Turning to the travel time and intensity fluctuation analyses, we again are at the halfway stage, where we have time series of many quantities of interest, but have not studied all the inter-correlations of the acoustics and environmental variables. In Figure 2, we show an example of the leading edge travel time (black dots – green is a higher threshold), pulse intensity centroid (blue), and full pulse width (red) over an approximately one day period, showing clear evidence of the diurnal tidal modulation of these signals. We currently have full time series for these travel time related variables, as well as most of the intensity variables. Time series for the prime oceanographic correlates to the acoustics (temperature and current) also exist, and so we should soon be able to comment in some detail about the forcing of the acoustics by the environment. Finally, we note that we also have noise time series at 700 Hz and above, and at 350-450 Hz, with more bands to be analyzed shortly. Again, these time series can be quickly correlated to environmental forcing.




Figure 2. Tracking of the leading edge travel time (black and green dots, using slightly different thresholds), pulse centroid (blue dots), and full pulse time spread (red dots) for one day of data from the ASIAEX experiment. (Courtesy T. Duda.)

IMPACT

Probably the greatest impact that the ASIAEX volume interaction experiment will have is in seeing just how well we can predict the acoustic field and its variability/fluctuation level for a case where we can check the answers against ground truth. Additionally, we can directly examine important quantities such as horizontal coherence length and the frequency dependence of acoustic propagation for the ASIAEX region. Finally, the oceanographic data taken will be of great use in calibrating Navy models, such as MODAS.

TRANSITIONS

One eventual transition of our data will be to ONR's Uncertainty DRI program, where the interest is in "the error bars" in ocean acoustic field and system performance prediction. Also, our ASIAEX data provide a unique database in a region where there are relatively few high-quality data.

RELATED PROJECTS

The SWARM acoustics/internal wave study and the PRIMER acoustics/shelfbreak front study were direct predecessors of ASIAEX, and examined some of the same scientific issues, only with fewer measurement resources. The Uncertainty DRI is also a closely related project to ASIAEX, as it will use some of the data.

PUBLICATIONS (ASIAEX, PRIMER, and other Shelfbreak Experiments)

J. Lynch, A. Newhall, B. Sperry, G. Gawarkiewicz, P. Tyack, and C.S. Chiu, “Spatial and temporal variations in acoustic propagation characteristics at the New England shelfbreak front”, submitted to *IEEE J. Oceanic Eng’g.* (2001).

J. Lynch and M. Orr, “Low frequency acoustic propagation through shallow water internal waves.” Review article/chapter draft submitted to ONR for forthcoming book on internal waves and acoustics. (2002).

G. Gawarkiewicz, F. Bahr, K. Brink, R.C. Beardsley, M. Caruso, J. Lynch, and C.S. Chiu, “A large amplitude meander of the shelfbreak front during summer south of New England: Observations from the shelfbreak PRIMER experiment.”, in press *J. Geophys. Res.* (2002).

P. Abbot, S. Celuzza, I. Dyer, B. Gomes, J. Fulford, and J. Lynch, “Effects of the East China Sea shallow water environment on acoustic propagation,” submitted to *IEEE J. Oceanic Eng’g.* (2002).

G. R. Potty, J.H. Miller, and J.F. Lynch, “Inversion for sediment geoacoustic properties at the New England Bight,” in press *J. Acoust. Soc. Am.* (2002).

W. Carey, P. Cable, W. Siegmann, J. Lynch and I. Rozenfeld, “Measurement of sound transmission and signal gain in the Strait of Korea.” In press, *IEEE J. Oceanic Eng’g.* (2002)

B. Sperry, J. Lynch, G. Gawarkiewicz, C. Chiu, and A. Newhall, “Characteristics of acoustic propagation to the eastern vertical line receiver during the summer 1996 New England shelfbreak PRIMER experiment,” submitted to *IEEE J. Oceanic Eng’g.* (2002).

J.F. Lynch, “Preliminary results from the 2001 ASIAEX South China Sea Experiment”, Proceedings of the IEEE Post Underwater Technology 2002 (UT’02) Workshop, pp 113-117, Taipei, Taiwan, April, 2002.

J. F. Lynch, A. Fredricks, J. Colosi, G. Gawarkiewicz, B. Sperry, A. Newhall, and C.S. Chiu, “An overview of the acoustic fluctuations from the summer 1996 New England Shelfbreak PRIMER experiment”, Conference on Undersea Technology 2002, pp. A1.1-A1.9, National Taiwan University, Taipei, Taiwan, April, 2002.

J.F. Lynch and the ASIAEX group, “The ASIAEX South China Sea Experiment”, WHOI 2001 Annual Report, pp. 7-8, 2002.

J. Lynch, A. Fredricks, J. Colosi, G. Gawarkiewicz, A. Newhall, C.S. Chiu, and M. Orr, “Acoustic effects of environmental variability in the SWARM, PRIMER and ASIAEX experiments.”

Conference proceedings paper for SACLANT conference on “Impact of littoral environmental variability on acoustic predictions and sonar performance,” September, 2002. To be published in Kluwer volume.