

An Observational and Modeling Study of Air-Sea Fluxes at Very High Wind Speeds

Kerry Emanuel
Room 54-1620, MIT
77 Massachusetts Avenue
Cambridge, MA 02139
phone: (617) 253-2462 fax: (425) 740-9133 email: emanuel@texmex.mit.edu

Peter Black
NOAA/ERL Hurricane Research Division
4301 Rickenbacker Causeway
Miami, FL 33149-1097
phone: (305) 361-4320 fax: (305) 361-4402 email: black@aoml.noaa.gov

Award Number N00014 01-1-0162
<http://www-paoc.mit.edu/~emanuel/home.html>

LONG-TERM GOALS

Our ultimate objective is to understand and be able to predict changes in the intensities of hurricanes. Over the last three decades, there has been considerable improvement in both our understanding and our ability to predict the tracks of hurricanes, but there is, by contrast, little skill in predicting hurricane intensity change.

OBJECTIVES

The objective of the work carried out under this grant is to understand and be able to model fluxes of enthalpy, moisture and momentum between the ocean and atmosphere in all ranges of wind speed, up to and including wind speeds encountered in severe hurricanes. Modeling such fluxes is critical to predicting hurricane intensity.

APPROACH

Our approach is to use atmospheric measurements collected in actual hurricanes to deduce surface fluxes of enthalpy and momentum. These measurements consist of wind, temperature and humidity measured directly from research reconnaissance aircraft and from GPS dropwindsondes deployed from those aircraft. Such measurements are first used to construct the azimuthally averaged radius-height distributions of angular momentum, moist static energy and radial velocity. Then, assuming that the storm is in an approximately steady state, the radial advections of moist static energy and angular momentum are calculated, and the surface fluxes of angular momentum and enthalpy are deduced as those needed to maintain a steady state. This approach has been taken (for angular momentum only) by (Hawkins and Rubsam, 1968), but they had to rely on measurements somewhat inferior to those available today. Our work is being carried out under the leadership of the principal investigators Kerry Emanuel of MIT, who is supervising the analysis of the aircraft and dropwindsonde data, and Peter Black of the NOAA/AOML Hurricane Research Division, who is providing the aircraft data and using

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2002		2. REPORT TYPE		3. DATES COVERED 00-00-2002 to 00-00-2002	
4. TITLE AND SUBTITLE An Observational and Modeling Study of Air-Sea Fluxes at Very High Wind Speeds				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) MIT,Room 54-1620,,77 Massachusetts Avenue,,Cambridge,,MA,02139				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Our ultimate objective is to understand and be able to predict changes in the intensities of hurricanes. Over the last three decades, there has been considerable improvement in both our understanding and our ability to predict the tracks of hurricanes, but there is, by contrast, little skill in predicting hurricane intensity change.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

the early results of our research to plan field operations during the second and third years. Analysis of the existing aircraft data is being carried out by William Ramstrom, an advanced graduate student at MIT. The approach we are taking under this grant compliments a laboratory modeling approach we are using under separate funding.

WORK COMPLETED


As planned for the first year of our work, we have completed an analysis of existing aircraft and GPS dropwindsonde data and published the results as a Master's Thesis (Ramstrom, 2001). During this past year, we conducted a preliminary field campaign in which we tested our strategy for deploying a dense line of GPS dropwindsondes across the eyewalls of Atlantic hurricanes, coupled with detailed aircraft measurements of turbulent fluxes across the top of the inflow layer. As of this writing, this field campaign is continuing; so far we have one good case of dense eyewall measurements in a strong, nearly axisymmetric storm (Isidore).

RESULTS

Until this year, the measurement programs were not developed with the kind of analysis we are now undertaking in mind, and consequently there are strong limitations to what can be done with them. Nevertheless, we were able to deduce surface drag coefficients in a number of hurricanes that show remarkable consistency with previous inferences and with very recent laboratory experiments. The 10 meter drag coefficients varied between 0.0027 and 0.0030 for wind speeds in the 40-60 m s⁻¹ range, with some tendency to increase with wind speed, although the spatial density of measurements did not allow us to infer a meaningful relationship between the drag coefficient and wind speed. As expected, we did not see much evidence for strong vertical turbulent fluxes of angular momentum out of the top of the boundary layer.

Deduction of the surface enthalpy fluxes, here attempted for the first time, proved more problematic, as in most of the storms we surveyed, the turbulent entropy flux out of the top of the boundary layer was comparable in magnitude to the inferred surface fluxes. These turbulent fluxes had to be inferred from aircraft data, but the aircraft flight tracks were not ideally suited to the task. Our best estimates of the 10 meter exchange coefficient for enthalpy are very comparable to the drag coefficient values, indicating that the ratio of the exchange coefficients for enthalpy and momentum is nearly unity.

We also developed a technique for calculating the ratio of the exchange coefficients directly from the radial distributions of angular momentum and moist static energy, without having to use the radial velocity, which is very sensitive to errors in the location of the center of the storm. The ratio can be estimated directly from the gradient of moist static energy with respect to potential radius, which is a measure of the angular momentum per unit mass. An example of such a calculation is shown below.



The most important findings of the first year research concern what is needed in the field experimental efforts that will take place in subsequent years. (Indeed, that was the primary objective of the first year's effort.) We are now in an excellent position to specify in some detail the field experimental requirements that will yield high quality estimates of exchange coefficients in hurricanes. In the second year, these findings were put into effect during the NOAA/HRD 2002 Atlantic hurricane field program. As of this writing, this field campaign is ongoing and useful measurements have been obtained in at least one storm.

IMPACT/APPLICATIONS

It is too soon to predict how the results obtained thus far will influence science, but as mentioned above, they will have a strong influence on the future conduct of this project in so far as planning the field experimental phase goes.

TRANSITIONS

These results are now being used to plan the next phase of the project.

RELATED PROJECTS

One of the P.I.'s (Emanuel) is conducting a series of experiments with an annular wind-wave flume. Equivalent 10 m winds speeds of 70 m s^{-1} can be achieved in this flume. By measuring the rate of spin down of the water mass after the airflow has been shut down, one can deduce rather precisely the drag coefficient. Similarly, the enthalpy flux coefficient is estimated by measuring the rate at which the water must be heated to keep it at room temperature. These experiments have shown that the drag coefficient increases linearly with wind speed up to about 30 m s^{-1} , but then levels off with no further increase. Experiments to deduce the enthalpy flux show a similar leveling off of the exchange coefficient at high wind speed.

Working with Ed Andreas, Emanuel has explored the effect of parameterized fluxes of momentum and enthalpy owing to sea spray on numerical simulations of tropical cyclones. This work has been published (Andreas and Emanuel, 2001).

Based on the findings of the first year's research and on the aforementioned flume experiments, the first PI submitted a theoretical paper presenting a similarity theory for air-sea fluxes at very high wind speeds (Emanuel, 2003).

SUMMARY

We have used aircraft measurements of wind, temperature and humidity in several hurricanes to deduce the drag that hurricane winds exert on the ocean surface, and the evaporation of seawater into a hurricane, which is the process that fuels the storm. Understanding the drag and the evaporation is crucial for understanding and predicting changes in hurricane wind speeds. While the estimates we made in this first year of the project are useful and consistent with independent estimates, their primary value is in designing a campaign for making much better measurements in hurricanes, which we began doing this 2002 hurricane season.

REFERENCES

Andreas, E. L. and K. Emanuel, 2001: Effects of sea spray on tropical cyclone intensity. *J. Atmos. Sci.*, **58**, 3741-3751

Emanuel, K., 2003: A similarity hypothesis for air-sea exchange at extreme wind speeds. *J. Atmos. Sci.*, **60**, conditionally accepted

Hawkins, H. F. and D. T. Rubsam, 1968: Hurricane Hilda, 1964: II. Structure and budgets of the hurricane on October 1, 1964. *Mon. Wea. Rev.*, **96**, 617-636

Ramstrom, W. D., 2001: Tropical cyclone energy and momentum fluxes. Earth, atmospheric and planetary sciences, Mass. Inst. Tech., 84 pp.

PUBLICATIONS

Emanuel, K., 2003: A similarity hypothesis for air-sea exchange at extreme wind speeds. *J. Atmos. Sci.*, **60**, conditionally accepted

Ramstrom, W. D., 2001: Tropical cyclone energy and momentum fluxes. Earth, atmospheric and planetary sciences, Mass. Inst. Tech., 84 pp.

Ramstrom, W. D., 2002: Tropical cyclone enthalpy and angular momentum fluxes. Preprints of the 22nd Conference on Hurricanes and Tropical Meteorology, Paper 16C.6, Amer. Meteor. Soc., Boston

PATENTS

None.