

UPPER OCEAN MIXING

Michael Gregg, PI
522 Henderson Hall
University of Washington
1013 NE 40th St.
Seattle, WA 98105-6698

Phone: (206)543-1353; fax (206)543-6785; email:gregg@apl.washington.edu
Award #: N00014-94-1-0079, N00014-94-1-0082, N00024-95-1-0406, N00014-93-1-0402, N00014-97-1-0136

Jack Miller, Co-PI
508 Henderson Hall
University of Washington
1013 NE 40th St.
Seattle, WA 98105-6698

Phone: (206)543-9958; fax (206)543-6785; email:miller@apl.washington.edu
Award #: N00014-94-1-0079, N00014-94-1-0082, N00024-95-1-0406, N00014-93-1-0402, N00014-97-1-0136

LONG-TERM GOAL

My long-term goal is to identify, quantify, and parameterize the major mixing processes in the upper ocean.

OBJECTIVES

My scientific objectives are to measure mixing directly with microstructure sensors and to relate these measurements to the larger scales producing the mixing in such a way that the results can be compared with either the large-scale changes resulting from the mixing or with theoretical predictions of mixing rates. My technological objectives are to develop instruments and sensors to measure the major mixing parameters.

APPROACH

Because a first-order understanding of mixing in the open ocean is rapidly being developed, our approach has shifted to obtaining a similar understanding of mixing near coasts and in estuaries. Because dissipation rates are higher in these waters, we have also shifted our technological developments to improve the spatial resolution of microstructure sensors and to adapt open-ocean measurements of finescale velocity to shallow water.

WORK COMPLETED

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Upper Ocean Mixing				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 N.E. 40th Street, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

In May and April 1997 we participated in the Coastal Mixing and Optics field program on the New England continental shelf. In August 1997 we observed shear and mixing over the narrow continental shelf and slope off Monterey, California as part of the Littoral Internal Waves Initiative. Both cruises were successful and sampled markedly different mixing regimes.

We finished and are writing up one phase of the analysis of our observations of flow and mixing in Knight Inlet, B.C. with a study of the three-dimensional flow field.

Analysis of observations taken in the Bosphorus is nearing completion. Gregg traveled to Turkey to work with colleagues at the Middle East Technical University, both to understand particulars of the observations and to put them into the context of earlier work in the Strait. Understanding how the flow is controlled is proving difficult owing to the complicated bathymetry of the Bosphorus and to the shallowness of surface currents in the southern end of the Strait.

RESULTS

Flows over sills often limit water exchanges between inland seas and estuaries with the open ocean. Previous studies have assumed that the flows are two-dimensional, but in Knight Inlet we find that cross-channel flows and variability are important and must be considered to understand the flow adequately. To our knowledge, this is the first observation of this type of recirculation over sills in channels.

When water is strongly stratified over the New England shelf, internal solibores generated at the shelf break by tidal currents propagate shoreward every tidal cycle. At the CMO site the solibores depress the thermocline, which slowly recovers to its original position. Superimposed on this structure are much more rapid oscillations resulting from the solitary wave packet evolving within the solibore. Background mixing rates are very low between solibores but increase at least 50-fold during the early stages of the solibore. Consequently, most of the mixing occurs while the solibores are transiting

IMPACT/APPLICATION

Owing to its simple geometry, Knight Inlet was more likely to be dominated by the along-channel flow than many other sites where two-dimensional regimes are assumed. Demonstration of the importance of the three-dimensional circulation in Knight Inlet will lead to three-dimensional studies in many other flow regimes influenced by topography.

TRANSITIONS

The identification of solibores as major mixing agents on continental shelves demonstrates that existing parameterizations of mixing on shelves are improperly formulated and should lead to improved parameterizations.

Instruments developed with ONR-core and CMO funding were used during the LIWI work as well as during a study of mixing in Monterey Canyon funded by NSF.

RELATED PROJECTS

- Shear and Mixing in Monterey Canyon, with Eric Kunze and Leslie Rosenfield and funded by NSF.
- Development of a thinistor for measuring temperature microstructure, proposed to NSF with Professor Fumio Ohuchi (University of Washington) based on preliminary work funded by ONR.
- Observations of intense mixing in the Banda Sea, proposed to NSF to use instruments developed with ONR funding.