

FIELD MEASUREMENTS OF SEDIMENT TRANSPORT PROCESSES IN STRATAFORM: EXTENDED DURATION OBSERVATIONS

Richard W. Sternberg
Andrea Ogston
School of Oceanography, Box 357940
University of Washington
Seattle, WA 98195-7940
phone: (206) 543-0589 fax: (206) 543-6073 email: rws@ocean.washington.edu
Award #N00014-95-1-0418, N00014-97-1-0377, N00014-97-1-1011

LONG-TERM GOALS

The long-term goal is to increase our understanding of the role of physical forcing mechanisms and associated sediment response that contribute to the formation of fine scale strata and cross-margin transport of suspended particulates on the Eel River continental margin.

SCIENTIFIC OBJECTIVES

The major objective is to carry out a measurement program over an extended period to document fluid motions and sediment transport at selected sites on the continental shelf and slope in the STRATAFORM study area (Eel River, California). The purpose of extended duration observations is to increase the probability of documenting the physical forcing and stratigraphic response to a wider range of major sediment transport events than possible in a single year. Extended duration observations began in September 1995 and have focused on two specific regions of interest along a cross-margin transect seaward of Arcata, California: the sand/mud transition (60 m depth); and the slope depositional region (450 m depth). The measurement program consists of:

1. Collection of a continuous time series of near-bed flow and suspended sediment concentration to document the causes and magnitude of shelf sedimentation events at the sand/mud transition and their linkages to the formation of event beds.
2. Collection of a continuous time series of flow and suspended sediment concentration at three elevations within the water column on the upper slope (450 m depth) to document off-shelf sediment transport and the causes and magnitudes of transport events.

These observations will be used to evaluate the nature of suspended particulate matter being introduced onto the shelf and the processes responsible for its subsequent transport and deposition. Sediment response is being evaluated in terms of frequency of motion, direction of transport, mass transport, and the relationship between suspension events and strata formation. The data are also being utilized by the modeling groups developing strata formation and preservation models.

APPROACH

1. Platform: Benthic tripod at sand/mud boundary (60 m depth)

Measurements: Current speed and direction at two elevations, waves and tides (pressure), bed elevation and changes (acoustic altimeter), suspended sediment concentration (optical backscatter sensor) at two elevations, in situ particle size (LISST), two video cameras (bed configuration, floc size, and settling velocity), CTD, temperature sensors at two elevations.

Deployment: Two identical tripods are being exchanged every four months beginning on 24 September 1995.

2. Platform: Mooring on upper continental slope (450 m depth)

Measurements: Three identical instrument packages attached at different elevations. Each instrument package consists of (1) an InterOcean S4a electromagnetic current meter, CTD, and

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Field Measurements of Sediment Transport Processes in Strataform: Extended Duration Observations				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington,School of Oceanography,Seattle,WA,98195				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

beam transmissometer, and (2) a time sequence sediment trap. The nearbed array is located 15 m above the seabed, the midwater array is located at approximately 160 m depth (depth of the shelf break), and the near surface array is positioned approximately 60 m below the sea surface. Additionally, ten temperature sensors are spaced over the mooring length to document internal wave activity.

Deployment: Two identical moorings are being exchanged on a six-month interval beginning on 27 September 1995.

TASKS COMPLETED

Data collection has spanned a two-year period and is still ongoing. Two aspects of the shelf measurement program have been investigated: (1) particle characteristics measured by the floc - camera; (2) sediment transport events occurring over an annual period.

RESULTS

Analysis of in-situ particle characteristics show that suspended sediment occurs as aggregates with mean sizes ranging from 130 μ (detection limit) to 760 μ . In terms of mass distribution the dominant aggregate size is 600 μ . Component particles comprising the flocs are 0.5% sand, 45.3% silt, 54.2% clay. Settling velocity of the floc population varies from 0.09 cm s^{-1} to 8.13 cm s^{-1} (3.8 cm s^{-1} for the dominant aggregate size).

Analysis of sediment transport events over the first year show a very energetic shelf with frequent suspension events and strong seasonal variability (Table 1).

TABLE 1. Seasonal Summary of Events

	ALL EVENTS	AUTUMN	WINTER	SPRING	SUMMER
Number of events	41 (21 south/ 20 north) (35 offshore / 6 onshore)	15 (8 south / 7 north) (12 offshore/ 3 onshore)	20 (11 south/ 9 north) (18 off / 2 on)	5 (1 south/ 4 north) (5 off / 0 on)	1 (1 south/ 0 north) (0 off / 1 on)
Event recurrence	7.1 days	3.3 days	4.5 days	18 days	73 days
Mean duration	3.1 days	1.9 days	3.1 days	4.8 days	7.5 days
Mean conc.	100 mg l^{-1}	80 mg l^{-1}	130 mg l^{-1}	60 mg l^{-1}	35 mg l^{-1}
Peak conc.	300 mg l^{-1}	150 mg l^{-1}	300 mg l^{-1}	90 mg l^{-1}	--- mg l^{-1}
Alongshelf transport	-46200 kg m^{-2} (south)	-1230 kg m^{-2}	-45100 kg m^{-2}	2960 kg m^{-2}	-2770 kg m^{-2}
Cross shelf transport	-38600 kg m^{-2} (offshore)	-4430 kg m^{-2}	-28300 kg m^{-2}	-6180 kg m^{-2}	390 kg m^{-2}

IMPACT ON SCIENCE

These results broaden our views of sedimentology related to energetic continental shelf environments with episodic sediment input. Particles being transported are large aggregates with extremely high settling velocities. The high settling velocity suggests that particles are deposited from the river plume relatively quickly, probably undergo numerous resuspension episodes in their transport offshore, and when undergoing transport remain close to the seabed with some evidence of fluid mud formation. This result is supported by physical observations which show frequent wave and current events capable of resuspending sediment and maintaining high concentration nearbed layers through the high river discharge months. Net particle flux is south and offshore supplying sediment to the upper slope and adjacent submarine canyons.

TRANSITIONS

These data are being used by other groups in STRATAFORM including the shelf and slope research groups and also represents primary input to the shelf and slope modeling groups.

REFERENCES

A.S. Ogston and R.W. Sternberg (in press) Sediment transport events on the Northern California continental shelf. Mar. Geol.

R.W. Sternberg, I. Berhane and A. Ogston (in press) Measurement of size and settling velocity of suspended aggregates on the Northern California continental shelf. Mar. Geol.