

SEQUENCE STRATIGRAPHIC INTERPRETATION AND MAPPING, NEW JERSEY MARGIN

Craig S. Fulthorpe and James A. Austin, Jr.

University of Texas, Institute for Geophysics, 4412 Spicewood Springs Road, Bldg. 600

Austin, TX 78759-8500

Phone: (512) 471-0459; Fax: (512) 471-8844

craig@utig.ig.utexas.edu

Phone: (512) 471-0450; Fax: (512) 471-8844

jamie@utig.ig.utexas.edu

Award #: N00014-96-1-0037

LONG-TERM GOALS

This project forms part of a long-term effort, linked to ONR's STRATAFORM program, to document the evolution of the middle Atlantic continental shelf and slope, offshore New Jersey, at a variety of spatial scales (lateral and vertical) and in three dimensions (3-D; Austin et al., 1996; 1997). The overall goal is to determine how short-term geological processes contribute to the formation of the preserved record of depositional sequences.

SCIENTIFIC OBJECTIVES

Depositional sequences are unconformity-bounded packets of sediment that are the building blocks of sedimentary deposits. They form in response to fluctuations in geological and environmental conditions, notably global sea-level (eustasy). We have delineated the 3-D geometries of Miocene clinoform sequences in order to investigate: 1) whether known, rapid Miocene shelf progradation occurred predominantly in response to point sources of sediment or to distributed cross-shelf sediment transport; 2) paleo-water depths at Miocene clinoform breakpoints, crucial to eustatic amplitude estimates and for understanding processes of sediment delivery to clinoform slopes during periods of rapid progradation; 3) the origins and spatial distribution of canyons during the Miocene.

APPROACH

A great deal of oil-industry multichannel seismic (MCS) data was acquired on the middle Atlantic margin beginning in the mid-1970s, when leasing there commenced. We selected a grid of 36-fold MCS data shot by Digicon in 1975, encompassing ODP sites on the slope and mid-outer shelf (Figure 1). We acquired the seismic profiles in digital form on magnetic tape in order to interpret and map the upper (seafloor - 1 sec.) interval using GeoQuest seismic interpretation software. While not true 3-D data, the dense grid allows mapping of the 3-D evolution of elements of the shelf sediment prism at the scale of individual sequences (i.e., vertical thickness • 100 m, horizontal dimensions of a few km to several tens of km).

WORK COMPLETED

The data tapes were not in SEG Y format and some had deteriorated or been damaged (some data bits in both traces and trace headers were degraded). Software was therefore developed to convert the data to SEG Y format and renumber common-depth points. Sequence boundaries approximately correlative with four slope reflections dated by ODP Leg 150 (m3, ~13.6 Ma; m2, ~12.5 Ma; m1, ~11.0 Ma; m0.5, ~8.0 Ma) have been mapped across the northern portion of the grid in a region extending ~70 km along strike and ~50 km down-dip (Figure 1; Fulthorpe and Austin, in press). This is the area of most immediate relevance to STRATAFORM objectives, linking: 1) ODP slope and shelf sites (Legs 150 and 174A), 2) 1993 UTIG Huntec 2-D/3-D survey, 3) MCS and SCS data acquired in support of Leg 150, and 4) 1995 MCS regional and hazards surveys acquired for STRATAFORM (Figure 1).

RESULTS

Miocene clinoform breakpoints are linear to gently arcuate in plan (Figure 2). A bulge in the margin migrated southward during deposition of successive clinoforms suggesting a progressive increase in volume of

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Sequence Stratigraphic Interpretation and Mapping, New Jersey Margin				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Texas at Austin, Institute for Geophysics (UTIG), 4412 Spicewood Springs Road, Bldg. 600, Austin, TX, 78759				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

sediment from southern sources relative to those in the north during the period ~13.6 to ~8. Ma. Miocene canyons were rare: only one V-shaped clinoform slope canyon was recognized. Mapping also confirmed the existence of rare planar-floored canyons seaward of clinoform toes. V-shaped and planar-floored canyon geometries are indicative of separate depositional histories. The lack of evidence for canyons breaching clinoform breakpoints suggests that Miocene rivers did not discharge directly at paleo-shelf edges during progradation. Comparison of reconstructed paleo-elevations of Miocene breakpoints with the Exxon eustatic curve supports the view that m3, m2 and m0.5 were not exposed during lowstands, though m1 may have lain close to sea level.

IMPACT/APPLICATIONS

Sequence stratigraphic models have assumed that sea level fell to, or below, the clinoform breakpoint at lowstands, so that sediment could be delivered directly to the slope. Our results, however, suggest that Miocene river mouths were generally located landward of breakpoints throughout each sea-level cycle. If correct, shelf sediment transport processes must have controlled both progradation and the efficient lateral dispersal of sediment and muting of point-source effects indicated by the near-linear breakpoint trends. In addition, paleo-water depth estimates at m3, m2 and m0.5 suggest that elevations of corresponding lowstands on the Haq et al. (1987) eustatic curve may be too high by several tens of meters.

TRANSITIONS

The mapped clinoforms will be the subject of a numerical modeling effort by G. Karner of Lamont-Doherty Earth Observatory.

RELATED PROJECTS

This work has close links to ODP drilling on the New Jersey margin. Fulthorpe participated in both Leg 150 (slope) and Leg 174A (shelf) and Austin was co-chief scientist of Leg 174A.

REFERENCES

- Austin, J.A., Jr., Fulthorpe, C.S., Mountain, G.S., Orange, D., and Field, M.E., 1996, Continental margin seismic stratigraphy: assessing the preservation potential of heterogeneous geologic processes operating on continental shelves and slopes, *Oceanography*, v. 9, p. 173-177
- Austin, J.A., Jr., Steckler, M.S., Goff, J., Field, M.E., Fulthorpe, C.S., Mayer, L.A., Mountain, G.S., Niedoroda, A.W., Orange, D., and Swift, D.J.P., 1997, Continental terraces hold clues to the generation and preservation of the stratigraphic record, *Eos*, v. 78, p. 1-2.
- Fulthorpe, C.S., and Austin, J.A., Jr., in press, The anatomy of rapid margin progradation: three-dimensional geometries of Miocene clinoforms offshore New Jersey, *AAPG Bulletin*.
- Haq, B.U., J. Hardenbol, and P.R. Vail, 1987, Chronology of fluctuating sea levels since the Triassic, *Science*, v. 235, p. 1156-1167.

WEB PAGE: <http://www.ig.utexas.edu/>




Figure 1. Bathymetric map showing locations of commercial seismic grid, including interpreted area of mapped Miocene clinoforms in the subsurface. Also shown are: a high-resolution MCS survey grid collected in 1995 for STRATAFORM, and locations of existing ODP Middle Atlantic Transect (MAT) sites drilled on Legs 150 and 174A, plus DSDP Site 612 (Leg 95). The COST B-2 well is approximately coincident with Site 1071.

m1c
Middle Miocene
(~11.0 Ma)





Figure 2. Buried clinoform m1c (middle Miocene, ~11.0 Ma). Lower: structure map showing seismic grid (Figure 1), existing drill sites 1071, 1072 (ODP Leg 174A), 902-904, 906 (ODP Leg 150), 612 (DSDP Leg 95), and additional proposed MAT drill sites. Units are milliseconds two-way travelt ime below present sea level. Upper panel: 3-D perspective shaded image with travelt ime contours (azimuth of artificial illumination = 220°). Both panels are viewed from an azimuth of 180° and an elevation of 30°. The rollover, or clinoform breakpoint, bifurcates in the southern part of the mapped area. Clinoform slope canyons are absent, but gullies may be present at the limits of seismic resolution.