

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 JUN 2007		2. REPORT TYPE N/A		3. DATES COVERED	
4. TITLE AND SUBTITLE Things that go boom: injuries from explosives				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Eastridge B. J. /				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) United States Army Institute of Surgical Research, JBSA Fort Sam Houston, TX 78234				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 1	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Things That Go Boom: Injuries From Explosives

COL Brian J. Eastridge, MC, USA

J Trauma. 2007;62:S38.

Explosive agents are materials that undergo rapid exothermic reaction when appropriately stimulated. The degree to which this reaction occurs is dependent upon the characteristics of the explosive agent. Low order explosives react by rapid burning or conflagration. On the other hand, high order explosives produce extreme heat and energy and result in the formation of a pressure wave or “blast wave.” This supersonic, superheated wave, known as “overpressure” comprises the blast front (Fig. 1). The blast wave is reflected and sustained by fixed structures and confined environments such as rooms, vehicles, etc., and may portend the effects of blast-related injury. By the same mechanism, water, which is a relatively noncompressible medium, sustains more of the energy from the blast energy and as such blast waves in water have a greater injurious effect propagated over a greater distance. The three main factors which characterize the blast wave are peak rate of pressure rise, peak pressure, and duration of the pressure rise.

Injuries from explosive agents are characterized by mechanism of injury associated with the explosive event. Primary blast injuries are those injuries associated with the blast wave itself. It should be noted that by this definition, true blast injury is only a subset of all injuries caused by explosions. The mechanism of injury in blast is the impartation of heat and energy to the body, particularly, air filled organs, by the blast wave itself. Survival and injury from primary blast injury is contingent upon a number of factors including energy of the blast, confined versus open space and standoff distance from the explosive source. Survivable blast injuries are characterized by tympanic membrane rupture, pulmonary barotraumas, bowel contusion and perforation, and concussive brain injury. The secondary explosive injuries




Fig. 1. The blast front.




Fig. 2. Standoff distance from the explosive source.

are fragmentation injuries from the casing of the explosive or secondary debris. Most are penetrating injuries and often are associated with multiple penetrations of the victim. Tertiary explosive injuries are those injuries caused by physical displacement of the victim, usually as a result of the attenuated energy from the blast wave. Lastly, quaternary effects are burns, inhalations and exacerbation of underlying conditions.

Copyright © 2007 by Lippincott Williams & Wilkins, Inc.

This article was written for the proceedings from a conference entitled *12th Annual San Antonio Trauma Symposium* in San Antonio, Texas. The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Brooke Army Medical Center, US Army Institute of Surgical Research and The University of Texas Health Science Center at San Antonio.

DOI: 10.1097/TA.0b013e318065414f