

MODELING SEDIMENTATION ON THE CONTINENTAL MARGIN

Donald J. P. Swift, and Alan Wm. Niedoroda* (adjunct),
Department of Oceanography, Old Dominion University, Norfolk, VA, 23908:
Ph (757) 683-4937; Fax -5303; dswift@odu.edu; Award No. N0014-95-1-0202

LONG TERM GOAL

We propose to develop a 3-level system of numerical models.

OBJECTIVES

We propose to build a boundary layer sedimentation model (**Resuspension-Bioturbation Model** (level 1), incorporate the Resuspension-Bioturbation Model as the 'driving element' in a 2-D, cross-shelf **Facies Model** (level 2), and Develop a time-averaged, cross-margin sediment transport model in which the morphodynamics of the continental margin surface is reproduced (**Sequence Stratigraphic model** (level 3).

APPROACH

We plan to calibrate the resuspension model against the tripod data sets that have been collected in the course of the Strataform program. The Facies model must be probabilistic in design, in order to develop realizations of the storm and flood climates. Modeling at the stratigraphic scale will require control of morphodynamics and lithospheric mechanics.

WORK COMPLETED

We (Yong Zhang, ODU) have developed a simple 1-D resuspension model suitable for driving the 2-D facies model. We (Alan Niedoroda and Chris Reed, WCI) have also developed a more sophisticated 1-D resuspension-bioturbation model that fully describes boundary layer resuspensive behavior, and computes the effects of bioturbation on the adjacent sea floor. We have an operational 2-D facies model, developed by Yong Zhang (ODU) that computes storm and flood bed successions the shelf floor. We have an operational version of our stratigraphic model that combines both the LDGO Geophysical routines and The WCI sediment transport algorithm.

* Woodward Clyde International
376 Hartfield Rd.
Tallahassee FL 32303

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Modeling Sedimentation on the Continental Margin				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Old Dominion University, Department of Oceanography, Norfolk, VA, 23529				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

Our Level 1 Models show a “lidding” effect in the combined flow boundary layer occurs during major storms and may have a major effect on sediment transport.

Results from the 2-D facies model (Level 2) show that the mud facies tract begins much closer to shore than it would if coastal erosion, not river flooding, were the primary source of sediment. It also shows that optimum storm bed thickness occurs on the seventy meter isobath of the Northern California Shelf. The model indicates that while most beds on the Northern California Study area are of more or less recent flood provenance, most of their material has undergone multiple resuspensions, and is storm packaged. Their sediment is in various stages of textural maturity, depending on the number of resuspensions to which it has been subjected. Muddy beds have been subjected to less storm reworking than sandy beds, but their thickness distributions are controlled by wave height, not storm intensity. Our stratigraphic model (level 3) demonstrates that continental shelf surfaces are surfaces of equilibrium, and that clinoform shelf structure reflect attempts by the shelf surface to regain equilibrium after conditions have changed rapidly.

IMPACT AND APPLICATIONS

The facies model described above will predict the geotechnical and acoustic properties of first 10 meters of the sea floor. The Stratigraphic model will predict seafloor structure at depths up to several kilometers, for foundation studies and petroleum exploration.

TRANSITIONS

We are, in structural terms, the most “downstream” component of Strataform in the sense that we use the results of other Strataform groups, but the consumers of our group will be will consist of the larger world outside of Strataform.

RELATED PROJECTS

We are calibrating the models against dynamical data sets collected by Cacchione, Sternberg, and Wright. We are validating the models by comparison with the sea floor observations of Drake, Wheatcroft, Borgeld, and Nittrouer. We are cooperating closely with ongoing Strataform modeling efforts by Syvitski and students (IAAR), and at Steckler at LDEO. We plan also to Collaborate with Wiberg (UVA).

PUBLICATIONS

Niedoroda, A.W., Reed, C.W., Swift, D.J.P., Arato, A. and K. Hoyanagi, 1995. Modeling shore-normal large scale coastal evolution. *Marine Geology*, v. 126, p. 181-199.

Steckler, M.S, Swift, D.J.P., Syvitski, J.P., Goff, J.A. and Niedoroda, A.W., 1996, Modeling the sedimentology and stratigraphy of continental margins, *Oceanography*, v. 9, p183-188

Syvitski, J., L. Pratson. P.L. De Boer, M. Garcia, M. Steckler., H. Lee, M. Perlmutter, G. Parker, P. Wiberg, and D. Swift, 1997. Earthworks: a large scale and complex numerical model to understand the flux and deposition of sediment over various time scales. in Pawlowsky-Glahn, *Proceedings of the Third Annual Conference of the international*

Association of mathematical Geologists. International Center for Numerical Methods, Universitat Politècnica de Catalunya, 08034, Barcelona, Spain

Zhang, Y., Swift, D. J. P., Niedoroda, A. W., Reid, C. W., and Thorne, J. A., 1997. Simulation of sedimentary facies on the northern California Shelf: implications for an analytical theory of facies differentiation. *Geology* v. 27, p. 635-638.

Carey, J. S. , D. J. P. Swift, M. Steckler, C. W. Reed and A. W. Niedoroda, In Press. High resolution sequence stratigraphy modeling: effects of sedimentation processes. *Numerical Experiments in Stratigraphy. Soc Econ, Paleon. Mineral. Spec. Publ.*

Reed, C.W., A. W. Niedoroda and D.J. P., Swift, In Press. Sediment entrainment and transport limited by

bed load armoring. Marine Geology.

Reed, C.W., A. W. Niedoroda, D. J. P. Swift, Y. Zhang, J. Carey, and M.S. Steckler, in press. Coastal evolution: modeling fabric and form at successive spatial scales. 1997 Coastal Dynamics congress, University of Plymouth, Plymouth, U.K.

Swift D.J. P., J. S. Carey, Y. Zhang, C. W Reed, A. W. Niedoroda, and M. Steckler, in press. Quaternary stratigraphy and sedimentation on the Northern California Continental Margins constrained by numerical modeling: preliminary results. Marine Geology.

Zhang, Y., D.J. P. Swift, C. R. Reed, A. Wm. Niedoroda and S. Fan, In Press. Two-dimensional numerical modeling of storm beds for sedimentary facies simulation. Marine Geology.

Zhang, Y., Swift, D.J.p., Yu, Z., and Jin, L., in press. Modeling of coastal profile evolution on the abandoned delta of the Huanghe River. Marine Geology