

Seabed Variability and its Influence on Acoustic Prediction Uncertainty

John A. Goff

University of Texas Institute for Geophysics

4412 Spicewood Springs Rd., Bldg. 600,

Austin, TX 78759

phone: (512) 471-0476; fax: (512) 471-0999; email: goff@ig.utexas.edu

Award Number: N00014-01-1-0891

<http://www.ig.utexas.edu/research/projects/uncertainty/uncertainty.htm>

LONG-TERM GOALS

The seabed variability group was incorporated into the ONR Defense Research Initiative on Capturing Acoustic Uncertainty in order to provide a focused investigation into the contribution of seabed inhomogeneity (both bathymetric and geoacoustic variability) to uncertainty in acoustic prediction and detection. My effort is focused on characterizing such inhomogeneity with existing data, with the goal of providing useful input to the acoustic modeling efforts.

OBJECTIVES

The primary scientific objective of my initial work for this project is to formulate a statistical model for bathymetric and geoacoustic variability for the seabed, and to generate from that model realizations that can be incorporated into acoustic modeling efforts.

APPROACH

In the first two years of the program, our seabed characterization efforts are focused on analysis of data collected within the ONR STRATAFORM/Geoclutter natural laboratory on the New Jersey shelf (Figure 1). A number of data sets are available: (1) swath multibeam bathymetry and sidescan data (collected by Larry Mayer as part of the STRATAFORM program; (2) chirp seismic reflection and geoacoustic estimation (data collected and being processed by Steve Schock as part of the Geoclutter program); (3) grab samples and short cores (collected by John Goff and Chris Sommerfield for Geoclutter); and (4) *in situ* velocity measurements (collected by Larry Mayer for Geoclutter).

The swath sidescan data are of particular interest to us because, in the very low slope environment of the continental shelf, backscatter intensity is primarily a reflection of sedimentological properties. Our first task was to investigate the correlation between backscatter intensity and the sediment properties measured (primarily grain size distribution and compressional velocity), in the hope that the highly detailed sidescan map could be used as a proxy for these important sediment properties. The grain size distribution of the outer New Jersey shelf is quite complex, often composed of identifiably separate contributions of mud, sand, and coarse content (either shell hash or gravel). Standard parameterizations of the distribution (i.e., mean, rms, skewness and kurtosis) are inadequate representations of the true nature of the sediments. Rather, we formulated the following parameters to characterize our samples: fine fraction (< 0.063 mm), coarse fraction (> 4 mm), and mean sand grain size (i.e., for grain sizes from 0.063 mm to 2 mm).

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2002		2. REPORT TYPE		3. DATES COVERED 00-00-2002 to 00-00-2002	
4. TITLE AND SUBTITLE Seabed Variability and its Influence on Acoustic Prediction Uncertainty				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Texas Institute for Geophysics,,4412 Spicewood Springs Rd., Bldg. 600,,Austin,,TX, 78759				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The seabed variability group was incorporated into the ONR Defense Research Initiative on Capturing Acoustic Uncertainty in order to provide a focused investigation into the contribution of seabed inhomogeneity (both bathymetric and geoacoustic variability) to uncertainty in acoustic prediction and detection. My effort is focused on characterizing such inhomogeneity with existing data, with the goal of providing useful input to the acoustic modeling efforts.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Our second priority was to statistically characterize bathymetry and sediment properties, either directly using available data or through the sidescan data as proxy. Data are statistically analyzed by fitting parameterized models to empirical representations of either the power spectrum, covariance function, or semi-variogram (e.g., Goff and Jordan, 1988). Once a statistical model is formulated, any number of independent random realizations can be generated, either unconditioned or conditioned to existing data constraints, using recognized Fourier methodologies (e.g., Goff and Jennings, 1999).

WORK COMPLETED

Measured seafloor properties in the STRATAFORM/Geoclutter natural laboratory have been compared against collocated backscatter data to investigate the possible use of backscatter data as a proxy for measuring spatial variability in sedimentary properties. Statistical characterizations have been produced for bathymetry, velocity and grain size properties. Random realizations have been generated for bathymetry and velocity for use in acoustic modeling work by collaborators in the seabed variability group (Kevin LePage and Bob Odom).

RESULTS

In the correlation analysis of sidescan backscatter and sedimentary properties, we found the following: (1) compressional velocity is primarily correlated to mean sand grain size and fine fraction; (2) backscatter intensity is correlated primarily to coarse fraction; and (3) compressional velocity and backscatter can be correlated only where coarse material is not a factor. Hence, the coarse material (shell hash, gravel) tends to dominate the backscatter when it is present in significant quantities (> ~5%). Unfortunately, this is true over much of the New Jersey outer shelf, and the backscatter map we observe (Figure 1) is largely a map of coarse material content. In contrast, acoustic velocity is not significantly affected by the coarse content, and the sidescan backscatter map cannot be used as a proxy for velocity. Nevertheless, the *in situ* velocity measurements are numerous and with separation distances that cover a range of scales – from 100's of meters to 10's of kilometers. This sort of distributed sampling is well suited to semi-variogram analysis using binning procedures (Goff et al., 2002). A von Kármán model function (Goff and Jordan, 1988) is fit to the velocity semi-variogram in Figure 3, and a 2-D realization is presented in Figure 4.

IMPACT/APPLICATIONS

Realizations of bathymetry and velocity statistical models are being used by LePage and Odom for use in acoustic modeling of the impact of seabed variability on acoustic prediction..

TRANSITIONS

It is hoped that others acoustic modelers within the Uncertainty DRI will also be able to make use of the seabed realizations.

RELATED PROJECTS

The ONR STRATAFORM and Geoclutter programs are providing critical data inputs for this project.

REFERENCES

Goff, J. A., and J. W. Jennings, Jr., Improvement of Fourier-based unconditional and conditional simulations for band limited fractal (von Kármán) statistical models, *Math. Geol.*, 31, 627-649, 1999.

Goff, J. A., and T. H. Jordan, Stochastic Modeling of Seafloor Morphology: Inversion of Sea Beam data for second-order statistics, *J. Geophys. Res.*, 93, 13,589-13,608, 1988.

Goff, J. A., Wheatcroft, R. A., Lee, H., Drake, D. E., Swift, D. J. P., and Fan, S., Spatial Variability of Shelf Sediments in the STRATAFORM Natural Laboratory, Northern California, *Cont. Shelf Res.*, 22, 1199-1223, 2002




Figure 1. *Sidescan backscatter map over the ONR STRATAFORM/Geoclutter natural laboratory, outer New Jersey shelf (higher backscatter = lighter shades), with locations of colocated grab samples and velocity measurements.*




Figure 2. Examples of the correlation analysis conducted for the New Jersey grab sample sediment distribution (mean sand grain size – top; weight % fines – middle, and weight % coarse – bottom), velocity measurements (right panels) and backscatter intensity (left panels). The coarse % vs. backscatter values indicate a non linear trend.




Figure 3. Semi-variogram (green) for velocity measurements conducted within the STRATAFORM/Geoclutter natural lab, with von Kármán statistical model overlain (blue dashed); H^2 is the variance, λ_0 is the characteristic scale, and D is the fractal dimension.




Figure 4. Realization of the von Kármán statistical model for seafloor velocity presented in Figure 3.