

Scattering of Internal Gravity Waves at Finite Topography

Peter Muller
University of Hawaii
Department of Oceanography
1000 Pope Road, MSB 429
Honolulu, HI 96822
phone: (808)956-8081 fax: (808)956-9164 email: pmuller@soest.hawaii.edu
Award Number: N00014-96-1-0489

LONG-TERM GOAL

Description and modeling of the kinematic structure and the dynamical processes of oceanic internal gravity waves.

Understanding the role of internal gravity waves in the redistribution of momentum, potential vorticity, heat, and salt.

Development of a numerical model that predicts the internal wave field and the internal wave induced transports globally and regionally.

SCIENTIFIC OBJECTIVES

The current research project investigates the interaction of internal gravity waves with topography. Its specific objectives are:

- to assess the role of reflection and scattering at topography in the redistribution of energy within the internal wave band and in providing energy for boundary mixing.
- to determine the modifications of the deep ocean internal wave field when it propagates up the continental slope onto the shelf and to compare this remotely forced wave field with the locally generated field.

APPROACH

The scientific approach combines theoretical analysis with numerical modeling. The specific approach differs for scattering in two and three dimensions.

In two dimensions the method of characteristics can be employed. A general theoretical framework was developed by Baines (1971 a, b) and by Sandstrom (1976). The major challenges are the implementation of the radiation condition and the singularities at points where the topographic slope equals the wave slope.

In three dimensions, no general approach has been developed as yet. The governing equation is a hyperbolic equation which is analogous to the ordinary wave equation with the vertical coordinate replacing the time coordinate. This difference causes difficulties in

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Scattering of Internal Gravity Waves at Finite Topography				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Hawaii, Department of Oceanography, 1000 Pope Rd, Honolulu, HI, 96822				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

solving this equation since the vertical space coordinate does not increase uniformly along the ray path in the bottom scattering problem.

WORK COMPLETED

In the two dimensional case we generalized Baines' (1971 a,b) theoretical results for an infinitely deep ocean to a finite-depth ocean. For applications a numerical model was developed for two types of topographic configurations: (a) a slope-shelf configuration with only one open sides and (b) a ridge configuration with two open sides. The model allows to study the scattering of an incoming internal wave at a smooth but otherwise arbitrary topography. Specifically, a suite of experiments is being conducted to study the scattering of an incoming deep-ocean internal wave field with a Garrett and Munk spectrum. The spectra of both the forward- and back-scattered internal wave field are calculated. From these spectra the redistribution of the wave energy in modenumber space is determined. If the energy flux is redistributed from low to high modenumbers a critical modenumber is determined, such that the Richardson number of all waves with modenumbers smaller than the critical modenumber equals the critical Richardson number $1/4$. The energy flux to modenumbers larger than the critical modenumber excites waves that are likely to break and cause mixing. The energy flux past this critical modenumber is thus interpreted as being available for internal wave induced boundary mixing.

In three dimensions, we have developed an approach based on the Green's function of the governing wave equation. Following Hurley (1972) this Green's function is obtained by analytic continuation of the Green's function of the elliptic problem into the hyperbolic range. With this Green's function the scattering problem can be cast in to an integral equation for the topographic source of the scattered wave field and the scattered wave field itself. The approach correctly reproduces the scattering at a straight slope.

RESULTS

The numerical model runs for the two-dimensional case indicate that internal waves interact in a complex manner with topography. The picture that is slowly emerging from the model runs is that the redistribution of the energy flux does not only depend on the height and slope of the topography but also on its curvature, as has been anticipated by Gilbert and Garrett (1989). Results obtained from the use of piecewise linear topography should thus be used with caution.

IMPACT/APPLICATION

The research will help to clarify the role of open ocean internal for (a) boundary mixing, and (b) for processes on the continental slope and shelf.

TRANSITIONS

RELATED PROJECTS

REFERENCES

Baines, P. G., 1971: The reflexion of internal/inertial waves from bumpy surfaces. *J. Fluid Mech.*, **46**, 273-291.

Baines, P. G., 1971: The reflexion of internal/inertial waves from bumpy surfaces. Part 2. Split reflexion and diffraction. *J. Fluid Mech.*, **49**, 113-131.

Gilbert, D. and C. Garrett, 1989: Implications for ocean mixing of internal wave scattering off irregular topography. *J. Phys. Oceanogr.*, **19**, 1716-1729.

Hurley, D. G., 1972: A general method for solving steady-state internal gravity wave problems. *J. Fluid Mech.*, **56**, 721-740.

Sandstrom, H., 1976: On topographic generation and coupling of internal waves. *Geophys. Fluid Dyn.*, **7**, 231-270.

Award Number: N00014-96-1-0489

Peter Muller
University of Hawaii
Department of Oceanography
1000 Pope Road, MSB 429
Honolulu, HI 96822
email: pmuller@soest.hawaii.edu
phone: (808)956-8081
fax: (808)956-9164

<total_students> 1
<total_women_students> none
<total_minority_students> none
<total_postdocs> none
<total_women_postdocs> none
<total_minority_postdocs> none
<patents_list> none
<patents_number> 0
<pubs_list> Firing, E., R. C. Lien and P. Müller, 1997: Observations of strong inertial oscillations
after the passage of tropical cyclone Ofa. *J. Geophys. Res.*, **103**, 3317-3322.
<pubs_number> 1
<presentations> none
<service> none
<honors> none
<other_orgs> none